

# East Anglia TWO Offshore Windfarm

# Chapter 12 Ornithology

Preliminary Environmental Information Volume 1 Document Reference: EA2-DEVWF-ENV-REP-IBR-000807

Prepared by:	Checked by:	Approved by:



Revis	sion Summary				
Rev	Date	Document Status	Prepared by	Checked by	Approved by
01	11/01/2019	For issue	Paolo Pizzolla	Julia Bolton	Helen Walker

Description of Revisions			
Rev	Page	Section	Description
01	n/a	n/a	Final Draft



## **Table of Contents**

12	Ornithology	1
12.1	Introduction	1
12.2	Consultation	1
12.3	Scope	8
12.4	Assessment Methodology	15
12.5	Existing Environment	24
12.6	Potential Impacts	34
12.7	Cumulative Impacts	105
12.8	Transboundary Impacts	153
12.9	Interactions	154
12.10	Inter-relationships	155
12.11	Summary	156
12.12	References	161



## The *Chapter 12 Ornithology* figure is presented in **Volume 2** and listed in the table below.

Figure number	Title
12.1	Ornithology Survey Area

## The *Chapter 12 Ornithology* appendix is presented in *Volume 3* and listed in the table below.

Appendix number	Title
12.1	Baseline Offshore Ornithology Technical Report



## Glossary of Acronyms

APEM	APEM is an environmental consultancy with specialist expertise in digital aerial
	survey
AR	Avoidance Rates
BDMPS	Biologically Defined Minimum Population Scale/size
BEIS	Business Environment and Industrial Strategy
BoCC	Birds of Conservation Concern
BTO	British Trust for Ornithology
CAA	Civil Aviation Authority
CIA	Cumulative Impact Assessment
CRM	Collision Risk Modelling
EA2	East Anglia TWO
EA3	East Anglia THREE
EATL	East Anglia THREE Limited
EC	European Commission
EIA	Environmental Impact Assessment
EMF	Electro-magnetic Field
EPP	Evidence Plan Process
ES	Environmental Statement
ESAS	European Seabirds at Sea database
ETG	Expert Topic Group
EU	European Union
FAME	Future of the Atlantic Marine Environment
GGOWL	Greater Gabbard Offshore Wind Farm Limited
GPS	Global Positioning System
HRA	Habitats Regulations Assessment
ICES	International Council for the Exploration of the Sea
IEEM	Institute of Ecology and Environmental Management
JNCC	Joint Nature Conservation Committee
KDE	Kernel Density Estimate
MAGIC	Multi-Agency Geographic Information for the Countryside
MCA	Maritime and Coastguard Agency
MLWS	Mean Low Water Springs
MMO	Marine Management Organisation
MRSea	A spatial modelling software package
MS	Method Statement
MW	Megawatt
NAF	Nocturnal Activity Factor
NE	Natural England
NGO	Non-Governmental Organisation
NPPF	National Planning Policy Framework
NPS	National Policy Statement
ORJIP	Offshore Renewables Joint Industry Programme
OWEZ	Offshore Wind Farm Egmond aan Zee, Netherlands
OWF	Offshore Windfarm
PAWP	Princess Amalia Wind Park, Netherlands
PBR	Potential Biological Removal



PCH	Potential Collision Height
PEI or PEIR	Preliminary Environmental Information Report
PVA	Population Viability Analysis
RSPB	Royal Society for the Protection of Birds
SAC	Special Area of Conservation
SAR	Search and Rescue
SE	Standard error (of the mean)
SNCB	Statutory Nature Conservation Body
SNH	Scottish Natural Heritage
SOSS	Strategic Ornithological Support Services
SPA	Special Protection Area (note, pSPA indicates a proposed site not yet designated)
SSSI	Site of Special Scientific Interest
UK	United Kingdom
WWT	Wildfowl and Wetlands Trust
ZAP	Zonal Appraisal and Planning
ZEA	Zonal Environmental Appraisal



## Glossary of Terminology

Applicant	East Anglia TWO Limited
As built	A term used for offshore windfarm developments that are operational and
	where the turbine array 'as built' is different to the worst case scenario in the
	Environmental Impact Assessment for the development (for example where a
	windfarm is built out with fewer turbines than the consented design envelope).
Construction	A fixed offshore structure required for construction, operation, and
operation and	maintenance personnel and activities.
maintenance	
platform	
East Anglia TWO	The proposed project consisting of up to 75 wind turbines, up to four offshore
project	electrical platforms, up to one construction operation and maintenance
	platform, inter-array cables, platform link cables, up to one operational
	meteorological mast, up to two offshore export cables, fibre optic cables,
	landfall infrastructure, onshore cables and ducts, onshore substation, and
East Anglia TMO	The red line boundary in which all wind turbines and ancillary infrastructure will
	he leasted
East Anglia Zono	The broader area defined for Pound 3 applications within which the East Applications
	TWO windfarm site is located together with East Anglia One. East Anglia
	THREE Norfolk Boreas and Norfolk Vanguard
European site	Sites designated for nature conservation under the Habitats Directive and Birds
Laropourrono	Directive, as defined in regulation 8 of the Conservation of Habitats and
	Species Regulations 2017 and regulation 18 of the Conservation of Offshore
	Marine Habitats and Species Regulations 2017. These include candidate
	Special Areas of Conservation, Sites of Community Importance, Special Areas
	of Conservation and Special Protection Areas.
Evidence Plan	A voluntary consultation process with specialist stakeholders to agree the
Process	approach to the EIA and information to support HRA.
Inter-array cables	Offshore cables which link the wind turbines to each other and the offshore
	electrical platforms, these cables will include fibre optic cables.
Landfall	The area (from Mean Low Water Springs) where the offshore export cables
	would make contact with land, and connect to the onshore cables.
Meteorological mast	An offshore structure which contains metrological instruments used for wind data acquisition.
Migration free	The breeding season for migratory seabird species is defined as a wider
breeding season	breeding season and a narrower window known as the migration free breeding
	season. In a given species, the timing of breeding will vary depending on the
	location of the breeding area; with the start of breeding usually later in more
	northerly locations. Thus, while birds at some colonies are beginning to nest,
	others may still be migrating to breeding sites. A core or migration free
	breeding season is defined as the period when all or the majority of breeding
	adults of a given species are present at breeding colonies.
Natura 2000 site	A site forming part of the network of sites made up of Special Areas of
	Habitats Directive and Birds Directive
Offshore	The East Anglia TWO windfarm site and offshore cable corridor (up to Mean
development area	High Water Springs).
	o -1 o-7



	The transmission assets required to export generated electricity to shore. This	
Offshore electrical	includes inter-array cables from the wind turbines to the offshore electrical	
infrastructure	platforms, offshore electrical platforms, platform link cables and export cables	
	from the offshore electrical platforms to the landfall.	
Offshore electrical	A fixed structure located within the windfarm area, containing electrical	
platform	equipment to aggregate the power from the wind turbines and convert it into a	
	more suitable form for export to shore.	
Offshore export	The cables which would bring electricity from the offshore electrical platforms	
cables	to the landfall, these cables will include fibre optic cables.	
Offshore	All of the offehere infractructure including wind turbings, platforms, and apples	
infrastructure	All of the offshole initiastructure including wind turbines, platforms, and cables.	
Offshore platform	A collective term for the construction operation and maintenance platform and	
	the offshore electrical platforms.	
Platform link cables	An electrical cable which links one or more offshore platforms, this will include	
	fibre optic cables.	
Safety zones	A marine area declared for the purposes of safety around a renewable energy	
	installation or works / construction area under the Energy Act 2004.	
Scour protection	Protective materials to avoid sediment being eroded away from the base of the	
	foundations as a result of the flow of water.	



## **12Ornithology**

## 12.1 Introduction

- This chapter is an assessment of the potential impacts that may arise from the construction, operation and decommissioning of the offshore components of the proposed East Anglia TWO project. It has been prepared by Royal HaskoningDHV from baseline survey work and data processing undertaken by APEM Ltd and data analyses by MacArthur Green.
- 2. The chapter describes the offshore components of the proposed project in relation to ornithology; the consultation that has been held with stakeholders; the scope and methodology of the assessment; the avoidance and mitigation measures that have been embedded through project design; the baseline data on birds and important sites and habitats for birds acquired through desk study and surveys and assesses the potential impacts on birds.
- 3. An ornithological assessment of the export cable landfall and onshore components of the project is included in *Chapter 23 Onshore Ornithology*.
- 4. Full details of the baseline data for the offshore ornithology assessment, acquired through the surveys specifically carried out within the East Anglia TWO windfarm site and a 4km buffer can be found in *Appendix 12.1 Baseline Offshore Ornithology Technical Report*.

## 12.2Consultation

- 5. Consultation is a key driver of the Development Consent Order (DCO) application process. To date, consultation regarding offshore ornithology has been conducted through formal submission of the Scoping Report submitted in November 2017 (SPR 2017) and through Expert Topic Group (ETG) Meetings in April 2017 and March 2018 involving Natural England and the Royal Society for the Protection of Birds (RSPB) as detailed in *section 5.3.3* of *Chapter 5 EIA Methodology*. Feedback received through this process has been considered in preparing incorporated into the PEIR where appropriate and this chapter will be updated following the next stage of consultation for the final assessment submitted with the Development Consent Order (DCO) application.
- 6. Responses from stakeholders have been captured in *Table 12.1* below and a reference included to where responses are addressed within this Preliminary Environmental Information Report (PEIR).
- 7. Correspondence received in relation to transboundary consultations is summarised in *section 12.8* below.



8. Further consultation will continue to be undertaken once the PEIR is made available and during further ETG meetings conducted between PEIR submission and the DCO application submission.

Consultee	Date / Document	Comment	Response / where addressed in the PEI
Natural England	08/12/2017 Scoping Response	NE maintains that a seasonal restriction is put in place from Nov – Feb for cable installation in order to mitigate against impacts to red- throated diver. This species has been particularly affected and displaced from large areas within the Outer Thames Estuary due to OWF construction. To reduce impacts further it would be a sensible option to cease works/activities that interact with the designated sites during this period.	A seasonal restriction has been considered further as part of the assessment, but is not considered to be necessary given the small predicted impact of disturbance on red-throated diver (see <b>section</b> <b>12.6.1.1.1</b> ). A best practice protocol for minimising disturbance to red-throated divers will be adopted as for East Anglia THREE.
Natural England	08/12/2017 Scoping Response	It appears that the Birds of Conservation Concern (BoCC) listing from BoCC 3 (Eaton et al. 2009) has been used. This listing has since been updated by BoCC 4, we advise the Applicant to see Eaton et al. (2015), available online at: <u>http://britishbirds.co.uk/wp- content/uploads/2014/07/BoCC4.pdf</u>	Updated to BoCC 4 for this assessment, see <i>Table 12.9</i> .
Natural England	08/12/2017 Scoping Response	We note the comments on the need for mitigation will be to some extent dependent on the results of site specific survey and the impact assessment. However, Natural England's advice at the EA3 hearing was that adverse effect on site integrity cannot be excluded in- combination with other plans or projects in respect of predicted mortality from collision on kittiwake from Flamborough Head and Bempton Cliff SPA and Flamborough and Filey Coast pSPA. Therefore Natural England would welcome any mitigation measures, such as raising the minimum hub height to be considered at the earliest	EA2 will review impact magnitudes as they become apparent and consider options for mitigation as they arise. Further work has been undertaken to refine realistic cumulative turbine numbers. This includes adjustments to reflect new information on nocturnal activity levels, and reference to revised collision risk figures for developments that are operational and where the turbine array 'as built' has a lower collision risk than the worst case scenario in the Environmental Impact Assessment (see assessment of cumulative impacts of collision in <i>section</i> 12.7.4).

### Table 12.1 Consultation Responses

## East Anglia TWO Offshore Windfarm Preliminary Environmental Information Report



Consultee	Date / Document	Comment	Response / where addressed in the PEI
		opportunity.	
Natural England	08/12/2017 Scoping Response	We agree with the use of Furness (2015) for use of definitions of biological seasons. However, further consideration will need to be given to lesser black-backed gull as the breeding season for individuals breeding at Alde Ore Estuary SPA will be wider than the May- July period stated as the breeding season only period, given that the project is within the foraging range of Alde-Ore Estuary SPA We suggest that this is discussed and agreed during Evidence Plan Process.	The breeding season collision risk estimate for lesser black-backed gull is based on the full breeding season (April to August; Furness, 2015) as opposed to the migration- free (core) breeding period. Although the project is within foraging range of lesser black- backed gulls breeding at the Alde Ore Estuary, tracking data from the SPA colony indicate that the proposed project is outwith the core foraging areas for this species during the breeding season ( <b>section 12.6.2.3</b> ). It is therefore considered unlikely that breeding birds from the SPA make regular use of the project.
Natural England	08/12/2017 Scoping Response	We note that surveys are planned between May to August 2018 to ensure there are 24 months of site- specific data available for assessment. We welcome the commitment to collect 24 months of site specific data at the EA2 windfarm site. We also acknowledge that additional contextual information will come from surveys undertaken for the former East Anglia Zone and the former East Anglia TWO windfarm site.	Noted.
Natural England	08/12/2017 Scoping Response	In addition to the RSPB tagging studies from Flamborough, there is tracking data of lesser black-backed gulls from Alde-Ore Estuary SPA from the Department for Business Environment and Industrial Strategy (BEIS) funded BTO study, and there is further tracking planned as part of Galloper's post construction monitoring which may be available during the examination.	Reference has been made to tracking data from the Alde-Ore Estuary SPA in the assessment (see <i>section 12.6.2.3.1.3</i> ). Where available we will attempt to obtain any additional relevant data for the assessment.



Consultee	Date / Document	Comment	Response / where addressed in the PEI
Natural England	08/12/2017 Scoping Response	We would like to clarify if it is planned to use MRSea on all the survey data, or whether reliable model based estimates require a minimum number of observations, and therefore may only be used for the more numerous species.	MRSea has not be used as there are limited data for many species. Design based outputs are however provided for all species.
Natural England	08/12/2017 Scoping Response	We agree that the species assessed will depend on the results of the surveys but will include: fulmar, gannet, kittiwake, lesser black- backed gull, great black-backed gull, herring gull, red-throated diver, guillemot, razorbill and puffin. We assume that other species assessed may include those that may pass through on migration but are only recorded in small numbers by snap shot aerial surveys, for example little gull. It is not clear in the Scoping Report (SPR 2017) if non-seabird migrants are also being considered.	An evidence plan supporting document on non-seabird migrants has been agreed based on nearby windfarm assessments, to justify scoping out non-seabird migrants, as these will have extremely low predicted impacts.
Natural England	08/12/2017 Scoping Response	We agree with the likely key issues listed in the Scoping Report (SPR 2017), although we would include lesser black-backed gull collision risk during the breeding season, in addition to the non-breeding season.	Noted and agreed, collision risk during the breeding season has been assessed for the project alone ( <i>section 12.6.2.3</i> ) and cumulatively ( <i>section 12.7.4.2</i> ).
Natural England	08/12/2017 Scoping Response	We agree with the list of expected features of the HRA however we recommend that impacts on other qualifying features of Flamborough and Filey Coast pSPA and other qualifying features from the Outer Thames Estuary pSPA are also likely be included in the HRA.	We will screen for other impacts in the HRA as suggested. See the HRA Screening Report
Natural England	08/12/2017 Scoping Response	We are content with the proposals for measuring flight height, and would expect there to be enough samples within the site specific surveys to get an adequate sample particularly if the historic digital	BTO generic flight height data have been used for the assessment of collision risk ( <i>section 12.6.2.3</i> ) the aerial survey contractors advised ScottishPower Renewables that the flight height estimates from



Consultee	Date / Document	Comment	Response / where addressed in the PEI
		aerial survey data can be used. We would expect flight heights to be provided with confidence intervals to enable them to be used with a stochastic collision risk model should that be available by the time the application is submitted.	specific baseline survey data were not reliable. Thus, these data have not been used in the assessment.
RSPB	20/12/2017 Scoping Response	Given the recent concerns around potential collision risk to breeding kittiwake and gannet from other windfarms in the former East Anglia Zone, we strongly recommend that the potential for impacts on these species during the breeding season from the proposed East Anglia TWO project is recognised in the table and the subsequent assessment.	Collision risk during the breeding season has been assessed for the project alone ( <i>section 12.6.2.3</i> ) and cumulatively ( <i>section</i> <i>12.7.4.2</i> ).
RSPB	20/12/2017 Scoping Response	If figures for the migration-free breeding season were to be used in the PEI as in the Scoping Report, we consider that it would be necessary to attribute birds in the crossover months to breeding and dispersal in order to ensure collision risk to breeding birds is not underestimated.	With the exception of lesser black- backed gull, the turbine array of the proposed project is considered to be outwith the foraging range of the seabird species considered in the assessment of collision risk. Tracking data from the Alde Ore SPA colony of lesser black-backed gulls also indicate that the proposed project is outwith the core foraging areas for this species during the breeding season ( <i>section 12.6.2.3</i> ).
RSPB	20/12/2017 Scoping Response	We recommend the following paper as a recent critique of the methods used to assess impacts of offshore windfarms on seabird populations: Green, R. E., Langston, R. H., McCluskie, A., Sutherland, R., and Wilson, J. D. (2016). Lack of sound science in assessing windfarm impacts on seabirds. <i>Journal of</i> <i>Applied Ecology.</i>	Full use of available literature and evidence has been made in assessing the project's potential impacts for the PEIR.
The Planning Inspectorate	02/12/2017 Scoping Response	The Inspectorate does not agree that the impact of disturbance due to lighting during operation and decommissioning can be scoped out as no information to support this	A review of the effects of operational lighting has been prepared (Furness, 2018). At the request of NE, construction and operational lighting is considered in



Consultee	Date / Document	Comment	Response / where addressed in the PEI
		approach and no evidence demonstrating clear agreement with relevant statutory bodies has been provided. The PEI should include an assessment of this matter.	the assessment ( <i>sections 12.6.1.1</i> and <i>12.6.2.1</i> ).
The Planning Inspectorate	02/12/2017 Scoping Response	The Inspectorate does not agree that the impact of transboundary impacts can be scoped out as no information to support this approach and no evidence demonstrating clear agreement with relevant statutory bodies has been provided.	Evidence is provided that Transboundary impacts can be screened out of the assessment (see <i>section 12.8</i> and the HRA screening report).
The Planning Inspectorate	02/12/2017 Scoping Response	It is noted that in the Scoping Report no ornithology surveys are proposed to be undertaken along the (cable corridor) AoS, based on conclusions drawn from existing survey information which was used to assess the potential impacts of East Anglia ONE and East Anglia THREE on red- throated diver, and that impacts are expected to be temporary and localised. No other bird species are referenced. The source of the data relied upon to support the conclusions in relation to the proposed East Anglia TWO project should be identified in the PEI and its relevance to bird species other than red- throated diver should be explained. The evidence demonstrating clear agreement with relevant statutory bodies that no surveys are required must be provided.	The methods and evidence (including source of data) for assessing potential impacts along the cable corridor during construction are set out in <b>section</b> <b>12.6.1.1</b> . As for East Anglia ONE and East Anglia THREE the assessment for the export cable corridor considers red-throated diver based on survey data collected for the Outer Thames Estuary SPA.
The Planning Inspectorate	02/12/2017 Scoping Response	Only 2 of 4 European sites are identified in the Method Statement in relation to HRA and are referenced under designated sites in the Scoping Report; Flamborough and Filey Coast pSPA and Alde-Ore Estuary SPA are omitted. In addition, although the little gull is identified in MS paragraph 46 as a feature of the Greater Wash pSPA,	The HRA screening and assessment provides a comprehensive review of potential connectivity of designated sites. This is reflected in the PEIR ( <i>section 12.5.2</i> ). Little gulls were recorded within the East Anglia TWO windfarm site although in small numbers ( <i>Table 12.11</i> ) and were not screened in for



Consultee	Date / Document	Comment	Response / where addressed in the PEI
		it is not included in the list of receptors likely to be affected by the Proposed Development provided in paragraph 44 of the MS. While the information in the ES should not duplicate that in the HRA Report, the Inspectorate expects it to be consistent between the two documents.	assessment for any potential impacts.
The Planning Inspectorate	02/12/2017 Scoping Response	It should be clearly explained in the PEI how the value of a feature will be taken into account in judging its sensitivity and the overall assessment of significance.	This is explained in the impact assessment methodology ( <i>section</i> <i>12.4.3</i> )
Natural England	18/05/2018 Comments on Expert Topic Group (ETG) meeting agreement points	Agree that BTO flight height data and presentation of Option1 and Option 2 outputs is acceptable given the uncertainty around site specific flight height data.	BTO flight height data and Option 1 and 2 outputs used in the assessment of collision risk. Option 1 outputs are provided in the Technical Appendix. See section 12.6.2.3 and Appendix 12.1 Technical Report.
Natural England	18/05/2018 Comments on ETG meeting agreement points	NE have advised SPR that the consequences of lighting for birds during all phases of the project (including construction) should be considered, so any potential impacts and mitigation can be explicitly stated.	A review of the effects of operational lighting has been prepared (Furness 2018). At the request of NE, construction and operational lighting is considered in the assessment ( <i>sections 12.6.1.1</i> and <i>12.6.2.1</i> ).
Natural England	18/05/2018 Comments on ETG meeting agreement points and evidence plan supporting document on non- seabird migrants	Impacts on migrating non-seabirds can be scoped out.	Migrating non-seabird species scoped out and not considered further.

Consultee	Date / Document	Comment	Response / where addressed in the PEI
Natural England	18/05/2018 Comments on ETG meeting agreement points	Agree that transboundary impacts on non-UK ornithology receptors can be scoped out subject to consultation with Scottish Natural Heritage (SNH))	Transboundary effects on non-UK receptors have been scoped out. Awaiting feedback from SNH.

- 9. Ongoing public consultation has been conducted through a series of Public Information Days (PIDs) and Public Meetings. PIDs have been held throughout Suffolk in November 2017, March 2018, and June / July 2018 with further events planned in 2019. A series of stakeholder engagement events were also undertaken in October 2018 as part of consultation phase 3.5. These events were held to inform the public of potential changes to the onshore substation location. This consultation aims to ensure that community concerns are well understood and that site specific issues can be taken into account, where practicable. Consultation phases are explained further in *Chapter 5 EIA Methodology.*
- 10. No public consultation feedback specific to Offshore Ornithology has been raised during any the public consultation undertaken to date. Full details of the proposed East Anglia TWO project consultation process will be presented in the Consultation Report, which will be submitted as part of the DCO application

## 12.3Scope

- 11. This chapter describes the ornithological interests of the windfarm site and the offshore cable corridor to landfall, and evaluates the potential impacts of the proposed East Anglia TWO project on these interests.
- 12. The baseline section describes the distribution and abundance of bird species recorded during surveys of the site. This includes flight characteristics (e.g. height and direction), ecology, seasonality and behaviour.
- 13. The predicted magnitude of impacts and significance of effects arising due to construction, operation and decommissioning of the windfarm on the ornithological interests of the site are assessed on the basis of the worst case development scenario. Measures to prevent or reduce significance of the possible effects are discussed where appropriate. Cumulative impacts arising from the site and offshore cable corridor and other offshore operations are assessed as appropriate.



## 12.3.1 Study Area

- 14. A study area was defined that was relevant to the consideration of potential impacts on offshore ornithological receptors. The suitability of the study area for the purpose of environmental impact assessment was agreed with Natural England and the RSPB during the Evidence Plan Process.
- 15. This study area includes the East Anglia TWO windfarm site and a 4km buffer placed around it (*Figure 12.1*). Monthly aerial surveys of the study area began in November 2015, ceased in April 2016, re-started in September 2016 until October 2017 (20 months in total). An additional four months of surveys was undertaken in May to August 2018 to provide a complete 24-month dataset. The final DCO submission will use all of these data, however the analysis and assessment in this PEIR has been undertaken prior to the data from the final aerial surveys being available, so is based on the first 21 monthly surveys and will be updated when 24 surveys are available.
- 16. The data collected during these aerial surveys have been used to identify the bird species present and their seasonal abundance.
- 17. In addition to the windfarm area covered by aerial surveys, the study area over which potential impacts on offshore bird species were considered included the offshore cable corridor to the Mean Low Water Spring (MLWS) at its landfall location in the vicinity of Sizewell and Thorpeness. Refer to *Chapter 23 Onshore Ornithology* for assessment of impacts above the MLWS.

## 12.3.2 Worst Case

- 18. The design of the proposed East Anglia TWO project (including number of wind turbines, layout configuration, requirement for scour protection, electrical design, etc.) is not yet fully determined, and may not be known until sometime after the DCO has been granted. Therefore, in accordance with the requirements of the Project Design Envelope (also known as the Rochdale Envelope) approach to Environmental Impact Assessment (EIA) (Planning Inspectorate 2018) (as discussed in *Chapter 5 EIA Methodology*), realistic worst case scenarios in terms of potential effects upon offshore ornithology are adopted to undertake a precautionary and robust impact assessment.
- 19. The worst-case scenarios for potential impacts of the proposed project on offshore ornithology receptors from the construction, operation and decommissioning phases are described and presented in *Table 12.2*. Where percentage areas affected have been calculated, these are based on a total windfarm site area of 255km<sup>2</sup> and an offshore cable corridor area of 123km<sup>2</sup> which results in a total offshore development area for the assessment of 378km<sup>2</sup>. As a worst case, the offshore cable corridor area has been calculated based on the northern route (see *Figure 9.2*) which has the largest area of the



two routes and from which the worst case export cable length was calculated. It would not be realistic to combine the areas for both route options as in reality only one of these routes will be used following final design of the project.

- 20. Definition of the worst case scenarios assumptions has been made from consideration of the proposed East Anglia TWO project parameters that are presented in *Chapter 6 Project Description*, alongside the mitigation measures that have been embedded in the design (*section 12.3.3*).
- 21. It should be noted that after collision risk modelling (CRM) was conducted for the proposed East Anglia ONE North project, the design envelope was changed so that the maximum number of wind turbines increased from 67 to 75 for the 12MW scenario, and from 53 to 60 for the 15MW scenario. The collision risk modelling presented in this assessment is for the previous scenarios of 67 12MW, 53 and 4819MW turbines (see *Appendix 12, Annex 3*). The collision risk model has not been re-run for the updated scenarios because of time constraints however an assessment of the updated parameters will be included within the ES. This model re-run will also incorporate the remaining three months of aerial survey data (see *section 12.3.1*).

Impact	Parameter	Rationale
Construction		
Impact 1 Disturbance and Displacement from increased vessel activity	It is anticipated that the installation of the offshore elements will take approximately 27 months. Construction works would be undertaken 24 hours a day and seven days a week offshore, dependent upon weather conditions.	The worst case scenario is based on the longest construction period and the maximum numbers of plant on site and operational at a given time.
	The maximum number of all types of vessels operating simultaneously within the offshore development area during construction would be 74.	
	There would be up to three foundation installation vessels (i.e. Dynamic Positioning Heavy Lift Vessels) on site at any one time.	
	Maximum of 1,005 helicopter round trips per annum assumed	
	Installation of the export cable would take place over a twelve month period split into two separate six month periods and up to two cable laying vessels operating	

 Table 12.2 Project Design: Realistic Worst case scenarios for the Proposed East Anglia TWO

 Project



Impact	Parameter	Rationale
	simultaneously.	
	Speed of cable laying vessels – maximum speed of 300m per hour for ploughing or jetting and 80m per hour if trenching (see <i>Chapter 6 Project</i> <i>Description</i> .	
Impact 2 Indirect effects as a result	Spatial worst case impact– maximum hammer energy of 4,000 kilo Joules (kJ).	See <b>Chapter 10 Fish and</b> <b>Shellfish Ecology</b> for a full breakdown of the maximum disturbed area of sea bed.
of displacement of prey species due to increased noise and disturbance to	Up to three foundation installation vessels (i.e. Dynamic Positioning Heavy Lift Vessels) on site at any one time.	
sea bed	Temporal worst case impact	
	No concurrent piling, 75 wind turbine foundations, five offshore electrical and construction operation and maintenance platforms, and one operational met mast.	
	The maximum worst case area of disturbance to benthic habitats during construction would be 10,543,760m <sup>2</sup> across the offshore development area, equivalent to 2.79% of the maximum offshore development footprint.	Breakdown is given in <i>Chapter 9</i> <i>Benthic ecology</i> .
	Disturbance / displacement from increased suspended sediment concentration from the excavation of up to 4,322,423m <sup>3</sup> of sediment in the offshore development area over the whole 27 month construction period.	Total sediment release over the construction period is given in <b>Chapter 9 Benthic Ecology</b> and <b>Chapter 7 Marine Geology and</b> <b>Physical Processes.</b> However, the release on a daily basis would be temporary and localised with sediment settling out quickly.
Operation	-	
Impact 3	A windfarm area of 255km <sup>2</sup> plus 4km	Maximum density of turbines and
Disturbance and displacement from offshore infrastructure	with a minimum spacing of 800m in row x 1200m between rows	project area, which maximises the potential for avoidance and
and due to increased vessel and helicopter activity	Maximum of 687vessel round trips per annum to support windfarm operations.	นเอยเลยอาเมียาน.
(Includes barrier effect)	Maximum of 1,005 helicopter round trips per annum for scheduled and	



Impact	Parameter	Rationale
	unscheduled maintenance.	
	Lighting requirements for the offshore windfarm will need to be consistent with maritime and aviation safety requirements, and are expected to consist of:	
	• Obstruction lighting compliant with CAA aviation safety requirements, as a minimum, requiring turbines on the periphery of the windfarm to be lit;	
	• Maritime navigational safety lighting compliant with Trinity House Light House service safety requirements which requires navigational lighting to be visible at a distance of at least 5nm, lights would be required to be placed low on the turbines and electrical platforms.	
	• Search and Rescue (SAR) lighting consistent with MCA safety requirements. These are most likely to be infra-red lighting which would only be activated during search and rescue operations.	
Impact 4 Collision risk	Maximum of 75 250m (12 MW) wind turbines, other scenarios are 60 300m (15 MW) or 48 300m (19MW) wind turbines.	CRM has been carried out for all turbine scenarios based on the turbine specifications in <i>Appendix 12.1, Annex 3, Table</i> <b>5</b> . The worst case scenario in terms of the highest collision risk varies between bird species. For each species, the turbine scenario which produces the highest collision risk has been used in the assessment (see <i>section 12.6.2.3</i> below).
Impact 5 Indirect effects due to habitat loss / change for	The maximum possible sea bed footprint of the project, and therefore habitat loss, would be:	The maximum possible above sea bed footprint of the project including scour or scour
key prey species	Windfarm Site Infrastructure	protection plus any cable
	2,028,406m <sup>2</sup> which constitutes 0.80% of the windfarm site (75 turbine foundations, five offshore electrical and construction operation and maintenance platforms, one meteorological mast, cable protection for platform link cables and	See Chapter 9 Benthic Ecology and Chapter 10 Fish and Shellfish Ecology.



Impact	Parameter	Rationale
	inter-array cables.	
	Export Cable	
	176,800m <sup>2</sup> , 0.14% of the northern offshore cable corridor which has been used a worst case as it has the largest area of the two cable route options.	
	Total	
	The overall total footprint which could be subject to permanent habitat loss would therefore be 2,205,206m <sup>2</sup> (0.58% of the offshore development area).	
Decommissioning		
Impact 6	Assumed similar to construction and	
Disturbance and Displacement from increased vessel activity	therefore a worst case would be as above in impact 1.	
Impact 7 Indirect effects as a result of displacement of prey species due to increased noise and disturbance to sea bed	There would be habitat disturbance effects over up to 2,677,414m <sup>2</sup> across the offshore development area (0.70% of maximum offshore development area). There would be limited noise disturbance to prey (as no piling and no use of explosives).	See Chapter 9 Benthic Ecology and Chapter 10 Fish and Shellfish Ecology.

## 12.3.3 Embedded Mitigation

- 22. A number of mitigation measures which are embedded into the proposed project design are relevant to offshore ornithology receptors.
- 23. The East Anglia TWO windfarm site was identified through the Zonal Appraisal and Planning process (*Chapter 4 Site Selection and Alternatives*). The windfarm area avoids European sites, although the cable corridor runs through the Outer Thames Estuary SPA and pSPA Extension (Royal HaskoningDHV 2018).
- 24. In order to reduce the spatial extent of potential disturbance and displacement impacts the decision was taken to use only one offshore cable corridor in the near shore for East Anglia TWO and East Anglia ONE North. This measure avoids potential impacts over a wider area.



## 12.3.3.1 Best practice protocol for red-throated divers

- 25. As has been accepted for East Anglia THREE, a best-practice protocol for minimising disturbance to red-throated divers during construction will be adopted. This would comprise some or all of the following measures.
  - restricting vessel movements to existing navigation routes (where the densities of divers are typically relatively low);
  - where it is necessary to go outside of established navigational routes, selecting routes that avoid known aggregations of birds;
  - maintaining direct transit routes (to minimise transit distances through areas used by divers);
  - avoidance of over-revving of engines (to minimise noise disturbance); and,
  - briefing of vessel crew on the purpose and implications of these vessel management practices (through, for example, tool-box talks).
- 26. Once further information is available about the port(s) that will be used for construction, operations and maintenance, then appropriate vessel traffic management measures including, where relevant, some or all of the above best practice examples can be formulated in agreement with Natural England.
- 27. If used, helicopters are a potential source of disturbance to red throated diver in the Outer Thames Estuary SPA. The minimum safe altitude for helicopters operating offshore is 1000 feet above the highest known obstacle within 5nm. It is considered that at these altitudes that any disturbance caused by the visual presence or noise of helicopters will be minimal and will not result in significant disturbance of red-throated diver.

## 12.3.4 Monitoring

28. Post-consent, the final detailed design of the proposed East Anglia TWO project and the development of the relevant management plan(s) will refine the worstcase parameters assessed in the EIA. It is recognised that monitoring is an important element in the management and verification of the impacts of the proposed East Anglia TWO project. Outline management plans, across a number of environmental topics, will be submitted with the DCO application. These outline management plans will contain key principles that provide the framework for any monitoring that could be required. The requirement for and final appropriate design and scope of monitoring will be agreed with the relevant stakeholders and included within the relevant management plan(s), submitted for approval, prior to construction works commencing.



## **12.4Assessment Methodology**

## 12.4.1 Legislation, Policy and Guidance

29. Legislation relevant to offshore ornithology is identified in *Table 12.3* along with a summary of important measures derived from it.

#### Table 12.3 Summary of Legislation and Relevant Measures

Legislation	Relevant Measures
Birds Directive - Council Directive 2009/147/EC on the Conservation of Wild Birds	This Directive provides a 'General System of Protection' for all species of naturally occurring wild birds in the EU. The most relevant provisions of the Directive are the identification and classification of Special Protection Areas (SPAs) for rare or vulnerable species listed in Annex I of the Directive and for all regularly occurring migratory species (required by Article 4). It also establishes a general scheme of protection for all wild birds (required by Article 5). The Directive requires national Governments to establish SPAs and to have in place mechanisms to protect and manage them. The SPA protection procedures originally set out in Article 4 of the Birds Directive have been replaced by the Article 6 provisions of the Habitats Directive.
	The UK has triggered article 50 of the Treaty of European Union and is currently in the process of withdrawing from the European Union (EU). Recent UK Government Guidance (September 2018) states that 'The EU Withdrawal Act 2018 will ensure all existing EU environmental law continues to operate in UK law, providing businesses and stakeholders with certainty as we leave the EU.'
Wildlife and Countryside Act 1981, as amended	The Wildlife and Countryside Act 1981 (as amended) is the principal mechanism for the legislative protection of wildlife in Great Britain. It provides protection for all species of wild birds and their nests and establishes the system of Sites of Special Scientific Interest (SSSI).
The Conservation of Offshore Marine Habitats and Species Regulations 2017	The Conservation of Offshore Marine Habitats and Species Regulations 2017 (as amended), (referred to here as the 'Offshore Regulations') transposes the Birds Directive and the Habitats Directive into national law in the offshore environment (beyond 12 nautical miles within British Fishery Limits and the UK Continental Shelf Designated Area. The Offshore Regulations place an obligation on 'competent authorities' to carry out an appropriate assessment of any proposal likely to affect a SAC or SPA, to seek advice from Natural England and / or JNCC, and to not approve an application that would have an adverse effect on the integrity of a SAC or SPA (except under very tightly constrained conditions).
The Conservation of Habitats and Species Regulations 2017	The Conservation of Habitats and Species Regulations 2017 (hereafter called the 'Habitats Regulations'), transposes the Birds Directive and the Habitats Directive into national law in the onshore environment and territorial waters out to 12 nautical miles, operating in conjunction with the Wildlife and Countryside Act 1981. The Habitats Regulations place an obligation on 'competent authorities' to carry out an appropriate assessment of any proposal likely to affect a



Legislation	Relevant Measures
	SAC or SPA, to seek advice from Natural England and / or JNCC, and to not approve an application that would have an adverse effect on the integrity of a SAC or SPA (except under very tightly constrained conditions).

30. Policy relevant to offshore ornithology is identified in *Table 12.4* along with a summary of important measures derived from it.

Policy	Relevant Measures
Overarching National Policy Statement (NPS) for Energy (NPS EN-1) (July 2011)	Paragraph 5.3.3 states that the Applicant should ensure that the ES clearly sets out any effects on internationally, nationally and locally designated sites of ecological or geological conservation importance, on protected species and on habitats and other species identified as being of principal importance for the conservation of biodiversity. Paragraph 5.3.4 states that the Applicant should also show how the proposed project has taken advantage of opportunities to conserve and enhance biodiversity and geological conservation interests. Paragraph 5.3.18 states that the Applicant should include appropriate mitigation measures as an integral part of the proposed development.
NPS for Renewable Energy Infrastructure (NPS EN-3) (July 2011)	Paragraph 2.6.64 states that the assessment of offshore ecology and biodiversity should be undertaken by the Applicant for all stages of the lifespan of the proposed offshore windfarm. Paragraph 2.6.102 states that the scope, effort and methods required for ornithological surveys should have been discussed with the relevant statutory advisor. Paragraph 2.6.104 states that it may be appropriate for the assessment to include collision risk modelling for certain bird species.
National Planning Policy Framework	The National Planning Policy Framework sets out the UK Government's planning policies for England and how these are expected to be applied. The document establishes a number of core land-use planning principles that should underpin both plan-making and decision-taking, including contributing to conserving and enhancing the natural environment.
	Paragraph 170 states that: "Planning policies and decisions should contribute to and enhance the natural and local environment byminimising impacts on and providing net gains for biodiversity, including by establishing coherent ecological networks that are more resilient to current and future pressures".
UK Post-2010 Biodiversity Framework	The 'UK Post-2010 Biodiversity Framework' succeeds the UK Biodiversity Action Plan. The Framework demonstrates how the work of the four countries and the UK contributes to achieving the Aichi Biodiversity Targets, and identifies the activities required to

## Table 12.4 Summary of Policy and Relevant Measures



Policy	Relevant Measures
	complement the country biodiversity strategies in achieving the targets.
UK Marine Policy Statement (MPS)	New systems of marine planning are being introduced in the UK. The MPS, adopted under section 44 of the Marine and Coastal Access Act 2009, is the framework for developing and implementing regional Marine Plans. It will contribute to the achievement of sustainable development in the United Kingdom marine area. High level objectives are for the protection, conservation and where appropriate recovery of biodiversity; healthy, resilient and adaptable marine and coastal ecosystems across their natural range; and oceans supporting viable populations of representative, rare, vulnerable and valued species.

- 31. The most relevant guidance on Environmental Impact Assessment (EIA) for marine ecology receptors, including birds, is the 'Guidelines for Ecological Impact Assessment in the UK and Ireland: Terrestrial, Freshwater, Coastal and Marine' published by the Chartered Institute of Ecology and Environmental Management (CIEEM 2018). The EIA methodology described in *section 12.4.3* and applied in this chapter is based on that CIEEM guidance.
- 32. Additional guidance on the assessment of the potential impacts of renewable energy generation on birds has been produced by a number of statutory bodies, NGOs and consultants including, but not limited to the following:
  - Assessment methodologies for offshore windfarms (Maclean et al., 2009);
  - Guidance on ornithological cumulative impact assessment for offshore wind developers (King et al. 2009);
  - Advice on assessing displacement of birds from offshore windfarms (SNCB 2017);
  - Collision risk modelling to assess bird collision risks for offshore windfarms (Band 2012);
  - Assessing the risk of offshore wind farm development to migratory birds (Wright et al. 2012);
  - Vulnerability of seabirds to offshore windfarms (Furness and Wade, 2012; Furness et al., 2013; Wade et al. 2016);
  - Mapping seabird sensitivity to Offshore Windfarms (Bradbury et al. 2014);
  - The avoidance rates of collision between birds and offshore turbines (Cook et al. 2014); and
  - Joint Response from the Statutory Nature Conservation Bodies to the Marine Scotland Science Avoidance Rate Review (JNCC et al. 2014).



## 12.4.2 Data Sources

#### 12.4.2.1 Site specific surveys

33. Site specific aerial surveys of the East Anglia TWO windfarm site (and 4km buffer) were conducted between November 2015 and April 2016, September 2016 and October 2017, and May to August 2018, to complete 24 months of site-specific data available for assessment. The analysis and assessment in this PEIR has been undertaken prior to the data from the final aerial surveys being available, so is based on the first 21 monthly surveys and will be updated when 24 surveys are available.

### 12.4.2.2 Other relevant surveys

- 34. The former East Anglia Zone has been subject to extensive ornithological surveys as described in this section. The survey data have not been used for quantitative analysis but provide context for the assessment.
- 35. For the purposes of Zonal Environmental Appraisal (ZEA), 18 months of high resolution aerial survey data were collected across the former East Anglia Zone, including;
  - The Crown Estate Enabling Action data (video aerial survey) from November 2009 to March 2010; and
  - APEM aerial survey data from April 2010 to April 2011.
- 36. Between November 2009 and March 2011 and September 2011 and December 2012, 33 months of aerial survey were completed of the south-west portion of the former East Anglia Zone. These surveys overlapped the East Anglia TWO windfarm site by 92% and therefore provide valuable additional data.
- 37. Surveys of the East Anglia ONE windfarm site to the east were conducted between November 2009 and October 2011, and for the East Anglia THREE windfarm site to the north-east between September 2011 and August 2013.

## 12.4.2.3 Desk based assessment

- 38. The desk based assessment has drawn on a wide variety of published literature, covering both peer reviewed scientific literature and the 'grey literature' such as windfarm project submissions and reports. It includes the published literature on seabird ecology and distribution and on the potential impacts of windfarms (both derived from expert judgement and post-construction monitoring studies). The key topics for which the literature has been examined include:
- 39. Potential impacts of windfarms (Garthe and Hüppop 2004; Drewitt and Langston 2006; Stienen et al. 2007; Speakman et al. 2009; Langston 2010;



Band 2012; Cook et al. 2012; Furness and Wade 2012; Wright et al. 2012; Furness et al. 2013; Johnston et al. 2014a and b).

- Bird population estimates (Mitchell et al. 2004; BirdLife International 2004; Holling et al. 2011; Holt et al. 2012; Musgrove et al. 2013; Furness 2015).
- Bird breeding ecology (Cramp and Simmons 1977-94; Del Hoyo et al. 1992-2011; Robinson 2005).
- Bird distribution (Stone et al. 1995; Brown and Grice 2005; Kober et al. 2010; Balmer et al. 2013).
- Bird migration and foraging movements (Wernham et al. 2002; Thaxter et al. 2012).
- Red-throated diver densities in the Outer Thames Estuary SPA (JNCC 2013) and data from an unpublished report on surveys carried out in 2013 by APEM for Natural England.
- East Anglia Offshore Wind: Zonal Assessment Report (APEM 2011).
- 40. Information on statutory sites and their interest features has been drawn from the web-based resource Multi-Agency Geographic Information for the Countryside (MAGIC www.magic.defra.gov.uk) and the Natural England and JNCC web sites (www.naturalengland.org.uk; www.jncc.defra.gov.uk).

## 12.4.3 Impact Assessment Methodology

- 41. The impact assessment methodology applied in this Chapter is based on that described *Chapter 5 EIA Methodology*, adapted to make it applicable to ornithology receptors and aligned with the key guidance document produced on impact assessment on ecological receptors (CIEEM 2018).
- 42. The methodology applied in this chapter has also been the subject of extensive consultation with Natural England and RSPB through the Evidence Plan process for the proposed East Anglia TWO project, and informed by discussion during the examination process for the consented East Anglia ONE and East Anglia THREE projects.
- 43. The assessment approach uses the conceptual 'source-pathway-receptor' model. The model identifies likely environmental impacts resulting from the proposed construction, operation and decommissioning of the offshore infrastructure. This process provides an easy to follow assessment route between impact sources and potentially sensitive receptors, ensuring a transparent impact assessment. The parameters of this model are defined as follows:



- Source the origin of a potential impact (noting that one source may have several pathways and receptors) e.g. an activity such as cable installation and a resultant effect such as re-suspension of sediments.
- Pathway the means by which the effect of the activity could impact a receptor e.g. for the example above, re-suspended sediment could settle and smother the sea bed.
- Receptor the element of the receiving environment that is impacted e.g. for the above example, bird prey species living on or in the sea bed are unavailable to foraging birds.

## 12.4.3.1 Sensitivity

44. Definitions of the different sensitivity levels for ornithology receptors, using the example of disturbance from construction activity, are included in *Table 12.5*.

Table 12.5 Definitions of the Different Sensitivity Levels for Ornithology Receptors in Relation to
Construction Disturbance

Sensitivity	Definition
High	Ornithology receptor (bird species) has <u>very limited</u> tolerance of sources of disturbance such as noise, light, vessel movements and the sight of people
Medium	Ornithology receptor (bird species) has <u>limited</u> tolerance of sources of disturbance such as noise, light, vessel movements and the sight of people
Low	Ornithology receptor (bird species) has <u>some</u> tolerance of sources of disturbance such as noise, light, vessel movements and the sight of people.
Negligible	Ornithology receptor (bird species) is <u>generally</u> tolerant of sources of disturbance such as noise, light, vessel movements and the sight of people.

## 12.4.3.2 Conservation Value

- 45. The conservation value of ornithological receptors is based on the population from which individuals are predicted to be drawn. This reflects current understanding of the movements of bird species. Therefore, conservation value for a species can vary through the year depending on the relative sizes of the number of individuals predicted to be at risk of impact and the population from which they are estimated to be drawn. Ranking therefore corresponds to the degree of connectivity which is predicted between the windfarm site and protected populations. Using this approach, the conservation importance of a species seen at different times of year may fall into any of the defined categories.
- 46. Example definitions of the value levels for ornithology receptors are given in **Table 12.6**. These are related to connectivity with populations that are protected as qualifying species of Special Protection Areas (SPAs). SPAs are



internationally designated sites which carry strong protection for populations of qualifying bird species and are a key consideration for the ornithology assessment.

Value	Definition
High	A species for which individuals at risk can be clearly connected to a particular Special Protection Area (SPA or pSPA).
Medium	A species for which individuals at risk are probably drawn from particular SPA or pSPA populations, although other populations (both SPA and non-SPA) may also contribute to individuals at risk
Low	A species for which individuals at risk on have no known connectivity to SPAs, or for which no SPAs are designated.

#### Table 12.6 Definitions of the Conservation Value Levels for an Ornithology Receptor

### 12.4.3.3 Magnitude

47. The definitions of the magnitude levels for ornithology receptors are set out in *Table 12.7*. This set of definitions has been determined on the basis of changes to bird populations.

Table 12.7 Definitions of the Magnitude Levels for Ornitho	logy Receptors
--	----------------

Value	Definition
High	A change in the size or extent of distribution of the relevant biogeographic population or the population that is the interest feature of a specific protected site that is predicted to irreversibly alter the population in the short-to-long term and to alter the long-term viability of the population and / or the integrity of the protected site. Recovery from that change predicted to be achieved in the long-term (i.e. more than 5 years) following cessation of the development activity.
Medium	A change in the size or extent of distribution of the relevant biogeographic population or the population that is the interest feature of a specific protected site that occurs in the short and long-term, but which is not predicted to alter the long-term viability of the population and / or the integrity of the protected site.
	Recovery from that change predicted to be achieved in the medium-term (i.e. no more than five years) following cessation of the development activity.
Low	A change in the size or extent of distribution of the relevant biogeographic population or the population that is the interest feature of a specific protected site that is sufficiently small-scale or of short duration to cause no long-term harm to the feature / population. Recovery from that change predicted to be achieved in the short-term (i.e. no more than one year) following cessation of the development activity.
Negligible	Very slight change from the size or extent of distribution of the relevant biogeographic population or the population that is the interest feature of a specific protected site. Recovery from that change predicted to be rapid (i.e. no more than circa 6 months) following cessation of the development related activity.



Value	Definition
No change	No loss of, or gain in, size or extent of distribution of the relevant biogeographic population or the population that is the interest features of a specific protected site.

## 12.4.3.4 Impact Significance

48. Following the identification of the receptor value and sensitivity and the determination of the magnitude of the effect, the significance of the impact will be determined. That determination will be guided by the matrix as presented in *Table 12.8*. Impacts shaded red or orange represent those with the potential to be significant in EIA terms (see *paragraph 51* below).

Sensitivity	Magnitude						
	High	Medium	Low	Negligible	No change		
High	Major	Major	Moderate	Minor	No change		
Medium	Major	Moderate	Minor	Negligible	No change		
Low	Moderate	Minor	Minor	Negligible	No change		
Negligible	Minor	Negligible	Negligible	Negligible	No change		

#### Table 12.8 Impact Significance Matrix

- 49. It is important that the matrix (and indeed the definitions of sensitivity and magnitude) is seen as a framework to aid understanding of how a judgement has been reached from the narrative of each impact assessment. It is not a prescriptive formulaic method. Expert judgement has been applied to the assessment of likelihood and ecological significance of a predicted impact.
- 50. In particular it should be noted that high conservation value and high sensitivity are not necessarily linked for a particular impact. A receptor could be of high conservation value (e.g. an interest feature of a SPA) but have a low or negligible physical/ecological sensitivity to an effect and vice versa. Potential impact significance will not be inflated simply because a feature is 'valued'. Similarly, potentially highly significant impacts will not be deflated simply because a feature is not "valued". The narrative behind the assessment is important here; the conservation value of an ornithological receptor can be used where relevant as a modifier for the sensitivity (to the effect) already assigned to the receptor.
- 51. For the purpose of this assessment the CIEEM (2018) guidance has been followed. This states that *'significance is a concept related to the weight that should be attached to effects when decisions are made... so that the decision*



maker is adequately informed of the environment consequences of permitting a project'. CIEEM (2018) defines significance as follows: 'In broad terms, significant effects encompass impacts on the structure and function of defined sites, habitats or ecosystems and the conservation status of habitats and species (including extent, abundance and distribution). Significant effects should be qualified with reference to an appropriate geographic scale, for example a significant effect on a Site of Special Scientific Interest ... is likely to be of national significance.'

- 52. Where possible, assessment is based upon quantitative and accepted criteria (for example, guidance from Statutory Nature Conservation Bodies (SNCBs) on collision risk modelling (Band 2012, and displacement (SNCB 2017), and /or biological removal thresholds determined through population modelling), together with the use of value judgement and expert interpretation to establish to what extent an impact is significant.
- 53. The assessment refers to and includes embedded mitigation (**section 12.3.3**). No further requirements for mitigation have been identified and thus there is no assessment of residual impacts post-mitigation.

## 12.4.4 Project Design Envelope

- 54. The project design envelope, which sets out a series of design options, is described in *section 6.1.1* of *Chapter 6 Project Description*. In accordance with the Planning Inspectorate (2018), this has a reasoned range and maximum extent for a number of key parameters (for example options for different numbers of wind turbines of different size). The project design envelope is used to establish the maximum extent to which the project would impact on the environment. The final detailed design of the project, including spatial, temporal and installation methodology, could then vary within this 'Rochdale Envelope' without rendering the assessment inadequate.
- 55. For ornithology receptors, on a precautionary basis, the assessment is based on the aspects of the design envelope considered to be worst case in terms of effects on birds. These are summarised in *Table 12.2* above. For example, for collision risk, the worst case scenario is the design option with the highest estimated collision risk for a given bird species.

## 12.4.5 Cumulative Impact Assessment

- 56. The impact assessment methodology applied in this Chapter is based on that described in *Chapter 5 EIA Methodology*, adapted to make it applicable to ornithology receptors.
- 57. The methodology has also been aligned with the approach to the assessment of cumulative impacts that has been applied by Ministers when consenting



offshore windfarms and confirmed in recent consent decisions. It also follows the approach set out in recent guidance from the Planning Inspectorate (Planning Inspectorate 2015) and from the renewables industry (RenewableUK 2013).

### 12.4.6 Transboundary Impact Assessment

- 58. The transboundary impact assessment methodology applied in this Chapter is based on that described in *Chapter 5 EIA Methodology*, adapted to make it applicable to ornithology receptors.
- 59. The potential for transboundary impacts is identified by consideration of potential linkages to non-UK protected sites and sites with large concentrations of breeding, migratory or wintering birds (including the use of available information on tagged birds).

## **12.5Existing Environment**

60. The characterisation of the existing or baseline environment is undertaken based on the site based surveys (listed in *section 12.4.2.1* above and as detailed in *Appendix 12.1*), the desk study (*section 12.4.2.3*), and other relevant literature.

## 12.5.1 Key Species

61. The bird species recorded during site-specific surveys (digital photographic aerial bird surveys of the windfarm site plus a 4km buffer, described in *Appendix 12.1*) of the East Anglia TWO windfarm site to date are listed in *Table 12.9* along with details of their conservation status. The locations of all species observed are plotted on figures in *Appendix 12.1*.

Species	Scientific name	Conservation Status
Red-throated diver	Gavia stellata	Birds of Conservation Concern (BoCC) <sup>1</sup> Green listed, Birds Directive Annex 1
Black-throated diver	Gavia arctica	BoCC Amber listed, Birds Directive Annex 1
Great northern diver	Gavia immer	BoCC Amber listed, Birds Directive Annex 1
Fulmar	Fulmarus glacialis	BoCC Amber listed, Birds Directive Migratory Species
Gannet	Morus bassanus	BoCC Amber listed, Birds Directive Migratory Species
Cormorant	Phalacrocorax carbo	BoCC Amber listed, Birds Directive Migratory Species

#### Table 12.9 Species Recorded in the East Anglia TWO Study Area and Their Conservation Status



Species	Scientific name	Conservation Status
Shag	Phalacrocorax aristotelis	BoCC Red listed, Birds Directive Migratory Species
Great skua	Stercorarius skua	BoCC Amber listed, Birds Directive Migratory Species
Puffin	Fratercula arctica	BoCC Red listed, Birds Directive Migratory Species
Razorbill	Alca torda	BoCC Amber listed, Birds Directive Migratory Species
Guillemot	Uria aalge	BoCC Amber listed, Birds Directive Migratory Species
'Commic' tern <sup>2</sup>	Tern species	BoCC Amber listed, Birds Directive Annex 1
Kittiwake	Rissa tridactyla	BoCC Red listed, Birds Directive Migratory Species
Black-headed gull	Chroicocephalus ridibundus	BoCC Amber listed, Birds Directive Migratory Species
Little gull	Hydrocoloeus minutus	BoCC Green listed, Birds Directive Migratory Species
Common gull	Larus canus	BoCC Amber listed, Birds Directive Migratory Species
Lesser black-backed gull	Larus fuscus	BoCC Amber listed, Birds Directive Migratory Species
Herring gull	Larus argentatus	BoCC Red listed, Birds Directive Migratory Species
Great black-backed gull	Larus marinus	BoCC Amber listed, Birds Directive Migratory Species

1. Eaton et al. 2015.

2. 'Commic tern' is used as a collective term where Arctic tern *Sterna paradisaea* and common tern *Sterna hirundo* could not be distinguished at distance or from aerial survey images.

- 62. For the offshore cable corridor, no site-specific ornithology surveys were carried out. The assessment for this component of the development has been carried out with reference to available data (from JNCC (2013) and data from an unpublished report on surveys carried out in 2013 by APEM for NE).
- 63. Species assessed for impacts are those which were recorded during surveys and which are considered to be at potential risk either due to their abundance,



potential sensitivity to windfarm impacts or due to biological characteristics (e.g. tendency to fly at rotor heights) which make them potentially susceptible.

- 64. Impacts have been assessed in relation to relevant biological seasons, as defined by Furness (2015). For the non-breeding period, the seasons and relevant biologically defined minimum population sizes (BDMPS) were taken from Furness (2015).
- 65. For the breeding period, the potential for connectivity to known breeding populations has been considered. However, it should be noted that bird abundance was low for most species during the breeding season, with many species absent in one or more of the summer months. This suggests that very few breeding birds utilise the East Anglia TWO windfarm site. The seasonal definitions in Furness (2015) include overlapping months in some instances due to variation in the timing of migration for birds which breed at different latitudes (i.e. individuals from breeding sites in the north of the species' range may still be on spring migration when individuals farther south have already commenced breeding). Due to the very low presence of breeding birds it was considered appropriate to define breeding as the migration-free breeding period (see Table 12.10), sometimes also referred to as the core breeding period. This ensured that any late or early migration movements which were observed were assessed in relation to the appropriate reference populations. One exception to this was lesser black-backed gull, for which there is potential that breeding adults from the Alde Ore Estuary SPA population may forage on the East Anglia TWO windfarm site. Hence for this species the full breeding season was applied in the attribution of potential impacts to relevant populations.

Species	Breeding	Migration- free breeding	Migration - autumn	Winter	Migration - spring	Non- breeding
Red-throated	Mar-Aug	May-Aug	Sep-Nov	Dec-Jan	Feb-Apr	-
diver			(13,277)	(10,177)	(13,277)	
Black-throated diver <sup>1</sup>	Apr-Aug	May-Aug				Aug-Apr
Great northern diver	-	-	Sep-Nov	Dec-Feb	Mar-May	Sep-May (200)
Fulmar	Jan-Aug	Apr-Aug	Sep-Oct (957,502)	Nov (568,736)	Dec-Mar (957,502)	-

## Table 12.10 Species Specific Definitions of Biological Seasons and BDMPS (from Furness 2015 or Other Sources) for Bird Species Recorded During Baseline Surveys for the Proposed East Anglia Two Project



Species	Breeding	Migration- free breeding	Migration - autumn	Winter	Migration - spring	Non- breeding
Gannet	Mar-Sep	Apr-Aug	Sep-Nov	-	Dec-Mar	-
			(456,298)		(248,385)	
Cormorant	Apr-Aug	May-Jul	Aug-Oct	Nov-Jan	Feb-Apr	Sep-Mar
Shag	Feb-Aug	Mar-Jul	Aug-Oct	Nov	Dec-Feb	Sep-Jan
Great skua	May-Aug	May-Jul	Aug-Oct	Nov-Feb	Mar-Apr	-
			(19,556)	(143)	(8,485)	
Puffin	Apr-Aug	May-Jun	Jul-Aug	Sep-Feb	Mar-Apr	Mid-Aug- Mar
						(231,957)
Razorbill	Apr-Jul	Apr-Jun	Aug-Oct	Nov-Dec	Jan-Mar	-
			(591,874)	(218,622)	(591,874)	
Guillemot	Mar-Jul	Mar-Jun	Jul-Oct	Nov	Dec-Feb	Aug-Feb
						(1,617,306)
'Commic' tern <sup>2</sup>	May-Aug	Jun	Jul-Sep	-	Apr-May	-
			(308,841)		(308,841)	
Kittiwake	Mar-Aug	May-Jul	Aug-Dec	-	Jan-Apr	-
			(829,937)		(627,816)	
Black-headed gull <sup>1</sup>	-	Apr-Jul	-	-	-	Aug-Mar
Little gull <sup>1</sup>	Apr-Jul	May-Jul	-	-	-	Aug-Apr
Common gull <sup>1</sup>	-	May-Jul	-	-	-	Aug-Apr
Lesser black-	Apr-Aug	May-Jul	Aug-Oct	Nov-Feb	Mar-Apr	-
backed gull			(209,007)	(39,314)	(197,483)	
Herring gull	Mar-Aug	May-Jul	Aug-Nov	Dec	Jan-Apr	Sep-Feb
						(466,511)
Great black-	Mar-Aug	May-Jul	Aug-Nov	Dec	Jan-Apr	Sep-Mar
Dacked gull	one not includ	ded within Europ	(2015): 600	ed on Natural	England 2012	(91,399) (Black

throated diver) or Birds of the Western Palearctic (other species).

2. Combined estimate for common and Arctic tern populations from Furness (2015).



66. The mean peak abundances within species-specific seasons (as defined in *Table 12.10*) recorded within the East Anglia TWO windfarm site are provided in *Table 12.11*. The mean peak in any given season was calculated as follows: (i) the population density and abundance for each survey was calculated using design-based estimation methods, with 95% confidence intervals calculated using non-parametric bootstrapping (see *Technical Appendix 12.1* for further details); (ii) the abundance for each calendar month was calculated as the mean of estimates for each month (e.g. mean of up to two values); (iii) the seasonal mean peak was taken as the highest from within the months falling in each season. In some cases the peak was recorded in a month which is included in overlapping seasons and therefore the same value has been identified in both seasons. These have been identified in italics in *Table 12.11*.

Table 12.11 Mean Peak Counts (and 95% Confidence Intervals) by Biological Season for Bird Species Within the East Anglia TWO Windfarm Site Recorded During Baseline Surveys. Figures in Italics Identify the Same Peak Occurring in Different Seasons Due to Overlapping Months

Species	Breeding	Migration- free breeding	Migration - autumn	Winter	Migration - spring	Non- breeding
Red-throated diver	130.2	7.7	7.7	28.1	130.2	-
	(0-258.9)	(0-44.6)	(0-37)	(0-87.1)	(0-258.1)	
Black-throated diver <sup>1</sup>	0	0	-	-	-	0
	(0-0)	(0-0)				(0-0)
Great northern diver <sup>2</sup>	-	-	0	0	0	0
			(0-0)	(0-0)	(0-0)	(0-0)
Fulmar	86.8	86.8	35.8	5.1	25.5	-
	(0-210.8)	(0-210.8)	(0-73.5)	(0-24.5)	(0-61.9)	
Gannet	145.6	145.6	646.1	-	143	-
	(0-209.2)	(0-209.2)	(0-1,297.7)		(0-363.2)	
Cormorant	0	0	0	0	5.1	-
	(0-0)	(0-0)	(0-0)	(0-0)	(0-27.7)	
Shag <sup>1</sup>	0	0	0	0	0	0
	(0-0)	(0-0)	(0-0)	(0-0)	(0-0)	(0-0)
Great skua	0	0	5.1	0	5.1	-
	(0-0)	(0-0)	(0-26)	(0-0)	(0-31.6)	
Puffin	12.8	0	0	0	12.8	0
	(0.54.7)	(0-0)	(0-0)	(0-0)	(0.54.7)	(0-0)


Species	Breeding	Migration- free breeding	Migration - autumn	Winter	Migration - spring	Non- breeding
Razorbill	114.9	114.9	28.1	71.5	145.6	-
	(0-318.4)	(0-318.4)	(0-73.2)	(0-137.6)	(10.7-233.2)	
Guillemot	1,299.8	1,299.8	423.9	674.2	1,236	1,236
	(11.7- 2,014.7)	(11.7- 2,014.7)	(0-557.5)	(135.9- 1,299)	(128.2- 1,424.2)	(0-1,424.2)
'Commic' tern	23	0	15.3	-	23	-
	(0-53.5)	(0-0)	(0-40.2)		(0-53.5)	
Kittiwake	229.8	201.7	86.8	-	229.8	-
	(0-510.4)	(0-294.7)	(0-196.9)		(0-510.4)	
Black-headed	-	-	0	-	-	84.3
gull			(0-0)			(0-225.1)
Little gull	5.1	0	-	-	-	38.3
	(0-27.3)	(0-0)				(0-111.5)
Common gull	-	0	-	-	-	5.1
		(0-0)				(0-33.4)
Lesser black-	20.4	20.4	28.1	15.3	5.1	-
backed gull	(0-75.3)	(0-75.3)	(0-91.9)	(0-58.1)	(0-31.6)	
Herring gull	15.3	15.3	17.9	0	5.1	17.9
	(0-39.5)	(0-39.5)	(0-73.5)	(0-0)	(0-25.2)	(0-73.5)
Great black-	28.1	28.1	12.8	17.9	20.4	20.4
backed gull	(0-64.5)	(0-64.5)	(0-55.1)	(0-55.9)	(0-61.9)	(0-61.9)

1. Recorded on only one occasion in baseline surveys.

2. Recorded on only two occasions in baseline surveys.

#### 12.5.2 Designated Sites

- 67. This section considers potential connectivity of the proposed East Anglia TWO Project with sites with statutory designation for nature conservation which have birds listed as qualifying features. The sites considered are SPAs, pSPAs, Ramsar sites and SSSIs.
- 68. Sites which may have connectivity to the East Anglia TWO windfarm site and offshore cable corridor include those designated for breeding seabirds and



those for terrestrial / coastal / marine bird interests (typically overwintering aggregations). Discussions held with Natural England and RSPB through the EPP have scoped out connectivity for migratory non-seabird species associated with coastal and terrestrial sites (*Table 12.1*).

- 69. The assessment therefore focuses on sites designated for breeding seabird colonies and coastal/offshore sites for overwintering seabirds.
- 70. The offshore ornithology section of the Habitats Regulations Assessment (HRA) Screening Report (Royal HaskoningDHV 2018) considers 86 offshore and coastal designated sites within or adjacent to the southern North Sea within 950km of the East Anglia TWO windfarm site. These comprise SPAs and Ramsar sites designated for bird interests, with terrestrial areas of coastal sites also designated as SSSIs (to Mean Low Water Springs).
- 71. The HRA screening identified four sites for further consideration in relation to potential effects. All remaining sites were not considered to be within range or to have a pathway for a potential effect in relation to the proposed East Anglia TWO project. Although the HRA is separate from the EIA, the screening carried out is also considered to be appropriate in terms of identifying potential connectivity for the ornithological impact assessment, so the same four sites are identified here. These are listed in *Table 12.12*.

Site	Designation	Ornithological interest features with potential connectivity to the proposed East Anglia TWO project	Minimum distance to project (km)
The Outer Thames Estuary	SPA	Wintering seabirds – red-throated diver	0 (windfarm site) 0 (offshore cable corridor)
Greater Wash	SPA	Wintering seabirds – red-throated diver, little gull	29 (windfarm site) 23 (offshore cable corridor)
Alde-Ore Estuary	SPA Ramsar SSSI	Breeding seabirds – lesser black- backed gull	36 (windfarm site) 3 (offshore cable corridor)
Flamborough and Filey Coast*	SPA SSSI	Breeding seabirds – gannet, kittiwake, razorbill, guillemot	246 (windfarm site) 237 (offshore cable corridor)
*The recently designate SPA and is a larger at	ed Flamborough and rea, so the latter site	Filey Coast SPA includes the Flamboroug is not considered as a separate entity a	h Head and Bempton Cliffs is it is encompassed in the

Table 12.12 Designated Sites for Birds with Potential Connectivity to the Proposed East AngliaTWO Project



Site	Designation	Ornithological interest features with potential connectivity to the proposed East Anglia TWO project	Minimum distance to project (km)
revised site			

- 72. Where a species that is a qualifying feature of one or more of the designated sites listed in *Table 12.12* is screened in for assessment in relation to a potential impact, the potential for connectivity with that site is considered in the assessment.
- 73. The assessment of likely significant effect on the interest features of the internationally designated sites (SPAs and Ramsar sites) is carried out through the HRA process and this is reported separately in the Draft Report to Inform the Habitats Regulations Assessment which accompanies this PEIR.

#### **12.5.3 Anticipated Trends in Baseline Condition**

- 74. Key drivers of seabird population size in western Europe are climate change (Sandvik et al. 2012; Frederiksen et al. 2004, 2013; Burthe et al. 2014; Macdonald et al. 2015; Furness 2016; JNCC 2016), and fisheries (Tasker et al. 2000; Frederiksen et al. 2004; Ratcliffe 2004; Carroll et al. 2017; Sydeman et al. 2017). Pollutants (including oil, persistent organic pollutants, plastics), alien mammal predators at colonies, disease, and loss of nesting habitat also impact on seabird populations but are generally much less important and often more local factors (Ratcliffe 2004; Votier et al. 2005, 2008; JNCC 2016).
- 75. Trends in seabird numbers in breeding populations are better known, and better understood than trends in numbers at sea within particular areas. Breeding numbers are regularly monitored at many colonies (JNCC 2016), and in the British Isles there have been three comprehensive censuses of breeding seabirds in 1969-70, 1985-88 and 1998-2002 (Mitchell et al. 2004) as well as single-species surveys (such as the decadal counts of breeding gannet numbers, Murray et al. 2015). In contrast, the European Seabirds at Sea (ESAS) database is incomplete, and few data have been added since 2000, so that current trends in numbers at sea in areas of the North Sea are not so easy to assess.
- 76. Breeding numbers of many seabird species in the British Isles are declining, especially in the northern North Sea (Foster and Marrs 2012; Macdonald et al. 2015; JNCC 2016). The most striking exception is gannet, which continues to increase (Murray et al. 2015), although the rate of increase has been slowing (Murray et al. 2015). These trends in British seabird populations seem likely to continue in the short to medium term future.



- 77. Climate change is likely to be the strongest influence on seabird populations in coming years, with anticipated deterioration in conditions for breeding and survival for most species of seabirds (Burthe et al. 2014; Macdonald et al. 2015; Capuzzo et al. 2018) and therefore further declines in numbers of most of our seabird populations are anticipated. It is therefore highly likely that breeding numbers of most of our seabird species will continue to decline under a scenario with continuing climate change due to increasing levels of greenhouse gases. Fisheries management is also likely to influence future numbers in seabird populations. The Common Fisheries Policy (CFP) Landings Obligation ('discard ban') will further reduce food supply for scavenging seabirds such as great black-backed gulls, lesser black-backed gulls, herring gulls, fulmars, kittiwakes and gannets (Votier et al. 2004; Bicknell et al. 2013; Votier et al. 2013; Foster et al. 2017). Recent changes in fisheries management that aid recovery of predatory fish stock biomass are likely to further reduce food supply for seabirds that feed primarily on small fish such as sandeels, as those small fish are major prey of large predatory fish. Therefore, anticipated future increases in predatory fish abundance resulting from improved management to constrain fishing mortality on those commercially important species at more sustainable levels than in the past are likely to cause further declines in stocks of small pelagic seabird 'food-fish' such as sandeels (Frederiksen et al. 2007; Macdonald et al. 2015). Lindegren et al. (2018) concluded that sandeel stocks in the North Sea, the most important prey fish stock for North Sea seabirds during the breeding season (Furness and Tasker 2000), have been depleted by high levels of fishing effort. These stocks are unlikely to recover fully even if fishing effort was reduced, because climate change has altered the North Sea food web to the detriment of productivity of fish populations. As a result, seabird populations are likely to continue to experience food shortages in the North Sea, especially for those species most dependent on sandeels as food.
- 78. Future decreases in kittiwake breeding numbers are likely to be particularly pronounced, as kittiwakes are very sensitive to climate change (Frederiksen et al. 2013; Carroll et al. 2015) and to fishery impacts on sandeel stocks near breeding colonies (Frederiksen et al. 2004; Carroll et al. 2017), and the species will lose the opportunity to feed on fishery discards as the Landings Obligation comes into effect. Gannet numbers may continue to increase for some years, but evidence suggests that this increase is already slowing (Murray et al. 2015), and numbers may peak not too far into the future. While the Landings Obligation will reduce discard availability to gannets in European waters, in recent years increasing proportions of adult gannets have wintered in west African waters rather than in UK waters (Kubetzki et al. 2009), probably because there are large amounts of fish discarded by west African trawl fisheries and decreasing amounts available in the North Sea (Kubetzki et al. 2009; Garthe et al. 2012). The flexible behaviour and diet of gannets probably



reduces their vulnerability to changes in fishery practices or to climate change impacts on fish communities (Garthe et al. 2012).

- 79. Fulmars, terns, common guillemot, razorbill and puffin appear to be highly vulnerable to climate change, so numbers may decline over the next few decades (Burthe et al. 2014). Strong declines in shag numbers are likely to continue as they are adversely affected by climate change, by low abundance of sandeels and especially by stormy and wet weather conditions in winter (Burthe et al. 2014; Frederiksen et al. 2008). Most of the red-throated divers and common scoters wintering in the southern North Sea originate from breeding areas at high latitudes in Scandinavia and Russia. Numbers of redthroated divers and common scoters wintering in the southern North Sea may possibly decrease in future if warming conditions make the Baltic Sea more favourable as a wintering area for those species so that they do not need to migrate as far as UK waters. There has been a trend of increasing numbers of sea ducks remaining in the Baltic Sea overwinter (Mendel et al. 2008; Fox et al. 2016; Ost et al. 2016) and decreasing numbers coming to the UK (Austin and Rehfisch 2005; Pearce-Higgins and Holt 2013), and that trend is likely to continue, although to an uncertain extent.
- 80. ESAS data indicate that there has already been a long-term decrease in numbers of great black-backed gulls wintering in the southern North Sea (S. Garthe et al. in prep.), and the Landings Obligation will probably result in further decreases in numbers of north Norwegian great black-backed gulls and herring gulls coming to the southern North Sea in winter. It is likely that further redistribution of breeding herring gulls and lesser black-backed gulls will occur into urban environments (Rock and Vaughan 2013), although it is unclear how the balance between terrestrial and marine feeding by these gulls may alter over coming years; that may depend greatly on the consequences of Brexit for UK fisheries and farming. Some of the human impacts on seabirds are amenable to effective mitigation (Ratcliffe et al. 2009; Brooke et al. 2018), but the scale of efforts to reduce these impacts on seabird populations has been small by comparison with the major influences of climate change and fisheries. This is likely to continue to be the case in future, and the conclusion must be that with the probable exception of gannet, numbers of almost all other seabird species in the UK North Sea region will most likely be on a downward trend over the next few decades, due to population declines, redistributions or a combination of both.
- 81. For offshore ornithology, the ecological impact assessment is therefore carried out against declining baseline populations of a number of receptor species. Where a receptor species is declining, the assessment takes into account whether a given impact is likely to exacerbate a decline in the relevant



reference population, and prevent a receptor species from recovery should environmental conditions become more favourable. Climate change has been identified as the strongest influence on future seabird population trends. In this context it is noted that a key component of global strategies to reduce climate change is the development of low-carbon renewable energy developments such as offshore windfarms.

### **12.6Potential Impacts**

- 82. Potential impacts to be included within the EIA have been agreed through consultation on a Method Statement (Scottish Power Renewables 2017, Appendix 2.4) with Natural England and RSPB during the Evidence Plan process. They are as follows:
  - In the construction phase:
    - Impact 1: Disturbance/displacement; and
    - Impact 2: Indirect impacts through effects on habitats and prey species.
  - In the operational phase:
    - Impact 3: Disturbance and displacement from offshore infrastructure and due to increased vessel and helicopter activity;
    - Impact 4: Collision risk; and
    - Impact 5: Indirect impacts through effects on habitats and prey species
  - In the decommissioning phase:
    - o Impact 6: Disturbance/displacement; and
    - Impact 7: Indirect impacts through effects on habitats and prey species.
- 83. In the assessment of potential impacts below they are assessed:
  - In the order of construction, operation and decommissioning;
  - Following the impact assessment methodology that is described in *section 12.4.3*;
  - On the basis of the worst case potential impacts set out in *section 12.3.2*; and
  - Accounting for the embedded mitigation that is described in *section 12.3.3.*

#### **12.6.1 Potential Impacts During Construction**

#### 12.6.1.1 Direct Disturbance and Displacement

84. The construction phase of the proposed project has the potential to affect bird populations in the marine environment through disturbance due to construction activity leading to displacement of birds from construction sites. This would effectively result in temporary habitat loss through reduction in the area



available for feeding, loafing and moulting. The worst case, outlined in **Table 12.2** describes the elements of the proposed project considered within this assessment.

- 85. The maximum duration of offshore construction for the proposed project would be 25 months which would overlap with a maximum of two breeding seasons, two winter periods and up to four spring/autumn migration periods.
- 86. The construction phase would require the mobilisation of vessels, helicopters and equipment and the installation of foundations, export cables and other infrastructure. These activities have the potential to disturb and displace birds from within and around the offshore elements of the proposed East Anglia TWO project, including the windfarm and the sub-sea cables. Causes of potential disturbance would comprise the presence of construction vessels and associated human activity, noise and vibration from construction activities and lighting associated with construction sites. The level of disturbance at each work location would differ dependent on the activities taking place, but there could be vessel movements at any time of day or night over the construction period.
- 87. Any impacts resulting from disturbance and displacement from construction activities would be short-term, temporary and reversible in nature, lasting only for the duration of construction activity, with birds expected to return to the area once construction activities have ceased. Construction related disturbance and displacement is most likely to affect foraging birds. Furthermore, modelling of the consequences of displacement for fitness of displaced birds suggests that even in the case of breeding seabirds that are displaced on a daily basis, there is likely to be little or no impact on survival unless the offshore windfarm is close to the breeding colony (Searle et al. 2014, 2017).
- 88. Bird species differ in their susceptibility to anthropogenic disturbance and in their responses to noise and visual disturbance stimuli. The principal source of noise during construction of the offshore windfarm would be subsea noise from piling works associated with the installation of foundations for wind turbines and associated offshore substations. While assessed for marine mammals and fish, subsea noise is not considered a risk factor for diving birds. Seabirds and other diving bird species will spend most of their time above or on the water surface, where hearing will detect sound propagated through the air. It has been speculated, based on what is known about the physiology of hearing in birds, and comparison to the underwater hearing ability of humans, that birds do not hear well underwater (Dooling and Therrien 2012). Anatomical studies of ear structure in diving birds suggest that there are adaptations for protection against the large pressure changes that may occur while diving, which may reduce hearing ability underwater but also protect the ear from damage due to acoustic



over-exposure (Dooling and Thierren 2012). Above water noise disturbance from construction activities is not considered in isolation as a risk factor for birds; but rather, combined with the presence of vessels, man-made structures, and human activity, part of the overall disturbance stimulus that causes birds to avoid boats and other structures – as discussed below.

- 89. Lighting of construction sites, vessels and other structures at night may potentially be a source of attraction (phototaxis), as opposed to displacement, for birds; however, the areas affected would be very small, and restricted to offshore construction areas which are active at a given time – a maximum of 72 offshore construction structures and a maximum of 74 offshore vessels may be active within the offshore development area. Phototaxis can be a serious hazard for fledglings of some seabird species, but occurs over short distances (hundreds of metres) in response to bright white light close to breeding colonies of these species. It is not seen over large distances or in older (adult and immature) seabirds (Furness 2018). Construction sites associated with the offshore development area would be far enough removed from any seabird breeding colonies as to render this risk negligible. Phototaxis of nocturnal migrating birds can be a problem, especially in autumn during conditions of poor visibility, but is generally seen where birds are exposed to intense white lighting such as from lighthouses; light from construction sites is likely to be one or two orders of magnitude less powerful than that from lighthouses (Furness 2018).
- 90. Considering variation between species in response to disturbance, gulls are not considered susceptible to disturbance, as they are often associated with fishing boats (e.g. Camphuysen 1995; Hüppop and Wurm 2000) and have been noted in association with construction vessels at the Greater Gabbard offshore windfarm (GGOWL 2011) and close to active foundation piling activity at the Egmond aan Zee (OWEZ) windfarm, where they showed no noticeable reactions to the works (Leopold and Camphuysen 2007). However, species such as divers and scoters have been observed to avoid shipping by several kilometres (Mitschke et al. 2001 from Exo et al. 2003; Garthe and Hüppop 2004; Schwemmer et al. 2011).
- 91. There are a number of different measures used to assess bird disturbance and displacement from areas of sea in response to activities associated with an offshore windfarm. Garthe and Hüppop (2004) developed a scoring system for such disturbance factors which they applied to seabird species in German sectors of the North Sea. This was refined by Furness and Wade (2012) and Furness et al. (2013) with a focus on seabirds using Scottish offshore waters. The approach uses information in the scientific and 'grey' literature, as well as expert opinion to identify disturbance ratings for individual species, alongside



scores for habitat flexibility and conservation importance. These factors were used to define an index value that highlights the sensitivity of a species to disturbance and displacement. As many of these references relate to disturbance from helicopter and vessel activities, these are considered relevant to this assessment.

- 92. Birds recorded during the species-specific spring and autumn migration periods are assumed to be moving through the area between breeding and wintering areas. As these individuals will be present in the site for a short time and the potential zone of construction displacement will be comparatively small, it has been assumed that there are negligible risks of impact at these times of year. Consequently, the following assessment focuses on the breeding and nonbreeding periods (seasons following Furness 2015).
- 93. In order to focus the assessment of disturbance and displacement, a screening exercise was undertaken to identify those species most likely to be at risk (*Table 12.13*). Any species recorded only in very small numbers within the Study Area (including the offshore cable corridor) or with a low sensitivity to displacement was screened out of further assessment.
- 94. The species screened in for assessment were red-throated diver, razorbill and guillemot. These were assessed for impacts during the periods and spatial locations where effects were potentially likely.

Species	Sensitivity to Disturbance and Displacement <sup>1</sup>	Screening Result (IN or OUT)	Rationale
Red-throated diver	Very High	IN	For the offshore export cable corridor only as this overlaps with the Outer Thames Estuary SPA for which red- throated diver is a qualifying species
Black-throated diver	Very High	OUT	Recorded on only one occasion in baseline surveys
Great northern diver	Very High	OUT	Recorded on only two occasions in baseline surveys
Fulmar	Low	OUT	Low susceptibility to disturbance
Gannet	Low	OUT	Low susceptibility to disturbance

<b>Table 12.13 Construction</b>	<b>Disturbance and Dis</b>	placement Screening
---------------------------------	----------------------------	---------------------



Species	Sensitivity to Disturbance and Displacement <sup>1</sup>	Screening Result (IN or OUT)	Rationale
Cormorant	High	OUT	Recorded on only one occasion in baseline surveys
Shag	Medium	OUT	Recorded on only one occasion in baseline surveys
Great skua	Low	OUT	Recorded in low numbers during passage migration periods
Puffin	Medium	OUT	Recorded on in low numbers during only one baseline survey
Razorbill	Medium	IN	Potentially susceptible to disturbance and abundant
Guillemot	Medium	IN	Potentially susceptible to disturbance and abundant
'Commic' tern <sup>2</sup>	Low	OUT	Low susceptibility to disturbance and recorded in low numbers
Kittiwake	Low	OUT	Low susceptibility to disturbance
Black-headed gull	Low	OUT	Low susceptibility to disturbance
Little gull	Low	OUT	Low susceptibility to disturbance
Common gull	Low	OUT	Low susceptibility to disturbance
Lesser black- backed gull	Low	OUT	Low susceptibility to disturbance
Herring gull	Low	OUT	Low susceptibility to disturbance
Great black-backed gull	Low	OUT	Low susceptibility to disturbance

1. With reference to Garthe and Hüppop, 2004; Furness and Wade, 2012; Furness et al., 2013; Wade et al., 2016.

2. 'Commic tern' is used where an arctic tern and common tern could not be distinguished at distance or from aerial survey images



#### 12.6.1.1.1 Red-throated Diver

- Red-throated diver has been identified as being particularly sensitive to human 95. activities in marine areas, including through the disturbance effects of ship and helicopter traffic (Garthe and Hüppop 2004; Schwemmer et al. 2011; Furness and Wade 2012; Furness et al. 2013; Bradbury et al. 2014). A selectivity index derived from aerial surveys in the German North Sea indicated that the numbers of divers (red- and black-throated divers could not be reliably distinguished during the surveys) were significantly lower in shipping lanes than in other areas, although there were insufficient data to estimate flush distances of divers from ships (Schwemmer et al. 2011); in this study it was assumed that the responses of red and black-throated divers to disturbance was similar. Observational studies of responses of marine birds to disturbance in Orkney inshore waters found that red-throated and black-throated divers showed similar flush behaviour from ferries (with respectively 75% (n=88) and 62% (n=21) of birds showing an evasive response within 300m of a passing ferry). Redthroated divers were highly likely to fly in response to marine activity whereas black-throated divers were more likely to swim away (although these differences may be related to differences in the timing of moult in the two species, which affects flight ability) (Jarett et al. 2018).
- 96. There is potential for disturbance and displacement of non-breeding redthroated divers resulting from the presence of vessels installing the offshore cable, including when it is laid through the Outer Thames Estuary SPA. The offshore cable corridor extends from the landfall approximately 5km north of Aldeburgh in a North-East orientation, passing through approximately 25km of the SPA in areas that are predominantly not shipping routes so may represent more important habitat for red-throated divers. Where it overlaps with the SPA, the offshore cable corridor width is between 2km and 4km, giving a total potential overlap between the export cable corridor and the SPA of approximately 132km<sup>2</sup> which represents an overlap with the SPA of approximately 3.5%, although this represents the area of search and the actual cable route itself will be much smaller. Cable-laying operations, utilising up to two vessels, have the potential to displace red-throated divers from an area around each vessel. However, cable laying vessels are static for large periods of time, and move slowly and over short distances as cable installation takes place. Offshore cable installation activity is also a relatively low noise emitting operation, particularly when compared to activities such as piling.
- 97. The assessment takes account of embedded mitigation in the form of a best practice protocol for minimising construction disturbance of red-throated divers (see *section 12.3.3.1*). The magnitude of disturbance to red-throated diver has been estimated on a 'Worst Case' basis. This assumes that there would be 100% displacement of those birds in a 2km buffer surrounding the source, in



this case a maximum of two cable laying vessels. This 100% displacement is consistent with the suggestion that all red-throated divers present fly away from approaching vessels at a distance of 1km or less (Bellebaum et al. 2006; Topping and Petersen 2011). This may be a very precautionary assumption, for example (as noted above) studies of responses of marine birds to disturbance in Orkney inshore waters found that 75% (n=88) red-throated divers flushed within 300m of ferries (Jarett et al 2018), implying that in this study not all birds were flushed within 300m of vessels.

- 98. To estimate the number of red-throated divers that would potentially be at risk of displacement from the offshore cable corridor during the cable laying process, the density of red-throated divers along the offshore cable corridor was estimated. Where the export cable overlaps with the SPA, the density of red-throated divers was assumed to be equivalent to the overall density within the Outer Thames Estuary SPA, estimated as 1.7 birds per km<sup>2</sup> (the designated population of 6,466 divided by the total SPA area (for red-throated divers<sup>1</sup>) of 3,792.7 km<sup>2</sup>).
- The 'worst case' area from which birds could be displaced was defined as a 99. circle with a 2km radius around each cable laying vessel, which is 25.2km<sup>2</sup> (2 x 12.6km<sup>2</sup>). If 100% displacement is assumed to occur within this area, then 43 divers would be displaced at any given time. It is considered reasonable to assume that birds will reoccupy areas following passage of the vessel. The cable laying vessels will move at a maximum speed of 300m per hour for ploughing or jetting and 80m per hour if trenching (Chapter 6 Project Description and see also Table 12.2). This represents a maximum speed of 7m per minute. For context, a modest tidal flow rate for the Outer Thames would be in the region of 30m per minute (0.5m per second, derived from DECC 2009). The tide would therefore be flowing at least four times faster than the cable laying vessel. Birds on the water surface are likely to be drifting with the tide and moving at the same speed as the tidal flow. Thus, even while moving, the vessels would be effectively stationary as far as birds are concerned, so the zone of impact around the vessel would be more or less fixed. Consequently, for the purposes of this assessment it can be assumed that the estimated number of red-throated divers displaced at any one time from cable-laying vessels represents the total number displaced over the course of a single winter.

<sup>&</sup>lt;sup>1</sup> Note that the total area of the Outer Thames Estuary SPA, including a recent extension to include coastal and riverine areas used for foraging by breeding little terns, is now 3,924.5km<sup>2</sup>. As the extension was not proposed for red-throated divers, the area of the original SPA has been used as the reference to calculate the density of this species.



- 100. Definitive mortality rates associated with displacement for any seabird are not known and precautionary estimates have to be used. There is no empirical evidence that displaced birds suffer any consequent mortality; any mortality due to displacement would be most likely a result of increased density in areas outside the affected area, resulting in increased competition for food where density was elevated. Such impacts are most likely to be negligible (Dierschke et al. 2017), and below levels that could be quantified. Impacts of displacement are also likely to be context-dependent. In years when food supply has been severely depleted, as for example by unsustainably high fishing mortality of sandeel stocks as has occurred several times in recent decades (ICES 2013; Lindegren et al. 2018), displacement of sandeel-dependent seabirds from optimal habitat may increase mortality. In years when food supply is good, displacement is unlikely to have any negative effect on seabird populations. Red-throated divers may feed on sandeels, but take a wide diversity of small fish prey, so would be buffered to an extent from fluctuations in abundance of individual fish species. It is not possible for the proposed East Anglia TWO project to predict future fishing effort. However, this assessment has assumed the precautionary maximum mortality rate associated with the displacement of red-throated diver in the wintering period is 1-5% (i.e. 1-5% of displaced individuals suffer mortality as a direct consequence<sup>2</sup>). At this level of mortality then only 0-2 birds would be expected to die across the entire winter period (September to April) as a result of any potential displacement effects from the offshore cable installation activities. The average annual mortality rate for redthroated diver, across age classes, is estimated as 0.228 (based on species specific data from Horswill and Robinson (2015); see Table 12.16 below). Based on this, 2320 birds would be expected to die each year from the winter BDMPS for this species (10,177; Furness 2015). The addition of a maximum of two birds to this would increase the mortality rate by 0.09%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Thus, this highly precautionary assessment generates an effect of negligible magnitude.
- 101. The construction works, specifically offshore cable laying, are temporary and localised in nature and the magnitude of effect on red-throated diver has been determined as negligible. As the species is of high sensitivity to disturbance, the impact significance is **minor adverse**.
- 102. Natural England has asked for consideration to be given to a seasonal restriction on construction works, between November and February, to minimise effects on red-throated divers (*Table 12.1*). Given the assessment above and the precautionary prediction that a maximum of two birds would die as a result

<sup>&</sup>lt;sup>2</sup> This assumption was applied for East Anglia THREE



of displacement over this period, a seasonal restriction is not considered to be required in addition to the measures set out in the best practice protocol for red-throated divers in *section 12.3.3.1* above.

#### 12.6.1.1.2 Razorbill

- 103. Razorbills were recorded in the East Anglia TWO windfarm site in most months throughout the year, in the highest numbers in the non-breeding season, peaking in January (mean density in windfarm site 0.57/km<sup>2</sup>) and at their lowest in June and August (mean density in windfarm site 0/km<sup>2</sup>). Razorbills are considered to have a medium general sensitivity to disturbance and displacement, based on their sensitivity to ship and helicopter traffic in Garthe and Hüppop (2004), Furness and Wade (2012), Furness et al. (2013) and Bradbury et al. (2014).
- 104. There is potential for disturbance and displacement of razorbills due to construction activities, including the construction of wind turbines and other infrastructure (offshore electrical platforms, construction operation and maintenance platforms and meteorological mast) and associated vessel traffic. However, construction will not occur across the whole of the proposed wind turbine array area simultaneously or every day but will be phased, with no more than three wind turbine foundations expected to be installed at any time. Consequently, the effects will occur only in the areas where vessels are operating at any given point and not the entire East Anglia TWO windfarm site.
- 105. For this precautionary assessment it has been assumed that 1-10% of displaced individuals could die as a result of displacement by construction vessels (as for displacement from the operational windfarm – see section 12.6.2.1 below).
- 106. During the autumn migration season, at a mean peak density of 0.11/km<sup>2</sup> and with a highly precautionary 2km radius of disturbance around each of three construction areas (wind turbines or other infrastructure), 4 individual birds (0.11 x 12.56 x 3) could be at risk of displacement, of which <1 bird (0-0.4 birds) would be predicted to die. The average annual mortality rate for razorbill, across age classes, is estimated as 0.174 (based on species specific data from Horswill and Robinson (2015); see *Table 12.16* below). Based on this, 102,986 birds would be expected to die each year from the winter BDMPS for this species (591,874; Furness 2015). The addition of 0.4 birds to this would increase the mortality rate by <0.001%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Thus this highly precautionary assessment generates an effect of negligible magnitude.



- 107. The construction works are temporary and localised in nature and the magnitude of effect has been determined as negligible. As razorbill is of medium sensitivity to disturbance, the impact significance is **negligible**.
- 108. During the winter period, at a mean peak density of 0.28/km<sup>2</sup> and with a highly precautionary 2km radius of disturbance around each of three construction areas (wind turbines or other infrastructure), 11 individual birds (0.28 x 12.56 x 3) could be at risk of displacement, of which 0-1 would be expected to die. Based on the average mortality for the species, a total of 38,040 birds would be expected to die each year from the winter BDMPS for this species (218,622; Furness 2015). The addition of a maximum of 1 bird would increase the mortality rate by <0.01%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Thus this highly precautionary assessment generates an effect of negligible magnitude.</p>
- 109. The construction works are temporary and localised in nature and the magnitude of effect has been determined as negligible. As razorbill is of medium sensitivity to disturbance, the impact significance is **negligible**.
- 110. During the spring migration season, at a mean peak density of 0.57/km<sup>2</sup> and with a highly precautionary 2km radius of disturbance around each of three construction areas (wind turbines or other infrastructure), 21 individual birds (0.57 x 12.56 x 3) could be at risk of displacement, of which 0-2 would be expected to die. Based on the average mortality for the species, a total of 102,986 birds would be expected to die each year from the spring migration BDMPS for this species (591,874; Furness 2015). The addition of a maximum of 2 birds would increase the mortality rate by <0.01%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Thus, this highly precautionary assessment generates an effect of negligible magnitude.
- 111. The construction works are temporary and localised in nature and the magnitude of effect has been determined as negligible. As razorbill is of medium sensitivity to disturbance, the impact significance is **negligible**.
- 112. During the breeding season, the maximum mean peak density on the site was 0.45/km<sup>2</sup> (April) which suggests that 17 individuals (0.45 x 12.56 x 3) could be at risk of displacement, of which 0-2 would be expected to die.
- 113. The mean maximum foraging range for breeding razorbill is 48.5km (Thaxter et al. 2012) which places the East Anglia TWO windfarm site considerably beyond the range of any razorbill breeding colonies. The nearest major breeding



colony is Flamborough Head, 246 km from the site (the minimum distance to the Flamborough and Filey Coast SPA, *Table 12.12*).

- It should be noted that some recent tagging studies have recorded larger 114. apparent distances than this (one razorbill was recorded travelling 312 km from Fair Isle) which might indicate connectivity to breeding colonies. However, further consideration of this apparent potential for connectivity indicates how exceptional this result is. A razorbill flies at about 16m per second (Pennycuick 1987) so would take almost eleven hours to complete this round trip even if it spent no time on the water or diving for food. This is incompatible with bringing enough food back to keep a chick alive as razorbill chicks receive about 3 feeds per day (Harris and Wanless 1989). Yet chicks are normally attended and protected by one adult at the nest site while the partner is foraging (Wanless and Harris 1989), so there are simply not enough hours in a single day to allow successfully breeding razorbills to make such long trips to provision a chick. At 16m per second the East Anglia TWO windfarm site is 4.3 hours direct flight time away from the nearest razorbill breeding colony (Flamborough Head). A return trip would take close to 9 hours, not allowing for foraging. As for the Fair Isle example, travelling such distances is incompatible with successful breeding. On the basis of 3 feeds per day, the furthest away a bird could fly per trip to achieve this in 24 hours is 115km (i.e. a round trip of 230km), with no allowance for foraging time. Even if the bird spends a maximum of only 30 minutes foraging, this reduces the farthest distance to 108km.
- 115. On the basis of the above evidence, it can be stated with certainty that there are no breeding colonies for razorbill within normal foraging range of the East Anglia TWO windfarm site, therefore it is reasonable to assume that individuals seen during the breeding season are nonbreeding (e.g. immature birds). Since immature seabirds are known to remain in wintering areas, the number of immature birds in the relevant population during the breeding season may be estimated as 43% of the total wintering BDMPS population (Furness 2015). This gives a breeding season population of 94,007 (BDMPS for the UK North Sea and Channel, 218,622 x 43%).
- 116. Based on the average mortality for the species, a total of 16,357 birds would be expected to die each year from the sub-adult component of the winter BDMPS for this species (94,007; Furness 2015). The addition of a maximum of two birds predicted to die from construction disturbance and displacement would increase the mortality rate by 0.01%. (Use of the average mortality produces a conservative estimate of % change, as the mortality of birds less than 2 years old is higher than (survival rates are lower than) that of adult birds, *Table 12.16*). This magnitude of increase in mortality would not materially alter the



background mortality of the population and would be undetectable. Thus this highly precautionary assessment generates an effect of negligible magnitude.

117. The construction works are temporary and localised in nature and the magnitude of effect has been determined as negligible. As the species is of medium sensitivity to disturbance, the impact significance is **negligible**.

#### 12.6.1.1.3Guillemot

- 118. Guillemots were recorded in the East Anglia TWO windfarm site year round, with densities peaking in January (mean density in windfarm site 4.84/km<sup>2</sup>) and at their lowest in June (mean density in windfarm site 0.040/km<sup>2</sup>). Guillemots are considered to have a medium general sensitivity to disturbance and displacement, based on their sensitivity to ship and helicopter traffic in Garthe and Hüppop (2004), Furness and Wade (2012), Furness et al. (2013) and Bradbury et al. (2014).
- 119. There is potential for disturbance and displacement of guillemots due to construction activities, including the construction of wind turbines and other infrastructure (offshore electrical platforms, construction operation and maintenance platforms and met mast) and associated vessel traffic. However, construction will not occur across the whole of the proposed wind turbine array area simultaneously or every day but will be phased, with no concurrent piling. Other installation vessels would be largely stationary and move short distances between turbine locations. Consequently, the effects will occur only in the areas where vessels are operating at any given point and not the entire East Anglia TWO windfarm site.
- 120. For this precautionary assessment, it has been assumed that 1-10% of displaced individuals could die as a result of displacement by construction vessels (as for displacement from the operational windfarm – see section 12.6.2.1.3 below).
- 121. During the nonbreeding season, at a mean peak density of 4.84/km<sup>2</sup> and with a highly precautionary 2km radius of disturbance around each of three active construction areas (wind turbines or other infrastructure), 182 individual birds (4.84 x 12.56 x 3) could be at risk of displacement, of which 2-18 would be expected to die. The average annual mortality rate for guillemot, across age classes, is estimated as 0.14 (based on species specific data from Horswill and Robinson (2015); see **Table 12.16** below). Based on this, 226,423 birds would be expected to die each year from the non-breeding season BDMPS for this species (1,617,306; UK North Sea and English Channel, Furness 2015). The addition of 2-18 birds to this would increase the mortality rate by <0.01%. This magnitude of increase in mortality would not materially alter the background



mortality of the population and would be undetectable. Thus, this highly precautionary assessment generates an effect of negligible magnitude.

- 122. The construction works are temporary and localised in nature and the magnitude of effect has been determined as negligible. As guillemot is of medium sensitivity to disturbance, the impact significance is **negligible**.
- 123. During the breeding season, the maximum mean peak density in the East Anglia TWO windfarm site was 5.09/km<sup>2</sup> (April) which suggests that 192 individuals (5.09 x 12.56 x 3) could be at risk of displacement, of which 2-19 would be expected to die.
- 124. The mean maximum foraging range for breeding guillemot is 84.2km (Thaxter et al. 2012) which places the East Anglia TWO windfarm site considerably beyond the range of any guillemot breeding colonies. The nearest breeding colony is Flamborough Head, 246 km from the site (the minimum distance to the Flamborough and Filey Coast pSPA, *Table 12.12*).
- It should be noted that some recent tagging studies have recorded larger 125. apparent distances than the mean maximum from Thaxter et al. 2012, for example one guillemot was recorded travelling 340km from Fair Isle. This might indicate connectivity to breeding colonies. At greater distances. However, further consideration indicates how exceptional this result is. The 340km figure is derived from an individual guillemot on Fair Isle in a year when the local sandeel stock collapsed and breeding success was close to zero (this bird's chick died). A common guillemot flies at about 19m per second (Pennycuick 1987) so would take almost 10 hours to complete this round trip even if it spent no time on the water or diving for food. This is incompatible with bringing enough food back to keep a chick alive. The species carries only one fish at a time and common guillemot chicks need about 5 feeds per day. Yet chicks are normally attended and protected by one adult at the nest site while the partner is foraging (Uttley et al. 1994), so there are simply not enough hours in the day to allow successfully breeding guillemots to make such long trips to provision a chick. At 19m per second the East Anglia TWO windfarm site is 3.6 hours direct flight time away from the nearest guillemot breeding colony (Flamborough Head). A return trip would take 7.3 hours, not allowing for foraging. As for the Fair Isle example, travelling such distances is incompatible with successful breeding. On the basis of 5 feeds per day, the furthest away a bird could fly per trip to achieve this in 24 hours is 164km (i.e. a round trip of 328km), with no allowance for foraging time. Even if the bird spends a maximum of only 30 minutes foraging, this reduces the farthest distance to 147km.



- 126. On the basis of the above evidence, it can be stated with confidence that there are no major breeding colonies for guillemot within foraging range of the East Anglia TWO windfarm site, therefore it is reasonable to assume that individuals seen during the breeding season are nonbreeding and that they are largely sub-adult birds. Since sub-adult seabirds are known to remain in wintering areas, the number of sub-adult birds in the relevant population during the breeding season may be estimated as 43% (the proportion of the wintering BDMPS population that is immature, Furness 2015). This gives a breeding season population of 695,441 (BDMPS for the UK North Sea and English Channel, 1,617,306 x 43%).
- 127. Based on the average mortality for the species, a total of 97,362 birds would be expected to die each year from the sub-adult component of the winter BDMPS for this species (94,007; Furness 2015). The addition of a maximum of 19 birds predicted to die from construction disturbance and displacement would increase the mortality rate by 0.02%. (Use of the average mortality produces a conservative estimate of % change, as the mortality of birds less than 3 years old is higher than (survival is lower than) that of adult birds, Table 12.16). This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Thus, this highly precautionary assessment generates an effect of negligible magnitude.
- 128. The construction works are temporary and localised in nature and the magnitude of effect has been determined as negligible. As the species is of medium sensitivity to disturbance, the impact significance is **negligible**.

#### 12.6.1.2 Indirect Impacts Through Effects on Habitats and Prey Species

Indirect disturbance and displacement of birds may occur during the 129. construction phase if there are impacts on prey species and the habitats of prey species. These indirect effects include those resulting from the production of underwater noise (e.g. during piling) and the generation of suspended sediments (e.g. during preparation of the sea bed for foundations) that may alter the behaviour or availability of bird prey species. Underwater noise may cause fish and mobile invertebrates to avoid the construction area and also affect their physiology and behaviour. Suspended sediments may cause fish and mobile invertebrates to avoid the construction area and may smother and hide immobile benthic prey. These mechanisms may result in less prey being available within the construction area to foraging seabirds. Such potential effects on benthic invertebrates and fish have been assessed in Chapter 9 Benthic Ecology and Chapter 10 Fish and Shellfish Ecology and the conclusions of those assessments inform this assessment of indirect effects on ornithology receptors.



- 130. With regard to noise impacts on fish, Chapter 10 Fish and Shellfish Ecology discusses the potential impacts upon fish relevant to ornithology as prey species of the proposed East Anglia TWO project. For species such as herring, sprat and sandeel, which are the main prey items of seabirds such as gannet and auks, underwater noise impacts (physical injury or behavioural changes) during construction are considered to be minor or negligible (see Tables 10.22 and 10.23). With a minor or negligible impact on fish that are bird prey species, it is concluded that the indirect impact significance on seabirds occurring in or around the proposed East Anglia TWO project during the construction phase is similarly a *minor or negligible adverse* impact.
- 131. With regard to changes to the sea bed and to suspended sediment levels, Chapter 8 Marine Geology and Physical Processes and Chapter 9 Benthic Ecology discusses the nature of any change and impacts on the sea bed and benthic habitats. Such changes are considered to be temporary, small scale and highly localised (see Chapter 9 Benthic Ecology, section 10.6.2). The consequent indirect impact on fish through habitat loss is considered to be minor or negligible (see Table 10.25) for species such as herring, sprat and sandeel which are the main prey items of seabirds such as gannet and auks. With a minor or negligible impact on fish that are bird prey species, it is concluded that the indirect impact significance on seabirds occurring in or around the proposed East Anglia TWO project during the construction phase is similarly a minor or negligible adverse impact.

#### **12.6.2 Potential Impacts During Operation**

#### 12.6.2.1 Direct Disturbance and Displacement:

- 132. The presence of wind turbines and associated infrastructure and operational activities have the potential to directly disturb and displace birds from within and around the offshore development area. This is assessed as an indirect habitat loss, as it has the potential to reduce the area available to birds for feeding, loafing and moulting, and may result in reduction in survival rates of displaced birds. The presence of wind turbines, associated ancillary structures, vessel activity and factors such as the lighting of wind turbines could also attract certain species of birds.
- 133. As offshore windfarms are relatively new features in the marine environment, there is limited robust empirical evidence about the disturbance and displacement effects of the operational infrastructure in the long term, although the number of available studies of post-construction monitoring is increasing (e.g. JNCC 2015, Dierschke et al. 2016, Vallejo et al. 2017, MMO 2018). Dierschke et al. (2016) reviewed evidence from 20 operational offshore windfarms in European waters. They found strong avoidance by divers, gannet, great crested grebe, and fulmar; less consistent displacement by razorbill,



guillemot, little gull and sandwich tern; no evidence of any consistent response by kittiwake, common tern and Arctic tern, evidence of weak attraction to operating offshore windfarms for common gull, black-headed gull, great blackbacked gull, herring gull, lesser black-backed gull and red-breasted merganser, and strong attraction for shags and cormorants. Thaxter et al. (2018) also found no evidence of macro-avoidance of offshore windfarms by lesser blackbacked gulls. Displacement is apparently stronger when wind turbines are rotating. For cormorants and shags the presence of structures for roosting and drying plumage is a factor in attraction, while other species appear to benefit from increases in food abundance within operational offshore windfarms.

- During operation, the wind turbine array and offshore platforms will have lights 134. for air safety and navigational safety. There would be other lighting for personnel working at night, however these would not be as bright as air and navigational safety lighting. Air safety lights will be placed high on the wind turbine structures, and as a minimum on wind turbines at the periphery of the windfarm. Navigational lights for shipping will be placed lower on wind turbine structures and other offshore structures. A review of the potential effects of operational lighting on birds considered eight categories of potential effect on birds: disruption of photoperiod physiology; extension of daytime activity; phototaxis of seabirds; phototaxis of nocturnal migrant birds; ability of birds to use artificial light to feed at night or to feed on prey aggregating under artificial lights; increased predation risk for nocturnal migrant birds; birds better able to avoid collision when structures are illuminated; displacement of birds due to avoidance of artificial lights (Furness 2018). The available evidence suggests that lights on offshore wind turbines in European shelf seas are extremely unlikely to have any detectable effect on birds as a consequence of any of the processes listed above. The effects of operational lighting are therefore not assessed separately.
- 135. There is no empirical evidence that birds displaced from windfarms, or exposed to barrier effects, suffer increased mortality. Any mortality due to displacement would most likely be a result of increased densities of foraging birds in locations outside the affected area, resulting in increased competition for food. This would be unlikely for seabirds that have large areas of alternative habitat available, but would be more likely to affect seabirds with highly specialised habitat requirements that are limited in availability (Furness and Wade 2012; Bradbury et al. 2014). Impacts of displacement are also likely to be dependent on other environmental factors such as food supply, and are expected to be greater in years of low prey availability (e.g. as could result from unsustainably high fisheries pressures or effects of climatic changes on fish populations). Furthermore, modelling of the consequences of displacement for fitness of displaced birds suggests that even in the case of breeding seabirds that are



displaced on a daily basis, there is likely to be little or no impact on survival unless the offshore windfarm is close to the breeding colony (Searle et al. 2014, 2017).

- 136. The assessment below is based on a guidance note on displacement from the UK Statutory Nature Conservation Bodies (SNCB 2017).
- 137. Displacement is defined as 'a reduced number of birds occurring within or immediately adjacent to an offshore windfarm' (Furness et al. 2013) and involves birds present in the air and on the water (SNCB 2017). Birds that do not intend to utilise a windfarm area but would have previously flown through the area on the way to a feeding, resting or nesting area, and which either stop short or detour around a development, are subject to barrier effects (SNCB 2017).
- 138. Birds are considered to be most at risk from operational disturbance and displacement effects when they are resident in an area, for example during the breeding season or wintering season, as opposed to passage or migratory seasons. Birds that are resident in an area may regularly encounter and be displaced by an offshore windfarm for example during daily commuting trips to foraging areas from nest sites, whereas birds on passage may encounter (and potentially be displaced from) a particular offshore windfarm only once during a given migration journey.
- 139. For the purposes of assessment of displacement for resident birds, it is usually not possible to distinguish between displacement and barrier effects for example to define where individual birds may have intended to travel to, or beyond an offshore windfarm, even when tracking data are available. Therefore, in this assessment the effects of displacement and barrier effects on the key resident species are considered together.
- 140. The small risk of impact to migrating birds resulting from flying around rather than through, the wind turbine array of an offshore windfarm is considered a potential barrier effect, and have been scoped out of the assessment. Masden et al. (2010, 2012) and Speakman et al. (2009) calculated that the costs of one-off avoidances during migration were small, accounting for less than 2% of available fat reserves. Therefore, the impacts on birds that only migrate through the site (including seabirds, waders and waterbirds on passage) are considered negligible and these have been scoped out of detailed assessment.
- 141. Following installation of the offshore cable, the required operational and maintenance activities (in relation to the cable) may have short-term and localised disturbance and displacement impacts on birds using the offshore development area. However, disturbance from operational activities would be



temporary and localised, and is unlikely to result in detectable effects at either the local or regional population level. Therefore, no impact due to cable operation and maintenance is predicted.

- 142. The focus of this section is therefore on the disturbance and displacement of birds due to the presence and operation of wind turbines, other offshore infrastructure and any maintenance operations associated with them. The methodology presented in the SNCB Advice Note (SNCB 2017) recommends a matrix is presented for each key species showing bird losses at differing rates of displacement and mortality. This assessment uses the range of predicted losses, in association with the scientific evidence available from post-construction monitoring studies, to quantify the level of displacement and the potential losses as a consequence of the proposed project. These losses are then placed in the context of the relevant population (e.g. SPA or BDMPS) to determine the magnitude of effect.
- 143. In order to focus the assessment of disturbance and displacement, a screening exercise was undertaken to identify those species most likely to be at risk (Table 12.14), focussing on the main species described in the Baseline Offshore Ornithology Technical Report (Appendix 12.1). The species identified as at risk were then assessed within the biological seasons within which effects were potentially likely to occur. Any species with a low sensitivity to displacement, or recorded only in very small numbers within the East Anglia TWO windfarm site during the breeding and wintering seasons, was screened out of further assessment. Table 12.14 presents the general sensitivity to disturbance and displacement for each species. Displacement rates (based on observations of macro-avoidance, that is avoidance at the level of the whole windfarm rather than the wind turbine) are derived from a review of monitoring reports at constructed windfarms (Krijgsveld et al., 2011, Leopold et al., 2011, Vanermen et al. 2013, Walls et al., 2013, Mendel et al. 2014, Braasch et al. 2015, Skov et al. 2018, Cook et al. in press).

Species	Sensitivity to Disturbance and Displacement <sup>1</sup>	Screening Result (IN or OUT)	Season(s)	Rationale
Red-throated diver	Very High	IN	Autumn migration, Midwinter, Spring migration	Recorded regularly outside the breeding season and sensitive to disturbance and displacement
Black-throated diver	Very High	OUT	N/A	Recorded on only one occasion in baseline

#### Table 12.14 Operational Disturbance and Displacement Screening

### East Anglia TWO Offshore Windfarm Preliminary Environmental Information Report



Species	Sensitivity to Disturbance and Displacement <sup>1</sup>	Screening Result (IN or OUT)	Season(s)	Rationale
				surveys
Great northern diver	Very High	OUT	N/A	Recorded on only two occasions in baseline surveys
Fulmar	Considered Low in some studies, but possibly high according to Dierschke et al. (2016)	OUT	N/A	the species has a maximum habitat flexibility score of 1 in Furness and Wade (2012), suggesting it utilises a wide range of habitats over a large area.
Gannet	Considered Low in some studies, but possibly high according to Dierschke et al. (2016), and has a high macro- avoidance rate for windfarms	IN	Autumn and Spring migration	Potentially susceptible to displacement from wind turbines and abundant
Cormorant	Considered high in some studies but species is attracted to offshore windfarm structures	OUT	N/A	Recorded on only one occasion in baseline surveys
Shag	Medium	OUT	N/A	Recorded on only one occasion in baseline surveys
Great skua	Low	OUT	N/A	Recorded in low numbers during passage migration periods
Puffin	Medium	OUT	N/A	Recorded on in low numbers during only one baseline survey
Razorbill	Medium	IN	Year round	Potentially susceptible to displacement from wind turbines and



Species	Sensitivity to Disturbance and Displacement <sup>1</sup>	Screening Result (IN or OUT)	Season(s)	Rationale
				abundant
Guillemot	Medium	IN	Year round	Potentially susceptible to displacement from wind turbines and abundant
'Commic' tern <sup>2</sup>	Low	OUT	N/A	Recorded in low numbers and not very susceptible to displacement
Kittiwake	Low	OUT	N/A	No clear evidence of displacement from wind turbines
Black-headed gull	Low	OUT	N/A	No clear evidence of displacement from wind turbines
Little gull	Low	OUT	N/A	No clear evidence of displacement from wind turbines
Common gull	Low	OUT	N/A	No clear evidence of displacement from wind turbines
Lesser black- backed gull	Low	OUT	N/A	No clear evidence of displacement from wind turbines
Herring gull	Low	OUT	N/A	No clear evidence of displacement from wind turbines
Great black- backed gull	Low	OUT	N/A	No clear evidence of displacement from wind turbines

1. With reference to Garthe and Hüppop, 2004; Furness and Wade, 2012, Furness et al., 2013, Wade et al., 2016, Dierschke et al., 2016)

2 'Commic tern' is used where Arctic tern and common tern could not be distinguished at distance or from aerial survey images

144. The site population estimate used for each species to assess the displacement effects was the relevant seasonal peak mean (i.e. the highest mean value for the months within each season). The seasonal peaks were calculated as follows: first the density for each calendar month was calculated (as the



average of the density in each survey undertaken in that month), then the highest value from the months within each season extracted. As per SNCB (2017), for divers, the assessment used all data recorded within the 4km buffer, for all other species the assessment used all data recorded within the 2km buffer. Seasonal site population estimates for species included in the displacement assessment are included in **Table 12.15**.

- 145. Birds are considered to be most at risk from operational disturbance and displacement effects when they are resident (e.g. during the breeding season or wintering season). The small risk of impact to migrating birds is better considered in terms of barrier effects. However, SNCB (2017) suggests that migration periods should also be assessed using the matrix approach and this has been undertaken where appropriate.
- 146. For each species and season assessed, the predicted mortality due to displacement was determined and the impact of this assessed in terms of the change in the baseline mortality rate of the relevant population. It has been assumed that all age classes are equally at risk of displacement in proportion to their presence in the population.
- 147. As no information on seasonal population age structure is available from site data, it is necessary to calculate an average baseline mortality rate for all age classes for each species screened in for assessment. These were calculated using empirical information on the survival rates for each age class and their relative proportions in the population.
- 148. Demographic rates for each species from Horswill and Robinson (2015) were entered into a matrix population model. This was used to calculate the expected proportions in each age class. To obtain robust stable age class distributions for less well studied species (e.g. divers) the rates were modified to obtain a stable population size. Each age class survival rate was multiplied by its proportion and the total for all ages summed to give the average survival rate for all ages. Taking this value from 1 gives the average mortality rate. The demographic rates and the age class proportions and average mortality rates calculated from them are presented in *Table 12.16*.



Species	Area considered for displacement	Breeding	Migration-free breeding	Migration - autumn	Winter	Migration - spring	Non-breeding
Red-throated diver	Windfarm + 4km buffer	382.47	6.17	49.35	37.01	382.47	-
Gannet	Windfarm + 2km buffer	156.7	156.7	914.78	-	237.17	-
Razorbill	Windfarm + 2km buffer	287.99	287.99	55.06	148.23	262.58	-
Guillemot	Windfarm + 2km buffer	2126.02	2126.02	-	-	-	2020.14

#### Table 12.15 Seasonal Peak Mean Populations for Species Assessed for Displacement



Species	Parameter	Age cl	ass		Productivity	Average			
		0-1	1-2	2-3	3-4	5-6	Adult		mortanty
Red-	Survival	0.6	0.62				0.84	0.571	0.228
diver	Proportion in population	0.179	0.145				0.678		
Gannet	Survival	0.424	0.829	0.891	0.895		0.912	0.7	0.191
	Proportion in population	0.191	0.081	0.067	0.06		0.6		
Guillemot	Survival	0.56	0.792	0.917	0.939	0.939	0.939	0.672	0.14
	Proportion in population	0.168	0.091	0.069	0.062	0.056	0.552		
Razorbill	Survival	0.63	0.63	0.895	0.895		0.895	0.57	0.174
	Proportion in population	0.159	0.102	0.065	0.059		0.613		

 Table 12.16 Average Annual Mortality Across Age Classes Calculated Using Age-Specific

 Demographic Rates and Age Class Proportions

149. Natural England advice is that displacement effects estimated in different seasons should be combined to provide an annual effect for assessment which should then be assessed in relation to the largest of the component BDMPS populations. Natural England has acknowledged that summing impacts in this manner almost certainly over-estimates the number of individuals at risk through double counting (i.e. some individuals may potentially be present in more than one season) and assessing against the BDMPS almost certainly under-estimates the population from which they are drawn (which must be at least this size and is likely to be considerably larger as a consequence of turnover of individuals). However, at the present time there is no agreed alternative method for undertaking assessment of annual displacement and therefore the above approach is presented, albeit with the caveat that the results are anticipated to be highly precautionary.

#### 12.6.2.1.1 Red-throated Diver

- 150. Red-throated divers are considered to have a very high general sensitivity to disturbance and displacement and they are prone to avoiding disturbed areas (Garthe and Hüppop 2004; Petersen et al. 2006; Furness and Wade 2012; Percival 2014 Dierschke et al. 2017).
- 151. Displacement rates of 60% to 80% were reported for Egmond aan Zee offshore windfarm (OWEZ) (Leopold et al. 2011) and the review by Dierschke et al. (2016) also suggested a figure in this range.



- 152. Monitoring studies of red-throated divers at the Kentish Flats offshore windfarm found an observable shift of birds away from the wind turbines, particularly within 500m of the site (Percival 2010). Further pre-construction and post-construction abundance and distribution studies have provided displacement values for both the site footprint and within distance bands away from the site boundary. Percival (2014) reported that while displacement within the windfarm boundary was around 80% (compared to pre-construction), this declined to 10% at 1km from the windfarm and was 0% beyond 2km. A similar within windfarm reduction in density was reported at Thanet, but there was no detectable displacement beyond the windfarm boundary (Percival 2013).
- 153. A study of pre-construction and post-construction abundance and distribution of birds conducted at Horns Rev offshore windfarm, Denmark, found that red-throated divers avoided areas of sea that were apparently suitable (favoured habitat, suitable depth and abundant food sources) following the construction of an offshore windfarm, and that this effect remained for a period of three years (Peterson et al. 2006).
- 154. Modelling of data from pre-construction, construction and post-construction for the London Array Windfarm considered 1km buffers extending around the wind farm up to 15km. Red-throated diver density close to the wind farm was found to decline significantly between the pre-construction and construction periods; preliminary data from the post-construction period, however, may suggest that divers recolonised the windfarm and surrounding areas after construction had been completed (APEM 2016). It was noted that the densities of divers in the study area may vary to a large extent between years, and, as well as the presence of offshore wind farms and shipping activities, the total numbers of birds present as well as changes in other environmental conditions will influence the distribution of birds in a given year.
- 155. A large-scale and long-term analysis of the distribution of red-throated divers in the German North Sea found decreases in abundance detectable as far as about 16km from the closest operational offshore wind farm (Mendel et al. 2018).
- 156. The displacement matrices in *Table 12.17* through

**Table** 12.19 have been populated with data for red-throated diver during the autumn migration, nonbreeding and spring migration periods within the site and a 4km buffer in line with recommendations (SNCB 2017). It should be noted that the inclusion of all birds within the 4km buffer, to determine the total number of birds subject to 80% displacement, is considered precautionary, as in reality displacement has been demonstrated to decline with distance from a site.



158. The cells highlighted in green are for displacement rates of 60% to 80% (based on data from the Egmond an Zee Offshore Windpark (OWEZ) and a review by Dierschke et al. 2016, suggesting the actual rate lies between these two figures) and a precautionary range of displacement mortality rates between 1-5%. However, the windfarm site is not within foraging range of any breeding areas for red-throated divers, and the largest numbers were recorded during the spring migration period, at which time there is likely to be a turnover of individuals passing through the area, rather than a resident population. Thus, a given individual might only be displaced once from the windfarm, as opposed to being displaced multiple times if it was resident over the three month spring migration period. In reality therefore, mortality associated with displacement of red-throated divers from the East Anglia TWO windfarm site is very unlikely to be as high as 1-5%, and may well be zero (Dierschke et al. 2017). This conclusion is also supported by modelling of individual energy budgets (Topping and Petersen 2011). The assessment therefore considers a range of 1-5 % and applies a rate of 5% mortality to the population of birds within the windfarm and 4km buffer, with the aim of balancing precaution and evidence in the assessment.

#### 12.6.2.1.1.1 Autumn Migration

- 159. Within the range of 60-80% displacement and 1-5% mortality, the number of individual red-throated divers which could potentially suffer mortality as a consequence of displacement from the East Anglia TWO windfarm site during the autumn migration period has been estimated as 0-2 individuals (*Table 12.17*). The BDMPS for red-throated diver in autumn is 13,277 (Furness, 2015).
- 160. At the average baseline mortality rate for red-throated diver of 0.228 (*Table 12.16*) the number of individuals expected to die in the autumn BDMPS is 3,027 (13,277 x 0.228). The addition of a maximum of two birds increases the mortality rate by 0.07%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. In addition, this estimate of increase in mortality is considered highly precautionary as during this period birds would be passing through the site during migration, and the upper range of 5% mortality of displaced birds due to displacement seems very unlikely (Dierschke et al. 2017). Therefore, during the autumn migration period, the magnitude of effect is assessed as negligible, even on the basis of this highly precautionary approach. As the species is of high sensitivity to disturbance, the impact significance is **minor adverse**.

#### 12.6.2.1.1.2 Midwinter

161. Within the range of 60-80% displacement and 1-5% mortality, the number of individual red-throated divers which could potentially suffer mortality as a



consequence of displacement from the East Anglia TWO windfarm site during the midwinter period has been estimated as 0-1 individual (*Table 12.18*). The BDMPS for red-throated diver in winter is 10,177 (Furness 2015).

162. At the average baseline mortality rate for red-throated diver of 0.228, the number of individuals expected to die in the midwinter BDMPS is 2,320 (10,177 x 0.228). The addition of a maximum of one to this increases the mortality rate by 0.04%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the midwinter period, the magnitude of effect is assessed as negligible. As the species is of high sensitivity to disturbance, the impact significance is **minor adverse**.

#### 12.6.2.1.1.3 Spring Migration

- 163. Within the range of 60-80% displacement and 1-5% mortality, the number of individual red-throated divers which could potentially suffer mortality as a consequence of displacement from the East Anglia TWO windfarm site during the spring migration period has been estimated as 2-15 individuals (*Table 12.19*). The BDMPS for red-throated diver in spring is 13,277 (Furness, 2015).
- 164. At an average mortality rate of 0.228, the number of individuals expected to die in the spring BDMPS is 3,027 (13,277 x 0.228). The addition of a maximum of 15 to this increases the mortality rate by 0.5%. This magnitude of increase in mortality is considered highly unlikely as during this period birds would be passing through the site during migration. Therefore, during the spring migration period, the magnitude of effect is assessed as negligible. As the species is of high sensitivity to disturbance, the impact significance is **minor adverse**.

#### 12.6.2.1.1.4 Year Round

- 165. Considering the year-round effects, the maximum number of red-throated divers expected to die as a result of displacement from the East Anglia TWO windfarm site, at a displacement rate of 60-80% and mortality of 1-5%, would be 3-19 (adding the numbers predicted to be displaced during autumn migration, winter, and spring migration in *Table 12.17, Table 12.18* and *Table 12.19*, and noting that the totals in each table and the combined total are expressed to the nearest integer). The biogeographic red-throated diver population with connectivity to UK waters is 27,000 (Furness 2015).
- 166. At the average baseline mortality rate for red-throated diver of 0.228, the number of individuals expected to die over one year is 6,156 (27,000 x 0.228). The addition of 3- 19 to this increases the mortality rate by 0.05-0.3%. Most of this mortality is predicted during the spring migration period, when birds would be passing through the site rather than resident in the area. This magnitude of increase in mortality would not materially alter the background mortality of the



population and would be undetectable. Therefore, the magnitude of effect is assessed as negligible. As the species is of high sensitivity to disturbance, the impact significance is **minor adverse**.



## Table 12.17 Displacement Matrix for Red-throated Diver During the Autumn Migration Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Autumn migration		Mortality rate										
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	0	0	0	0	0	0	1	1	2	4	5
	20%	0	0	0	0	0	1	2	3	5	8	10
	30%	0	0	0	1	1	1	3	4	7	12	15
	40%	0	0	1	1	1	2	4	6	10	16	20
	50%	0	0	1	1	1	2	5	7	12	20	25
	60%	0	1	1	1	1	3	6	9	15	24	30
	70%	0	1	1	1	2	3	7	10	17	28	35
	80%	0	1	1	2	2	4	8	12	20	32	39
	90%	0	1	1	2	2	4	9	13	22	36	44
	100%	0	1	1	2	2	5	10	15	25	39	49



## Table 12.18 Displacement Matrix for Red-throated Diver During the Winter Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Winter		Mortality rate										
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	0	0	0	0	0	0	1	1	2	3	4
	20%	0	0	0	0	0	1	1	2	4	6	7
	30%	0	0	0	0	1	1	2	3	6	9	11
	40%	0	0	0	1	1	1	3	4	7	12	15
	50%	0	0	1	1	1	2	4	6	9	15	19
	60%	0	0	1	1	1	2	4	7	11	18	22
	70%	0	1	1	1	1	3	5	8	13	21	26
	80%	0	1	1	1	1	3	6	9	15	24	30
	90%	0	1	1	1	2	3	7	10	17	27	33
	100%	0	1	1	1	2	4	7	11	19	30	37

### SCOTTISHPOWER RENEWABLES

# Table 12.19 Displacement Matrix for Red-throated Diver During the Spring Migration Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Spring migration		Mortality rate										
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	0	1	1	2	2	4	8	11	19	31	38
	20%	1	2	2	3	4	8	15	23	38	61	76
	30%	1	2	3	5	6	11	23	34	57	92	115
	40%	2	3	5	6	8	15	31	46	76	122	153
	50%	2	4	6	8	10	19	38	57	96	153	191
	60%	2	5	7	9	11	23	46	69	115	184	229
	70%	3	5	8	11	13	27	54	80	134	214	268
	80%	3	6	9	12	15	31	61	92	153	245	306
	90%	3	7	10	14	17	34	69	103	172	275	344
	100%	4	8	11	15	19	38	76	115	191	306	382



#### 12.6.2.1.2Gannet

- 167. Gannets show a low level of sensitivity to ship and helicopter traffic (Garthe and Hüppop 2004, Furness and Wade 2012, Furness et al. 2013), but appear to be more sensitive to displacement from structures such as offshore wind turbines (Wade et al. 2016) and on this basis SNCB (2017) indicates that a detailed assessment of potential displacement should be carried out as standard.
- 168. Cook et al. (in press) review a number of studies of displacement of gannets from offshore windfarms. Where quantified, macro-avoidance rates (the % of birds taking action to avoid entering the wind turbine array) of 64% to 100% were reported. Some studies however reported no displacement response of gannets, possibly in areas where low densities of birds were present. Cook et al. (in press) recommended that the lowest of the quantified macro-avoidance rates, 64% for Egmond aan Zee offshore windfarm (Krijgsveld et al 2011) was appropriate for this species. A study of seabird flight behaviour at Thanet offshore windfarm, not included in the above review, found a macro-avoidance rate of 79.7% for gannets approaching within 3km of the windfarm (Skov et al. 2018).
- 169. Displacement effects for gannets for the East Anglia TWO windfarm site were assessed during the autumn and spring migration periods, based on respective peak mean populations of 915 and 237 individual birds (

**Table** 12.20) calculated for the windfarm site and a 2km buffer in line with recommendations within the SNCB guidance (SNCB 2017). The inclusion of all birds within the 2km buffer, to determine the total number of birds subject to displacement, is precautionary, as in reality the avoidance rate is likely to fall with distance from the site. This has been demonstrated in a recent study of gannet distribution in relation to the Greater Gabbard windfarm (APEM 2014).

171. Although gannets were recorded within the windfarm and a 2km buffer within the breeding season, and the site lies within potential foraging range of the nearest breeding colony at Bempton Cliffs, tracking data suggest that breeding adults from that colony make very little if any use of the East Anglia TWO windfarm site during the breeding season (Langston et al. 2013). Thus, it has been assumed that birds present during the breeding season are sub-adults or non-breeding adults, and any displacement of such non-breeding birds would not affect the Bempton Cliffs breeding population. Therefore, the displacement of gannets in summer is more appropriately assessed against the relevant BDMPS component of immature birds. However, the estimated number of birds that might die due to displacement in summer (60-80% displaced, 0-1% mortality of displaced birds) would be one bird so this impact is negligible at the population level.


172. Displacement matrices for gannets during the autumn and spring migration periods (calculated for the site and a 2km buffer) are presented in

Table 12.20 and

173. **Table** 12.21. For this species, the assessment considers predicted displacement rates of 60-80%, based on the recommendations of Cook et al. (*in press*) and also the findings of Skov et al. (2018) (see **paragraph 168** above). Mortality rates of displaced birds are assumed to be a maximum of 1%, as this species has high habitat flexibility (Furness and Wade 2012) indicating that displaced birds are predicted to readily find alternative habitats including foraging areas.

#### 12.6.2.1.2.1 Autumn Migration

175. Based on displacement rates of 60% to 80% and mortality rates of 0-1%, the maximum number of individual gannets which could potentially suffer mortality as a consequence of displacement during the autumn migration period has been estimated as 7 individuals (cells highlighted

#### *Table* 12.20).

177. The BDMPS for gannet in autumn is 456,298 (Furness 2015). At the average baseline mortality rate for gannet of 0.191 (the number of individuals expected to die in the autumn BDMPS is 87,153 (456,298 x 0.191). The addition of a maximum of 7 to this increases the mortality rate by 0.008%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the autumn migration period, the magnitude of effect is assessed as negligible. As the species is of low to medium sensitivity to disturbance, the impact significance is **negligible**.

#### 12.6.2.1.2.2 Spring Migration

- 178. Within the range of 60-80% displacement and 0-1% mortality, the maximum number of individual gannets which could potentially suffer mortality as a consequence of displacement during the spring migration period has been estimated as 2 individuals (*Table 12.16*).
- 179. The BDMPS for gannet in spring is 248,385 (Furness 2015).
- 180. At the average baseline mortality rate for gannet of 0.191 (

**Table** 12.22) the number of individuals expected to die in the spring BDMPS is 47,441 (248,385 x 0.191). The addition of a maximum of 2 to this increases the mortality rate by 0.004%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the spring migration period, the magnitude of



effect is assessed as negligible. As the species is of low to medium sensitivity to disturbance, the impact significance is negligible.

### 12.6.2.1.2.3 Year Round

182. Considering the year-round effects, the maximum number of gannets expected to die as a result of displacement from the East Anglia TWO windfarm site, at a displacement rate of 60-80% and mortality of 0-1%, would be 10 (7 during autumn migration, 2 during spring migration, and 1 during the breeding season,

*Table* 12.22). The biogeographic gannet population with connectivity to UK waters is 1,180,000 (Furness 2015).

184. At the average baseline mortality rate for gannet of 0.191 the number of individuals expected to die over one year is 225,380 (1,180,000 x 0.191). The addition of a maximum of 10 to this increases the mortality rate by 0.004%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, the magnitude of effect is assessed as negligible. As the species is of low to medium sensitivity to disturbance, the impact significance is **negligible**.

## SCOTTISHPOWER RENEWABLES

## Table 12.20 Displacement Matrix for Gannet During the Autumn Migration Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Autumn migration	Mortality rate											
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	1	2	3	4	5	9	18	27	46	73	91
	20%	2	4	5	7	9	18	37	55	91	146	183
	30%	3	5	8	11	14	27	55	82	137	220	274
	40%	4	7	11	15	18	37	73	110	183	293	366
	50%	5	9	14	18	23	46	91	137	229	366	457
	60%	5	11	16	22	27	55	110	165	274	439	549
	70%	6	13	19	26	32	64	128	192	320	512	640
	80%	7	15	22	29	37	73	146	220	366	585	732
	90%	8	16	25	33	41	82	165	247	412	659	823
	100%	9	18	27	37	46	91	183	274	457	732	915



## Table 12.21 Displacement Matrix for Gannet During the Spring Migration Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Spring	, T	Mortality r	Aortality rate											
migration														
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%		
Displacement	10%	0	0	1	1	1	2	5	7	12	19	24		
	20%	0	1	1	2	2	5	9	14	24	38	47		
	30%	1	1	2	3	4	7	14	21	36	57	71		
	40%	1	2	3	4	5	9	19	28	47	76	95		
	50%	1	2	4	5	6	12	24	36	59	95	119		
	60%	1	3	4	6	7	14	28	43	71	114	142		
	70%	2	3	5	7	8	17	33	50	83	133	166		
	80%	2	4	6	8	9	19	38	57	95	152	190		
	90%	2	4	6	9	11	21	43	64	107	171	213		
	100%	2	5	7	9	12	24	47	71	119	190	237		

## Table 12.22 Displacement Matrix for Gannet During the Breeding Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Breeding	Mortality rate											
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	0	0	0	1	1	2	3	5	8	13	16
	20%	0	1	1	1	2	3	6	9	16	25	31
	30%	0	1	1	2	2	5	9	14	24	38	47
	40%	1	1	2	3	3	6	13	19	31	50	63
	50%	1	2	2	3	4	8	16	24	39	63	78
	60%	1	2	3	4	5	9	19	28	47	75	94
	70%	1	2	3	4	5	11	22	33	55	88	110
	80%	1	3	4	5	6	13	25	38	63	100	125
	90%	1	3	4	6	7	14	28	42	71	113	141
	100%	2	3	5	6	8	16	31	47	78	125	157





### 12.6.2.1.3Auks (Razorbill and Guillemot)

- 185. Auks are considered to have medium sensitivities to disturbance and displacement from operational offshore windfarms based on available monitoring data and information on their responses to man-made disturbance, for example for ship and helicopter traffic (Garthe and Hüppop 2004; Schwemmer et al. 2011; Furness and Wade 2012; Furness et al. 2013; Bradbury et al. 2014; MMO 2018).
- 186. Available pre- and post-construction data for offshore windfarms have yielded variable results; they indicate that auks may be displaced to some extent by some windfarms, but displacement is partial and apparently negligible at others (Dierschke et al. 2016).
- 187. Common guillemots were displaced at Blighbank (Vanermen et al. 2012, 2014), were displaced only in a minority of surveys at two Dutch windfarms (OWEZ and PAWP; Leopold et al. 2011, Krijgsveld et al. 2011), but were not significantly displaced at Horns Rev (although the data suggest that slight displacement was probably occurring; Petersen et al. 2006) or Thornton Bank (Vanermen et al. 2012). Razorbills were displaced in one out of six surveys at two Dutch windfarms (OWEZ and PAWP; Leopold et al. 2011, Krijgsveld et al. 2011), but not at Horns Rev (Petersen et al. 2006) or Thornton Bank (Vanermen et al. 2012). At Blighbank, razorbills were found to be significantly displaced when considering the windfarm area and a 3km buffer, or the buffer alone (0.5-3km from the windfarm; Vanermen et al. 2014).
- 188. Following statutory guidance (SNCB 2017) the abundance estimates for each auk species for the windfarm and a 2km buffer for the most relevant biological periods have been placed into individual displacement matrices.
- 189. Each matrix displays displacement rates and mortality rates for each species. For the purpose of this assessment a displacement rate range of 30 to 70% and a mortality rate range of 1 to 10% are highlighted in each matrix (based on advice from Natural England), with the 70% / 10% combination representing a precautionary worst case scenario.
- 190. As noted previously, there are no breeding colonies for guillemot or razorbill within foraging range of the East Anglia TWO windfarm site. Therefore, it is reasonable to assume that individuals seen during the breeding season are nonbreeding individuals (e.g. immature birds). Since immature seabirds are known to remain in wintering areas, the number of immature birds in the relevant populations during the breeding season may be estimated as 43% of the total wintering BDMPS population for guillemot and razorbill (based on modelled age structures for these species populations in Furness, 2015). This



gives breeding season populations of non-breeding individuals of 695,441 guillemots (BDMPS for the UK North Sea and Channel, 1,617,306 x 43%), and 94,007 razorbills (BDMPS for the UK North Sea and Channel, 218,622 x 43%). For guillemot, there is only one defined nonbreeding season (August - February), while for razorbill there are three (August - October, November - December and January - March; *Table 12.10*. The number of birds which could potentially be displaced has been estimated for each species-specific relevant season.

### 12.6.2.1.4Razorbill

### 12.6.2.1.4.1 Autumn Migration

- 191. The estimated number of razorbills subject to mortality during the autumn migration period due to displacement from the East Anglia TWO windfarm site is between zero and four individuals (within the range of displacement/mortality of 30%/1% to 70%/10%, *Table* 12.23). The BDMPS for the UK North Sea and Channel is 591,874 (Furness 2015).
- 192. At the average baseline mortality rate for razorbill of 0.174 (*Table 12.16*) the number of individuals expected to die in the autumn migration period is 102,986 (591,874 x 0.174). The addition of a maximum of four individuals to this increases the mortality rate by < 0.01%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the autumn migration period, the magnitude of effect is assessed as negligible. As the species is of medium sensitivity to disturbance, the impact significance is **negligible**.

#### 12.6.2.1.4.2 Winter

193. The estimated number of razorbills subject to mortality during the winter period due to displacement from the East Anglia TWO windfarm site is between zero and 10 individuals (within the range of displacement/mortality of 30%/1% to 70%/10%,

*Table* 12.24). The BDMPS for the UK North Sea and Channel is 218,622 (Furness 2015).

195. At the average baseline mortality rate for razorbill of 0.174 the number of individuals expected to die in the winter period is 38,040 (218,622 x 0.174). The addition of a maximum of 10 individuals to this increases the mortality rate by 0.03%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the winter period, the magnitude of effect is assessed as negligible. As the species is of medium sensitivity to disturbance, the impact significance is **negligible**.



### 12.6.2.1.4.3 Spring Migration

196. The estimated number of razorbills subject to mortality during the spring migration period due to displacement from the East Anglia TWO windfarm site is between one and 18 individuals (within the range of displacement/mortality of 30%/1% to 70%/10%,

*Table* 12.25). The BDMPS for the UK North Sea and Channel is 591,874 (Furness 2015).

198. At the average baseline mortality rate for razorbill of 0.174 the number of individuals expected to die in the spring migration period is 102,986 (591,874 x 0.174). The addition of a maximum of 18 individuals to this increases the mortality rate by 0.02%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the spring migration period, the magnitude of effect is assessed as negligible. As the species is of medium sensitivity to disturbance, the impact significance is **negligible**.

### 12.6.2.1.4.4 Breeding Season

- 199. The estimated number of razorbills subject to mortality during the breeding period due to displacement from the East Anglia TWO windfarm site is between one and 20 individuals (from 30%/1% to 70%/10%, *Table* 12.26 and *Table 12.28*). The BDMPS is 94,007 non-breeding individuals (see *paragraph 190* above).
- 200. At the average baseline mortality rate for razorbill of 0.174, the number of individuals expected to die in the breeding season is 16,357 (94,007 x 0.174). The addition of a maximum of 20 to this increases the mortality rate by 0.12%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the nonbreeding migration period, the magnitude of effect is assessed as negligible. As the species is of medium sensitivity to disturbance, the impact significance is **negligible**.

## 12.6.2.1.4.5 Year Round

The estimated number of razorbills subject to displacement mortality throughout the year is between 2 and 53 individuals (summing the range of displacement/mortality of 30%/1% to 70%/10% from *Table 10.23* through *Table 10.26*).

201. At the average baseline mortality rate for razorbill of 0.174, the number of individuals expected to die from the largest BDMPS population throughout the year is 102,986 (591,874 x 0.174). The addition of a maximum of 53 individuals to this increases the mortality rate by 0.05%. This magnitude of increase in



mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the non-breeding season, the magnitude of effect is assessed as negligible. As the species is of medium sensitivity to disturbance, the impact significance is **negligible**.



## Table 12.23 Displacement Matrix for Razorbill During the Autumn Migration Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Autumn		Mortality r	rtality rate											
migration														
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%		
Displacement	10%	0	0	0	0	0	1	1	2	3	4	6		
	20%	0	0	0	0	1	1	2	3	6	9	11		
	30%	0	0	0	1	1	2	3	5	8	13	17		
	40%	0	0	1	1	1	2	4	7	11	18	22		
	50%	0	1	1	1	1	3	6	8	14	22	28		
	60%	0	1	1	1	2	3	7	10	17	26	33		
	70%	0	1	1	2	2	4	8	12	19	31	39		
	80%	0	1	1	2	2	4	9	13	22	35	44		
	90%	0	1	1	2	2	5	10	15	25	40	50		
	100%	1	1	2	2	3	6	11	17	28	44	55		

## Table 12.24 Displacement Matrix for Razorbill During the Winter Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Winter		Mortality rate										
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	0	0	0	1	1	1	3	4	7	12	15
	20%	0	1	1	1	1	3	6	9	15	24	30
	30%	0	1	1	2	2	4	9	13	22	36	44
	40%	1	1	2	2	3	6	12	18	30	47	59
	50%	1	1	2	3	4	7	15	22	37	59	74
	60%	1	2	3	4	4	9	18	27	44	71	89
	70%	1	2	3	4	5	10	21	31	52	83	104
	80%	1	2	4	5	6	12	24	36	59	95	119
	90%	1	3	4	5	7	13	27	40	67	107	133
	100%	1	3	4	6	7	15	30	44	74	119	148





## Table 12.25 Displacement Matrix for Razorbill During the Spring Migration Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Spring		Mortality r	rtality rate									
migration												
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	0	1	1	1	1	3	5	8	13	21	26
	20%	1	1	2	2	3	5	11	16	26	42	53
	30%	1	2	2	3	4	8	16	24	39	63	79
	40%	1	2	3	4	5	11	21	32	53	84	105
	50%	1	3	4	5	7	13	26	39	66	105	131
	60%	2	3	5	6	8	16	32	47	79	126	158
	70%	2	4	6	7	9	18	37	55	92	147	184
	80%	2	4	6	8	11	21	42	63	105	168	210
	90%	2	5	7	9	12	24	47	71	118	189	236
	100%	3	5	8	11	13	26	53	79	131	210	263



## Table 12.26 Displacement Matrix for Razorbill During the Breeding Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Breeding		Mortality r	ortality rate											
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%		
Displacement	10%	0	1	1	1	1	3	6	9	14	23	29		
	20%	1	1	2	2	3	6	12	17	29	46	58		
	30%	1	2	3	3	4	9	17	26	43	69	86		
	40%	1	2	3	5	6	12	23	35	58	92	115		
	50%	1	3	4	6	7	14	29	43	72	115	144		
	60%	2	3	5	7	9	17	35	52	86	138	173		
	70%	2	4	6	8	10	20	40	60	101	161	202		
	80%	2	5	7	9	12	23	46	69	115	184	230		
	90%	3	5	8	10	13	26	52	78	130	207	259		
	100%	3	6	9	12	14	29	58	86	144	230	288		



## 12.6.2.1.5 Guillemot

### 12.6.2.1.5.1 Non-Breeding

- 202. The estimated number of guillemots subject to mortality during the nonbreeding period due to displacement from the East Anglia TWO windfarm site is between six and 141 individuals (within the range of displacement/mortality of 30%/1% to 70%/10%, *Table 12.27*). The BDMPS for the UK North Sea and Channel is 1,617,306 (Furness 2015).
- 203. At the average baseline mortality rate for guillemot of 0.140 (*Table 12.16*) the number of individuals expected to die in the non-breeding season is 226,423 (1,617,306 x 0.140). The addition of a maximum of 141 individuals to this increases the mortality rate by 0.06%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the non-breeding season, the magnitude of effect is assessed as negligible. As the species is of medium sensitivity to disturbance, the impact significance is **negligible**.

#### 12.6.2.1.5.2 Breeding Season

- 204. The estimated number of guillemots subject to mortality during the breeding period due to displacement from the East Anglia TWO windfarm site is between six and 149 individuals (from 30%/1% to 70%/10%, *Table 12.28*). The BDMPS is 695,441 non-breeding individuals (see *paragraph 190* above).
- 205. At the average baseline mortality rate for guillemot of 0.140, the number of individuals expected to die in the breeding season is 97,362 (695,441 x 0.140). The addition of a maximum of 149 to this increases the mortality rate by 0.15%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the breeding period, the magnitude of effect is assessed as negligible. As the species is of medium sensitivity to disturbance, the impact significance is negligible.

#### 12.6.2.1.5.3 Year Round

- 206. The estimated number of guillemots subject to displacement mortality throughout the year is between 12 and 290 individuals (summing the range of displacement/mortality of 30%/1% to 70%/10% from **Table 12.27** and **Table 12.28**).
- 207. At the average baseline mortality rate for guillemot of 0.140, the number of individuals expected to die from the largest BDMPS population throughout the year is 226,423 (1,617,306 x 0.140). The addition of a maximum of 290 individuals to this increases the mortality rate by 0.13%. This magnitude of increase in mortality would not materially alter the background mortality of the population and would be undetectable. Therefore, during the breeding season,



the magnitude of effect is assessed as negligible. As the species is of medium sensitivity to disturbance, the impact significance is **negligible**.



## Table 12.27 Displacement Matrix for Guillemot During the Non-Breeding Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Non-breeding		Mortality r	Aortality rate									
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	2	4	6	8	10	20	40	61	101	162	202
	20%	4	8	12	16	20	40	81	121	202	323	404
	30%	6	12	18	24	30	61	121	182	303	485	606
	40%	8	16	24	32	40	81	162	242	404	646	808
	50%	10	20	30	40	51	101	202	303	505	808	1010
	60%	12	24	36	48	61	121	242	364	606	970	1212
	70%	14	28	42	57	71	141	283	424	707	1131	1414
	80%	16	32	48	65	81	162	323	485	808	1293	1616
	90%	18	36	55	73	91	182	364	545	909	1455	1818
	100%	20	40	61	81	101	202	404	606	1010	1616	2020



## Table 12.28 Displacement Matrix for Guillemot During the Breeding Period. The cells show the number of birds predicted to be die (rounded to the nearest integer) at a given rate of displacement and mortality.

Breeding Mortality rate												
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	2	4	6	9	11	21	43	64	106	170	213
	20%	4	9	13	17	21	43	85	128	213	340	425
	30%	6	13	19	26	32	64	128	191	319	510	638
	40%	9	17	26	34	43	85	170	255	425	680	850
	50%	11	21	32	43	53	106	213	319	532	850	1063
	60%	13	26	38	51	64	128	255	383	638	1020	1276
	70%	15	30	45	60	74	149	298	446	744	1191	1488
	80%	17	34	51	68	85	170	340	510	850	1361	1701
	90%	19	38	57	77	96	191	383	574	957	1531	1913
	100%	21	43	64	85	106	213	425	638	1063	1701	2126



### 12.6.2.2 Indirect Impacts Through Effects on Habitats and Prey Species

- 208. Indirect disturbance and displacement of birds may occur during the operational phase of the proposed East Anglia TWO project if there are impacts on prey species and the habitats of prey species. These indirect effects include those resulting from the production of underwater noise (e.g. the turning of the wind turbines), electro-magnetic fields (EMF) and the generation of suspended sediments (e.g. due to scour or maintenance activities) that may alter the behaviour or availability of bird prey species. Underwater noise and EMF may cause fish and mobile invertebrates to avoid the operational area and also affect their physiology and behaviour. Suspended sediments may cause fish and mobile invertebrates to avoid the operational area and may smother and hide immobile benthic prey. These mechanisms could result in less prey being available within the operational area to foraging seabirds. Changes in fish and invertebrate communities due to changes in presence of hard substrate (resulting in colonisation by epifauna) may also occur, and changes in fishing activity could influence the communities present.
- 209. With regard to noise impacts on fish, *Chapter 10 Fish and Shellfish Ecology* discusses the potential impacts upon fish relevant to ornithology as prey species. With regard to behavioural changes related to underwater noise impacts on fish during the operation of the proposed East Anglia TWO project, *Chapter 10 Fish and Shellfish Ecology, section 10.6.2.4* identifies that the sensitivity of fish and shellfish species to operational noise is considered to be low and the magnitude of effect negligible. It concludes a minor adverse impact on fish (see *Chapter 10 Fish and Shellfish Ecology, section 10.6.2.4*). With a negligible impact on fish that are bird prey species, it could be concluded that the indirect impact on seabirds occurring in or around the proposed TWO offshore development area during the operational phase is similarly a **negligible adverse impact**.
- 210. With regard to changes to the sea bed and to suspended sediment levels, *Chapter 9 Benthic Ecology* discusses the nature of any change and impact. It identifies that the small quantities of sediment released due to scour processes would rapidly settle within a few hundred metres of each wind turbine or cable protection structure. Therefore, the magnitude of the impact is likely to be negligible to low (see *Chapter 9 Benthic Ecology, section 9.6.2.3*) and that smothering due to increased suspended sediment during operation of the project would result in an impact of minor adverse significance. With a minor impact on benthic habitats and species, it could be concluded that the indirect impact on seabirds occurring in or around the East Anglia TWO windfarm site during the operational phase is similarly a **minor adverse impact**.



- 211. With regard to EMF effects, these are identified as highly localised with the majority of cables being buried to up to 5m depth, further reducing the effect of EMF (see *Chapter 9 Benthic Ecology, section 9.6.2.5*). The magnitude of impact is considered negligible on benthic invertebrates and low on fish. With a minor or negligible impact on invertebrates and fish, it could be concluded that the indirect impact on seabirds occurring in or around the offshore development area during the operational phase is similarly a **minor or negligible adverse** impact.
- 212. Very little is known about potential long-term changes in invertebrate and fish communities due to colonisation of hard substrate and changes in fishing pressures in the offshore development area. Whilst the impact of the colonisation of introduced hard substrate is seen as a minor adverse impact in terms of benthic ecology (as it is a change from the baseline conditions), the consequences for seabirds may be positive or negative locally, but are unlikely to be significant at a wider scale.

#### 12.6.2.3 Collision Risk

- 213. Birds flying through the wind turbine arrays of offshore windfarms may collide with rotor blades. This would result in fatality or injury to birds which fly through the East Anglia TWO windfarm site, during migration, whilst foraging for food, or commuting between breeding sites and foraging areas.
- 214. Collision Risk Modelling (CRM) has been used in this assessment to estimate the risk to birds associated with the East Anglia TWO windfarm site. CRM, using the Band model (Band 2012) has been used to produce predictions of mortality for particular species across biological seasons and annually. The approach to CRM is summarised here and further details are provided in *Appendix 12.1*.
- 215. The assessment is based on collision risk for each key seabird species from the Band CRM Option 2. This option uses generic estimates of flight height for each species based on the percentage of birds flying at Potential Collision Height (PCH) derived from data from a number of offshore windfarm sites, presented in Johnston et al. (2014a, 2014b).
- 216. Collision estimates for Band CRM option 1 are also included in *Appendix 12.1* (for information only as agreed through the ETG). This option uses flight height data for the East Anglia TWO study area. Following a review of data collection and analysis methods, the aerial survey contractors advised ScottishPower Renewables that the flight height estimates from baseline survey data were not reliable. Thus, these data have not been used in the assessment.



- 217. To take account of uncertainty in the parameters on which the model is based, CRM has been run as a stochastic simulation to incorporate uncertainty in seabird density, flight height (based on the generic dataset in Johnston et al. 2014a, 2014b), avoidance rates and nocturnal activity rates. This approach has been requested by NE because many of the CRM input parameters include both natural variation (e.g. seabird densities) and measurement error.
- 218. The most efficient method for incorporating uncertainty in multiple parameters is to generate multiple random values for each of the parameters from appropriate distributions and calculate the collision mortality for each combination of random values. To achieve this the Band model equations (Band 2012) were scripted in the R programming language (R Core Team 2016) to enable the Band model to be run as multiple simulations. Summary outputs calculated across the simulations can then be presented (e.g. median and confidence which incorporate the uncertainty in all the parameters intervals) simultaneously. However, as this approach has not been commonly used to date, and to assist readers to understand how variation in each of the parameters contributes to the overall variation, simulations were also conducted with only one of the parameters randomised at a time. In addition, a set of results obtained with no randomised parameters has been included, which are identical to those which are obtained using the Band (2012) spreadsheet.
- 219. The input parameters are provided in *Technical Appendix 12.1 Annex 3* and complete CRM results are provided in *Technical Appendix 12.1 Annexes 4* and 5. For both options 1 and 2 the following model runs were undertaken:
  - Uncertainty in seabird density, avoidance rate, flight height (Option 2 only) and nocturnal activity (gannet, kittiwake, large gulls only);
  - Uncertainty in seabird density only;
  - Uncertainty in collision avoidance rates only;
  - Uncertainty in flight height (Option 2) only;
  - Uncertainty in nocturnal activity only (gannet, kittiwake, large gulls only); and
  - No uncertainty in any parameter (i.e. a deterministic run)
- 220. The densities of birds in flight were calculated from the survey data. To obtain randomised values a nonparametric bootstrap resampling method was applied to each survey's dataset. This generated 1,000 resampled density estimates for each species on each survey. Density values were drawn at random from the resampled data. Runs which did not include uncertainty in density used the median density for each month (i.e. this was the median across all survey data for that month).



221. The avoidance rates used were as advised by Natural England and are set out in *Table 12.29* below. These were those recommended by the SNCBs (JNCC et al. 2014) following the review conducted by the British Trust for Ornithology (BTO) on behalf of Marine Scotland (Cook et al. 2014). When modelled with uncertainty the variations recommended in JNCC et al. (2014) were used.

Species	Avoidance Rate	Standard Deviation (SD)
Gannet	98.9%	0.2%
Kittiwake	98.9%	0.2%
Herring gull		
Lesser black-backed gull	99.5%	0.1%
Great black-backed gull		
Little gull		
Common gull	99.2%	0.2%
Black-headed gull		
All other species	98%	0.2%

Table	12 29	Avoidance	Rates	Used in (	RM
Iable	12.23	AVUIUAIICE	nates		21/101

- 222. It should be noted that further work on avoidance rates for offshore windfarms is underway. For example, a study on gannet behaviour in relation to offshore windfarms (APEM 2014) gathered evidence which suggests this species may exhibit a higher avoidance rate than the current recommended rate of 98.9%. This work, conducted during the autumn migration period, indicated an overall wind turbine avoidance of 100%, although a suitably precautionary rate of 99.5% was proposed (for the autumn period at least). Although this rate has not been applied to the estimates presented in this assessment, it indicates that gannet collision mortality estimated at 98.9% is likely to overestimate the risk for this species. The difference between the two rates sounds like a small change, but applying the higher avoidance rate of 99.5% would reduce predicted collision rates considerably, perhaps by as much as 50%. Indeed, as noted in Cook et al. (2014), all the recommended avoidance rates remain precautionary and thus the results presented in this assessment are worst case estimates.
- 223. A bird flight behaviour study has been conducted for the Offshore Renewables Joint Industry Programme (ORJIP). The study provides further evidence relating to the precautionary nature of current avoidance rates and other parameters used in windfarm assessment (Skov et al. 2018). Empirical avoidance rates were estimated for five seabird species as follows: 99.9% for gannet and herring gull, 99.8% for kittiwake and lesser black-backed gull, and 99.6% for



great black-backed gull. The predicted collision rate for gannet is consistent with the findings of the APEM (2014) study reported above, and all other empirical avoidance rates are higher than the currently recommended rates for CRM for a given species (*Table 12.29*). A thermal animal detection system of daylight and thermal imaging cameras recorded 6 collisions of birds with rotor blades during the course of the study. These were all gulls (not all identified to species), including one kittiwake.

- The nocturnal activity parameter used in the CRM defines the level of nocturnal 224. flight activity of each seabird species, expressed in relation to daytime flight activity levels. For example, a value of 50% for the nocturnal activity factor is appropriate for a species which is half as active at night as during the day. This factor is used to enable estimation of nocturnal collision risk from survey data collected during daylight, with the total collision risk the sum of those for day and night. The values typically used for each species were derived from reviews of seabird activity reported in Garthe and Hüppop (2004). This review ranked species from 1 to 5 (1 low, 5 high) for relative nocturnal activity, and these were subsequently modified for the purposes of CRM into 1 = 0% to 5 =100%. This approach was not anticipated by Garthe and Hüppop (2004), who considered that their 1 to 5 scores were simply categorical and were not intended to represent a scale of 0 to 100% of daytime activity (not least because the lowest score given was 1 and not 0). This is clear from their descriptions of the scores: for example, for score 1 'hardly any flight activity at nighť.
- 225. Recently however, a number of studies have deployed loggers on seabirds, and data from those studies can provide empirical evidence of the actual flight activity level. These studies indicate that the rates derived from Garthe and Hüppop (2004) almost certainly overestimate the levels of nocturnal activity in the species studied. For example, across four studies of gannet, nocturnal activity relative to daytime was reported as between 0% and 2%, across four studies of kittiwake nocturnal activity relative to daytime was reported as between 0% and 12% and in one study of lesser black-backed gull nocturnal activity relative to daytime was reported as 25%. These compare to the much higher values recommended by SNCBs for use in CRM of 25%, 50% and 50% for gannet, kittiwake and lesser black-backed gull respectively.
- 226. As the relative proportion of daytime to night-time varies considerably during the year at the UK's latitude, the effect of changes in the nocturnal activity factor for CRM outputs depends on the relative abundance of birds throughout the year. The extent of mortality reduction obtained by reducing the categorical score for five species (gannet, kittiwake, lesser black-backed gull, herring gull and great black-backed gull) by 1 (i.e. from 3 to 2 for kittiwake) has been



investigated previously (EATL 2015). This work revealed annual mortality estimate reductions of between 14.5% (lesser black-backed gull) and 27.7% (gannet). This indicated that current nocturnal activity factors based on arbitrary conversions of Garthe and Hüppop (2004) scores into percentages are over-estimated, and consequently CRM outputs are highly precautionary in this regard.

- 227. In the light of this, recent advice from Natural England has suggested that CRM should use upper and lower nocturnal activity rates of 0% and 25% for gannet and 25% and 50% for kittiwake, lesser black-backed gull, great black-backed gull and herring gull, rather than just the higher value as used previously.
- 228. In order to more accurately estimate nocturnal activity for gannet, a review of evidence from tracking studies has been undertaken (Furness et al. 2018). This recommended precautionary nocturnal flight activity rates for gannet in the breeding and nonbreeding seasons of 8% and 4% respectively. However, the actual average rates from the study were 7.1% and 2.3% respectively. Furthermore, the breeding season value was very heavily influenced by the results from the smallest study in the review, which was based on three tagged birds in Shetland (Garthe et al. 1999). That study yielded a nocturnal activity rate of 20.9% (compared to daytime) but the total duration of flight activity recorded was 215 hours, which was less than 3% of the > 8,000 hours covered by the remaining studies. If the average rate is calculated without this study a breeding season rate of 4.3% is obtained. This is considered to be more robust. Thus, for the proposed East Anglia TWO project, appropriate (and still precautionary) values for the breeding and nonbreeding seasons are considered to be 4.3% (SE 2.7%) and 2.3% (SE 0.4%). A similar review and analysis has been conducted for kittiwake (Furness et al. in prep.) which has identified values for the breeding and nonbreeding seasons respectively of 20% (SE 5%) and 17% (SE 1.5%). These values have considerably more merit, being based on empirical evidence, when compared with the categorical values which have been applied in CRM. Therefore, they have been used in the stochastic simulations for gannet and kittiwake in the current assessment. For the large gulls, uncertainty in nocturnal activity was modelled by selecting either 25% or 50% at random for each simulation. For all other species the previous nocturnal activity levels have been used (with no random variation in any run). In CRM runs which did not include uncertainty in nocturnal activity, the previously recommended values of 25% for gannet and 50% for kittiwake and the large gulls were used.
- 229. The CRM modelling results for all wind turbine scenarios are provided in *Technical Appendix 12.1 Annexes 4* and *5*. The assessment presented here uses the outputs for the worst case wind turbine layout for each species,



calculated using CRM option 2. As explained in paragraph 20, after the collision risk modelling was completed the design envelope was changed so that the maximum number of wind turbines increased from 67 to 75 for the 12MW scenario, and from 53 to 60 for the 15MW scenario. The results presented here are for previous scenarios of scenarios of 67 12MW, 53 15MW and 48 19MW turbines (see Appendix 12, Annex 3) rather than the scenarios presented in Table 12.2. The collision risk model has not been re-run for the updated scenarios because of time constraints, however an assessment of the updated parameters will be included within the ES. This model re-run will also incorporate the remaining three months of aerial survey data (see section 12.3.1).

- 230. As the numbers of turbines have been increased for the 12MW and 15MW scenarios then the collision risk for these scenarios would also increase, although changes are not considered likely to be of a magnitude that would change the conclusions of the assessment for the project alone and cumulative assessments as presented here.
- 231. Seasonal mortality predictions have been compared to the relevant BDMPS populations and the predicted increase in background mortality which could result has been estimated.
- 232. The full CRM results for the proposed project are presented in *Appendix 12.1*. The following sections provide a summary of the outputs for assessment, using the seasons defined in *Table 12.10*. Annual collision risk estimates for all species assessed are presented in *Table 12.30*. This table also includes an assessment of the species sensitivity to collision risk, based on available data on the % time spent flying at heights within the rotor diameter of offshore wind turbines, flight agility, the percentage of time flying, the extent of nocturnal flight activity and conservation importance (with reference to Garthe and Hüppop, 2004; Furness and Wade, 2012, Furness et al., 2013, Wade et al., 2016).
- 233. Several species had very low predicted annual collision risks at East Anglia TWO (i.e. worst case median prediction was below approximately one bird per year). These were red-throated diver, fulmar, great skua, black-headed gull, little gull, common gull and herring gull. As the magnitudes of predicted impact were so small, even for the worst case, no further assessment is considered necessary for these species (although additional outputs for these species are provided in *Technical Appendix 12.1*). The predicted annual collision risk for lesser black-backed gull was also very low, but this species was taken forward to assessment on a precautionary basis. The East Anglia TWO windfarm site is 37km from the Alde-Ore Estuary SPA at the nearest point, and within the mean maximum foraging range (141km, Thaxter et al. 2012). Thus, lesser black-



backed gulls breeding at the Alde-Ore Estuary SPA might forage within or pass through the development and be at risk of collision.

- 234. The seasonal collision estimates for lesser black-backed gull and species with higher estimated annual collisions (gannet, kittiwake, and great black-backed gull) are presented in *Table 12.31*. These species are considered in the assessment of collision risk.
- 235. The full stochastic results have been used for the following assessment of potential effects as these are considered to be the most robust figures, while still being precautionary.
- 236. Impacts during the non-breeding periods have been assessed in relation to the relevant BDMPS (Furness 2015). Where there is potential for impacts during the breeding season, these have been assessed in relation to reference populations calculated as described in the assessment for a given species.

Table 12.30 Annual Collision Risk Estimates.Values are the Median number of birds and 95%Confidence Intervals.For species screened in for Assessment, Shaded Cells Indicate the DesignOption with the Highest Estimated Collision Risk

Species (sensitivity to collision impacts)	Model run type	Annual 12MW	Annual 15MW	Annual 19MW		
Red throated diver	Full stochastic	0	0	0		
(Low)		(0-5.18)	(0-5.22)	(0-5.16)		
	Density only	0	0	0		
		(0-4.43)	(0-4.75)	(0-4.12)		
	Avoidance rate	0	0	0		
	only	(0-0)	(0-0)	(0-0)		
	Flight height only	0	0	0		
		(0-0)	(0-0)	(0-0)		
	Nocturnal activity	0	0	0		
	only	(0-0)	(0-0)	(0-0)		
	Deterministic	0	0	0		
		(0-0)	(0-0)	(0-0)		
Fulmar	Full stochastic	Full stochastic	Full stochastic 0	0.18	0.17	0.16
(Low)		(0-1.72)	(0-1.66)	(0-1.74)		
	Density only	0.2	0.2	0.2		
		(0-1.28)	(0-1.26)	(0-1.26)		



Species	Model run type	Annual 12MW	Annual 15MW	Annual 19MW	
(sensitivity to collision impacts)					
	Avoidance rate	0.21	0.2	0.2	
	only	(0.17-0.25)	(0.16-0.24)	(0.16-0.24)	
	Flight height only	0.18	0.18	0.19	
		(0.05-0.44)	(0.05-0.46)	0.05-0.45)	
	Nocturnal activity	0.2	0.2	0.2	
	only	(0.2-0.2)	(0.2-0.2)	(0.2-0.2)	
	Deterministic	0.2	0.2	0.2	
Croot Skup	Full stoppostic	0	0	0	
(Modium)	Full Stochastic	0	(0, 2, 24)	0	
(Medium)		(0-2.18)	(0-2.24)	(0-2.1)	
	Density only	0	0	0	
		(0-2.1)	(0-2.02)	(0-2.02)	
	Avoidance rate	0	0	0	
	oniy	(0-0)	(0-0)	(0-0)	
	Flight height only	0 (0-0)	0 (0-0)	0 (0-0)	
	Nocturnal activity only	0 (0-0)	0 (0-0)	0 (0-0)	
	Deterministic	0	0	0	
Gannet	Full stochastic	18.56	17.82	16.63	
(High)		(3.17-110.67	(2.5-102.75)	(2.6-106.77)	
	Density only	25.39	23.72	24.42	
		(7.07-69.86)	(6.54-64.38)	(7.02-65.64)	
	Avoidance rate	25.35	23.49	23.36	
	only	(17.54-39.95)	(16.11-33.11)	(16.15-33.19)	
	Flight height only	24.11	21.93	22.23	
		(6.21-56.84)	(6.18-53.8)	(5.24-53.52)	
	Nocturnal activity	25.65	23.68	23.68	
	only	(25.29-26.31)	(23.36-24.27)	(23.36-24.27)	
	Deterministic	25.7	23.71	23.71	



Species	Model run type	Annual 12MW	Annual 15MW	Annual 19MW
(sensitivity to collision impacts)				
Kittiwake	Full stochastic	25.79	25.25	25.87
(Medium)		(3.6-114.96)	(3.8-111.09)	(3.6-113.26)
	Density only	31.48	30.81	31.87
		(8.13-73.14)	(8.24-71.91)	(7.61-72.62)
	Avoidance rate	31.66	30.96	30.93
	only	(21.47-44.14)	(21.1-43.53)	(21.22-43.14)
	Flight height only	29.43	28.72	29.11
		(7.67-70.89)	(7.66-66.82)	(7.57-68.55)
	Nocturnal activity	31.78	31.23	31.28
	only	(30.42-33.45)	(29.91-32.97)	(29.92-32.87)
	Deterministic	31.82	31.31	31.31
Black-headed gull	Full stochastic	0.08	0.24	0.14
(Medium)		(0-13.29)	(0-13.18)	(0-12.66)
	Density only	1.3	0.86	0.86
		(0-8.78)	(0-8.37)	(0-8.37)
	Avoidance rate only	1	0.99	1.02
		(0.57-1.53)	(0.56-1.51)	(0.55-1.56)
	Flight height only	0.92	0.95	0.91
		(0.26-2.26)	(0.24-2.09)	(0.24-2.22)
	Nocturnal activity	1.02	1.01	1.01
	oniy	(1.02-1.02)	(1.01-1.01)	(1.01-1.01)
	Deterministic	1.02	1.01	1.01
Little gull	Full stochastic	0.5	0.5	0.47
(Low)		(0-4.39)	(0-5.03)	(0-4.72)
	Density only	0.65	0.64	0.64
		(0-3.52)	(0-3.57)	(0-3.57)
	Avoidance rate	0.64	0.66	0.65
	only	(0.37-1.01)	(0.36-1.03)	(0.39-1.02)



Species	Model run type	Annual 12MW	Annual 15MW	Annual 19MW
(sensitivity to collision impacts)				
	Flight height only	0.59	0.6	0.61
		(0.16-1.37)	(0.18-1.45)	(0.15-1.44)
	Nocturnal activity	0.66	0.67	0.67
	only	(0.65-0.65)	(0.67-0.67)	(0.67-0.67)
	Deterministic	0.66	0.75	0.67
Common gull	Full stochastic	0	0	0
(Medium)		(0-5.82)	(0-5.57)	(0-5.95)
	Density only	0	0	0
		(0-4.82)	(0-4.71)	(0-4.72)
	Avoidance rate	0	0	0
	oniy	(0-0)	(0-0)	(0-0)
	Flight height only	0	0	0
		(0-0)	(0-0)	(0-0)
	Nocturnal activity	0	0	0
	oniy	(0-0)	(0-0)	(0-0)
	Deterministic	0	0	0
Lesser black-	Full stochastic	0.48	0.43	0.49
(High)		(0-12.65)	(0-12.39)	(0-11.71)
	Density only	0.73	0.7	0.7
		(0-10.9)	(0-9.89)	(0-10.46)
	Avoidance rate	0.72	0.7	0.7
	oniy	(0.48-1.01)	(0.46-0.99)	(0.45-1.02)
	Flight height only	0.69	0.68	0.63
		(0.15-1.56)	(0.12-1.5)	(0.13-1.47)
	Nocturnal activity	0.73	0.7	0.7
	oniy	(0.66-0.73)	(0.63-0.7)	(0.63-0.7)
	Deterministic	0.73	0.7	0.7
Herring gull	Full stochastic	0	0	0



Species (sensitivity to collision impacts)	Model run type	Annual 12MW	Annual 15MW	Annual 19MW
(High)		(0-3.41)	(0-3.59)	(0-3.47)
	Density only	0	0	0
		(0-2.84)	(0-3.41)	(0-2.72)
	Avoidance rate	0	0	0
	only	(0-0)	(0-0)	(0-0)
	Flight height only	0	0	0
		(0-0)	(0-0)	(0-0)
	Nocturnal activity	0	0	0
	only	(0-0)	(0-0)	(0-0)
	Deterministic	0	0	0
Great black- backed gull (High)	Full stochastic	2.7	2.24	2.45
		(0-35.77)	(0-32.83)	(0-31.98)
	Density only	3.57	3.38	3.38
		(0-29.23)	(0-28.03)	(0-27.6)
	Avoidance rate	3.52	3.35	3.34
	only	(2.3-5.1)	(2.21-4.88)	(2.16-4.81)
	Flight height only	3.35	3.08	3.23
		(0.76-7.67)	(0.53-7.19)	(0.66-6.97)
	Nocturnal activity	3.1	3.38	2.94
	only	(3.1-3.57)	(2.94-3.38)	(2.94-3.38)
	Deterministic	3.57	3.38	3.38
* with reference to G al., 2016	arthe and Hüppop, 20	004; Furness and Wa	de, 2012, Furness et	al., 2013, Wade et



Species (worst case layout)	Model run type	Breeding season	Autumn migration	Non-breeding	Spring migration	Annual (worst case)
Gannet	Full Stochastic	8.83	8.58		1.15	18.56
(12 MW)		(2.15-21.8)	(1.02-57.74)		(0-17.14)	(3.17-110.67)
	Density only	10.3	13.47		1.62	25.39
		(4.73-21.8)	(2.34-35.77)		(0-12.29)	(7.07-69.86)
	Avoidance rate only	10.19	13.54		1.62	25.35
		(7.06-14.65)	(9.39-19.02)		(1.09-2.28)	(17.54-35.95)
	Flight height only	9.75	12.84		1.52	24.11
		(2.72-23.17)	(3.12-30.05)		(1.6-1.65)	(6.21-56.84)
	Nocturnal activity	10.27	13.76		1.62	25.65
	only	(10.08-10.71)	(13.61-13.95)		(1.6-1.65)	(25.29-26.31)
	Deterministic	10.3	13.78		1.62	25.7
Kittiwake	Full stochastic	13.61	2.86		9.32	25.79
(12MW)		(2.51-52.77)	(0-18.89)		(1.09-43.4)	(3.6-114.96)
	Density only	16.58	3.69		11.21	31.48
		((5.55-32.63)	(0-13.9)		(2.58-26.61)	(8.13-73.14)
	Avoidance rate only	16.62	3.71		11.33	31.66
		(11.32-23.07)	(2.46-5.2)		(7.69-15.87)	(21.47-44.14)

#### Table 12.31 Seasonal Collision Risk Estimates. Values are the Median Number of Birds and 95% Confidence Intervals



Species (worst case layout)	Model run type	Breeding season	Autumn migration	Non-breeding	Spring migration	Annual (worst case)
	Flight height only	15.36	3.44		10.63	29.43
		(3.97-37.3)	(0.87-8.06)		(2.83-25.53)	(7.67-70.89)
	Nocturnal activity	16.65	3.73		11.4	31.78
	only	(16.03-17.41)	(3.56-3.93)		(10.83-12.11)	(30.42-33.45)
	Deterministic	16.67	3.73		11.42	31.82
Lesser black-	Full stochastic	0.48	0	0	0	0.48
backed gull (12 MW)		(0-5.64)	(0-3.26)	(0-1.63)	(0-2.12)	(0-12.65)
	Density only	0.73	0	0	0	0.73
		(0-4.36)	(0-3.07)	(0-1.6)	(0-1.87)	(0-10.9)
	Avoidance rate only	0.72	0	0	0	0.72
		(0.48-1.01)	(0-0)	(0-0)	(0-0)	(0.48-1.01)
	Flight height only	0.69	0	0	0	0.69
		(0.15-1.56)	(0-0)	(0-0)	(0-0)	(0.15-1.56)
	Nocturnal activity	0.73	0	0	0	0.73
	only	(0.66-0.73)	(0-0)	(0-0)	(0-0)	(0.66-0.73)
	Deterministic	0.73	0	0	0	0.73
Great Black-backed	Full stochastic	2.25		0.45		2.7
gull (12 MW)		(0-17.6)		(0-18.17)		(0-35.8)



Species (worst case layout)	Model run type	Breeding season	Autumn migration	Non-breeding	Spring migration	Annual (worst case)
	Density only	2.84		0.73		3.57
		(0-12.93)		(0-16.3)		(0-29.23)
	Avoidance rate only	2.8		0.72		3.52
		(1.84-4.05)		(0.46-1.05)		(2.3-5.1)
	Flight height only	2.69		0.66		3.35
		(0.61-6.16)		(0.15-1.51)		(0.76-7.67)
	Nocturnal activity	2.56		0.54		3.1
	only	(2.56-2.84)		(0.54-0.73)		(3.1-3.57)
	Deterministic	2.84		0.73		3.57



### 12.6.2.3.1 Breeding Season Reference Populations for Collision Assessment

#### 12.6.2.3.1.1 Gannet

237. The nearest gannet breeding colony to the proposed development is Bempton Cliffs within the Flamborough and Filey Coast SPA. The SPA is 246km from the East Anglia TWO windfarm site at the nearest point (*Table 12.12*). This is outside the mean maximum foraging range of gannets, estimated as 229 km (Thaxter et al. 2012), the usual measure used to identify potential connectivity between a breeding seabird colony and foraging areas, although it is within the estimated maximum foraging range of 590 km. Tracking studies of gannets from Bempton Cliffs during 2010-2012 suggest very little if any use of the East Anglia TWO windfarm site during the breeding season (Langston et al. 2013).

#### 12.6.2.3.1.2 Kittiwake

The East Anglia TWO windfarm site is beyond the range of kittiwake from any 238. breeding colonies. It is therefore very unlikely that birds present during the breeding season are breeding. While RSPB's Future of the Atlantic Marine Environments (FAME) studies have shown some extremely long foraging trips for this species (as reported in various publications such as Fair Isle Bird Observatory annual reports) those extreme values tend to occur at colonies where food supply is extremely poor and breeding success is low (for example Orkney and Shetland). Daunt et al. (2002) point out that seabirds, as central place foragers, have an upper limit to their potential foraging range from the colony, set by time constraints. For example, they assess this limit to be 73km for kittiwake based on foraging flight speed and time required to catch food, based on observations of birds from the Isle of May. This means that kittiwakes would be unable to consistently travel more than 73km from the colony and provide enough food to keep chicks alive. Hamer et al. (1993) recorded kittiwake foraging ranges exceeding 40km in 1990 when sandeel stock biomass was very low and breeding success at the study colony in Shetland was 0.0 chicks per nest, but <5km in 98% of trips in 1991 when sandeel abundance was higher and breeding success was 0.98 chicks per nest. Kotzerka et al. (2010) reported a maximum foraging range of 59km, with a mean range of around 25km for a kittiwake colony in Alaska. Consequently, the breeding season impact on kittiwake has been assessed against a reference population estimated using the same approach as that for the displacement assessment (section 12.6.1.1). This is based on the observation that immature birds tend to remain in wintering areas. Thus, the number of immature birds in the relevant populations during the breeding season may be estimated as the proportion of the relevant BDMPS (the one immediately preceding the breeding season) which are sub-adults. This can be calculated as 47.3% of the spring BDMPS populations of kittiwake (Furness 2015). This yields a breeding season



population of nonbreeding kittiwake of 296,956 (Spring BDMPS for the UK North Sea and Channel, 627,816 x 47.3%).

### 12.6.2.3.1.3 Lesser black-backed gull

- 239. Lesser black-backed gulls breed at the Alde-Ore Estuary SPA which is within the 141km mean maximum foraging range (Thaxter et al. 2012) of this species from the East Anglia TWO windfarm site. Thus, there is potential for connectivity with the East Anglia TWO project during the breeding season.
- 240. In addition to the Alde-Ore colony, non-SPA colonies of lesser black-backed gulls located within foraging range of the East Anglia TWO windfarm site include rooftop nesting gulls in several towns in Suffolk and Norfolk. Potential connectivity with breeding colonies of lesser black-backed gulls in the Netherlands, within foraging range, was considered. This was ruled out however based on colour-ring and tracking studies which indicate that breeding lesser black-backed gulls from the Netherlands normally remain on the continental side of the North Sea. The recent assessment for Norfolk Vanguard (Norfolk Vanguard Ltd 2018) estimated a breeding reference population of 25,700 individuals of lesser black-backed gulls with potential connectivity to the proposed development. This comprised breeding adults and non-breeding adults and sub adults associated with the Alde-Ore SPA, and coastal and urban areas of Suffolk and Norfolk with, based on the JNCC Seabird Monitoring Programme (http://jncc.defra.gov.uk/smp/) and a survey of Suffolk breeding colonies in 2012 (Piotrowski 2012). For the purposes of this assessment, the estimated collision risk mortality at the East Anglia TWO windfarm site during the breeding season has been compared against this reference population.
- 241. Tracking data for lesser black-backed gulls breeding at the Alde-Ore Estuary SPA indicate that birds sometimes travel as far as the East Anglia TWO windfarm site, but the core foraging areas for this breeding colony do not overlap with the proposed project (Thaxter et al. 2015).

#### 12.6.2.3.1.4 Great Black-backed Gull

242. There are no breeding colonies for this species within foraging range of the East Anglia TWO windfarm site. Consequently, the breeding season impact on great black-backed gull has been assessed against a reference population estimated using the same approach as that for the displacement assessment (*section 12.6.1.1*). This is based on the observation that immature birds tend to remain in wintering areas. Thus, the number of immature birds in the relevant populations during the breeding season may be estimated as the proportion of the relevant BDMPS (the one immediately preceding the breeding season) which are sub-adults. Thus, the breeding season reference population can be calculated as 57.8% of the nonbreeding BDMPS populations of great black-backed gull (Furness 2015). This yields a breeding season population of



nonbreeding great black-backed gull of 52,829 (nonbreeding BDMPS for the UK North Sea and Channel, 91,399 x 57.8%).

- 12.6.2.3.2 Nonbreeding Season Reference Populations for Collision Assessment
- 243. As advised by Natural England, the non-breeding season reference populations were taken from Furness (2015).

### 12.6.2.3.3 Collision Impacts

- 244. The impacts of mortality caused by collisions on the populations are assessed in terms of the change in the baseline mortality rate which could result. It has been assumed that all age classes are equally at risk of collisions (i.e. in proportion to their presence in the population), therefore it is necessary to calculate an average baseline mortality rate for all age classes for each species assessed. These were calculated using the different survival rates for each age class and their relative proportions in the population.
- 245. The first step is to calculate an average survival rate. The demographic rates for each species were taken from reviews of the relevant literature (e.g. Horswill and Robinson, 2015) and recent examples of population modelling (e.g. EATL 2016). The rates were entered into a matrix population model to calculate the expected proportions in each age class. For each age class the survival rate was multiplied by its proportion and the total for all ages summed to give the average survival rate for all ages. Taking this value away from 1 gives the average mortality rate. The demographic rates and the age class proportions and average mortality rates calculated from them are presented in **Table 12.32**.

Species	Parameter	Age class					Productivity	Average
		0-1	1-2	2-3	3-4	Adult		montainty
Gannet	Survival	0.424	0.829	0.891	0.895	0.912	0.7	0.191
	Proportion in population	0.191	0.081	0.067	0.06	0.6		
	Survival	0.79	0.854	0.854	0.854	0.854	0.69	0.156
Kittiwake	Proportion in population	0.155	0.123	0.105	0.089	0.527		
Lesser	Survival	0.82	0.885	0.885	0.885	0.885	0.53	0.126
black- backed gull	Proportion in population	0.134	0.109	0.085	0.084	0.577		
Great	Survival	0.815	0.815	0.815	0.815	0.815	1.139	0.185

# Table 12.32 Average Annual Mortality Across Age Classes Calculated Using Age-SpecificDemographic Rates and Age Class Proportions



black-	Proportion in	0.194	0.156	0.126	0.102	0.422	
backed	population						
9							

- 246. The percentage increases in background mortality rates of seasonal and annual populations due to predicted collisions with the East Anglia TWO wind turbines are shown in *Table 12.33*.
- 247. The median and upper 95% confidence interval collision predictions for all species in all seasons and also summed across the year resulted in increases in background mortality of 0.4% or less. Increases of such small magnitude would not materially alter the background mortality of the population and would be undetectable. Therefore, the magnitude of effects due to collision mortality for gannet, kittiwake, lesser black-backed, gull and great black-backed gull are considered to be negligible, resulting in impact significances of **negligible to minor adverse** (based on species' sensitivities to collision risk in *Table 12.30*).
Table 12.33 Precautionary Estimates of Percentage Increases in the Background Mortality Rate of Seasonal and Annual Populations Due to Predicted Collisions (Option 2) Calculated with Stochasticity in Density, Avoidance Rate, Flight Height And Nocturnal Activity for Species Specific Worst Case Project Assumptions. Note that the Annual Mortalities Have Been Assessed Against Both the Biogeographic Populations and The Largest BDMPS (As Advised by Natural England) in Order to Indicate the Range of Likely Effects.

Species		Gannet (12 MW)			Kittiwake (12 MW)			Lesser black-backed gull			Great black-backed gull		
(Worst case layout)								(12 MW)			(12 MW)		
		Median	Lower c.i.	Upper c.i.	Median	Lower c.i.	Upper c.i.	Median	Lower c.i.	Upper c.i.	Median	Lower c.i.	Upper c.i.
Baseline average annual mortality		0.191			0.156	0.156		0.126			0.185		
Breeding Reference population		44,637			296,956	296,956		25,970			52,829		
	Seasonal mortality	8.83	2.15	35.79	13.61	2.51	52.77	0.48	0	5.64	2.95	0	11.14
	Increase in background mortality (%)	0.10%	0.03%	0.42%	0.03%	0.01%	0.11%	0.01%	0%	0.17%	0.02%	0%	0.18%
Autumn	Reference population	456,298	1	1	829,937	1	1	209,007	1	1	N/A	1	1
	Seasonal mortality	8.58	1.02	57.74	2.86	0	18.89	0	0	3.26			
	Increase in background mortality	0.01%	0.00%	0.07%	0%	0%	0.01%	0%	0%	0.01%			



Species		Gannet (12 MW)			Kittiwake (12 MW)		Lesser black-backed gull			Great black-backed gull			
(Worst case layout)								(12 MW)			(12 MW)		
	(%)												
Winter	Reference population	ce N/A on			N/A		39,316			91,399			
	Seasonal mortality							0	0	1.63	0.45	0	18.17
	Increase in background mortality (%)							0%	0%	0.03%	0%	0%	0.11%
Spring	Reference population	248,835			627,816		197,483			N/A			
	Seasonal mortality	1.15	0	17.14	9.32	1.09	43.4	0	0	2.12			
	Increase in background mortality (%)	0%	0%	0.04%	0.01%	0%	0.04%	0%	0%	0.01%			
Annual largest BDMPS	Reference population	456,298		829,937		209,007			91,399				
	Seasonal mortality	18.56	3.17	110.67	25.79	3.6	114.96	0.48	0	12.65	2.7	0	35.77



Species (Worst case layout)		Gannet (12 MW)		Kittiwak	Kittiwake (12 MW)		Lesser black-backed gull (12 MW)			Great black-backed gull (12 MW)			
	Increase in background mortality (%)	0.02%	0%	0.13%	0.02%	0%	0.09%	0%	0%	0.05%	0.02%	0%	0.21%
Annual biogeographic	Reference population	Reference         1,180,000         5,100,79000           population         5,100,79000         5,100,79000		9000 854,000		,000		235,000		·			
	Seasonal mortality	18.56	3.17	110.67	25.79	3.6	114.96	0.48	0	12.65	2.7	0	35.77
	Increase in background mortality (%)	0.01%	0%	0.05%	0%	0%	0.01%	0%	0%	0.01%	0.01%	0%	0.08%



### 12.6.3 Potential Impacts During Decommissioning

- 248. There are two potential impacts that may affect bird populations during the decommissioning phase of the proposed project that have been screened in. These are:
  - Disturbance / displacement; and
  - Indirect impacts through effects on habitats and prey species.
- 249. Any effects generated during the decommissioning phase of the proposed East Anglia TWO project are expected to be similar, or of reduced magnitude, to those generated during the construction phase, as certain activities such as piling would not be required. This is because it would generally involve a reverse of the construction phase through the removal of some structures and materials installed.
- 250. Potential impacts predicted during the decommissioning phase include those associated with disturbance and displacement and indirect effects on birds through effects on habitats and prey species.
- 251. It is anticipated that any future activities would be programmed in close consultation with the relevant statutory marine and nature conservation bodies, to allow any future guidance and best practice to be incorporated to minimise any potential impacts.

#### 12.6.3.1 Direct Disturbance and Displacement

- 252. Disturbance and displacement is likely to occur due to the presence of working vessels and crews and the movement, noise and light associated with these. Such activities have already been assessed for relevant bird species in the construction section above and have been found to be of negligible to minor negative magnitude.
- 253. Any impacts generated during the decommissioning phase of the proposed East Anglia TWO project are expected to be similar, but likely of reduced magnitude compared to those generated during the construction phase; therefore, the magnitude of effect is predicted to be negligible. This magnitude of impact on a range of species of low to high sensitivity to disturbance is of negligible to minor adverse significance.

### 12.6.3.2 Indirect Impacts Through Effects on Habitats and Prey Species

254. Indirect effects such as displacement of seabird prey species are likely to occur as structures are removed. Such activities have already been assessed for relevant bird species in the construction section above and have been found to be of negligible magnitude.



255. Any impacts generated during the decommissioning phase of the proposed project are expected to be similar, but likely of reduced magnitude compared to those generated during the construction phase; therefore, the magnitude of effect is predicted to be negligible. This magnitude of impact on a range of species of low to high sensitivity to disturbance is of negligible to minor adverse significance.

## **12.7Cumulative Impacts**

### **12.7.1 Screening for Cumulative Impacts**

- 256. The potential effects from the proposed East Anglia TWO project that were screened in for assessment for the project alone were further screened for the potential for cumulative effects with other projects (as defined in *section 12.7.2* below).
- 257. Two potential effects, operational displacement and collision risk, were screened in for cumulative assessment (*Table 12.34*).

Impact	Potential for cumulative impact	Data confidence <sup>1</sup>	Rationale
Construction			
Direct disturbance and displacement:	No	High	The likelihood that there would be a cumulative impact is low because the contribution from the proposed project is small and it is dependent on a temporal and spatial co- incidence of disturbance / displacement from other plans or projects.
Indirect impacts through effects on habitats and prey species	No	High	The likelihood that there would be a cumulative impact is low because the contribution from the proposed project is small and it is dependent on a temporal and spatial co- incidence of disturbance / displacement from other plans or projects.

### Table 12.34 Potential Cumulative Impacts

# East Anglia TWO Offshore Windfarm Preliminary Environmental Information Report



Impact	Potential for cumulative impact	Data confidence <sup>1</sup>	Rationale
Operation			
Direct disturbance and displacement:	Yes	High	There is a sufficient likelihood of a cumulative impact to justify a detailed, quantitative cumulative impact assessment.
Indirect impacts through effects on habitats and prey species	No	High	The likelihood that there would be a cumulative impact is low because the contribution from the proposed project is small
Collision risk	Yes	High	There is a sufficient likelihood of a cumulative impact to justify a detailed, quantitative cumulative impact assessment.
Decommissioning			
Direct disturbance and displacement:	No	High	The likelihood that there would be a cumulative impact is low because the contribution from the proposed project is small and it is dependent on a temporal and spatial co- incidence of disturbance / displacement from other plans or proposed projects.
Indirect impacts through effects on habitats and prey species	No	High	The likelihood that there would be a cumulative impact is low because the contribution from the proposed project is small and it is dependent on a temporal and spatial co- incidence of disturbance /



Impact	Potential for cumulative impact	Data confidence <sup>1</sup>	Rationale						
			displacement from other plans or projects.						
1. Indicates the degree of confidence; medium / low reflects lower confidence in older assessments which used variable methods.									

## **12.7.2 Projects Considered for Cumulative Impacts**

- 258. The classes of projects that could potentially be considered for the cumulative assessment of offshore ornithological receptors include:
  - Offshore windfarms;
  - Marine aggregate extraction;
  - Oil and gas exploration and extraction;
  - Sub-sea cables and pipelines; and
  - Commercial shipping.
- 259. Of these, only offshore windfarms are considered to have potential to contribute to cumulative operational displacement and collision risk, the effects screened in for cumulative assessment. Thus, the cumulative assessment is focused on offshore windfarms.
- 260. The identification of offshore windfarms to include in the cumulative assessment of offshore ornithological receptors has been based on:
  - Approved plans;
  - Constructed projects;
  - Approved but as yet unconstructed projects; and
  - Projects for which an application has been made, are currently under consideration and may be consented before the proposed East Anglia TWO project.
- 261. In addition, other 'foreseeable' projects are included: those for which an application has not been made but have been the subject of consultation by the developer, or those are listed in plans that have clear delivery mechanisms. For such projects, the absence of robust or relevant data could preclude a quantitative cumulative assessment being carried out.
- 262. The windfarms listed in *Table 12.35* have been assigned to Tiers following the approach proposed by Natural England and JNCC (Natural England, 2013) as follows:



- 1. Built and operational projects;
- 2. Projects under construction;
- 3. Consented;
- 4. Application submitted and not yet determined;
- 5. In planning (scoped), application not yet submitted; and,
- 6. Identified in Planning Inspectorate list of projects.



Project	Tier	Status	Development status	Distance from East Anglia TWO windfarm site (km)	Distance from offshore cable corridor (km)	Project data status	Included in CIA	Rationale
Greater Gabbard	1	Built and operational	Fully commissioned Aug 2013	13	20	Complete for the ornithology receptors being assessed	Yes	Included as an operational project that does not yet form part of the baseline.
Gunfleet Sands	1	Built and operational	Fully commissioned Jun 2010	67	52	Complete for the ornithology receptors being assessed	Yes	Included as an operational project that does not yet form part of the baseline.
Kentish Flats	1	Built and operational	Fully commissioned Dec 2005	96	85	Complete but limited quantitative species assessment	Yes	Operational for a sufficiently long time that its effects will have been incorporated in surveys but not yet in population responses
Lincs	1	Built and operational	Fully commissioned Sep 2013	143	128	Complete but limited quantitative species assessment	Yes	Included as an operational project that does not yet form part of the baseline.

### Table 12.35 Summary of Projects Considered for the CIA in Relation to Ornithology



Project	Tier	Status	Development status	Distance from East Anglia TWO windfarm site (km)	Distance from offshore cable corridor (km)	Project data status	Included in CIA	Rationale
London Array	1	Built and operational	Fully commissioned Apr 2013	58	52	Complete but limited quantitative species assessment	Yes	Included as an operational project that does not yet form part of the baseline.
Lynn and Inner Dowsing	1	Built and operational	Fully commissioned Mar 2009	144	129	Complete but limited quantitative species assessment	Yes	Included as an operational project that does not yet form part of the baseline.
Scroby Sands	1	Built and operational	Fully commissioned Dec 2004	41	33	Complete but limited quantitative species assessment	Yes	Operational for a sufficiently long time that its effects will have been incorporated in surveys but not yet in population responses
Sheringham Shoal	1	Built and operational	Fully commissioned Sep 2012	107	98	Complete but limited quantitative species assessment	Yes	Included as an operational project that does not yet form part of the baseline.
Beatrice (demonstrator)	1	Built and operational	Fully commissioned July 2007	721	713	Complete but limited quantitative	Yes	Due to be decommissioned



Project	Tier	Status	Development status	Distance from East Anglia TWO windfarm site (km)	Distance from offshore cable corridor (km)	Project data status	Included in CIA	Rationale
						species assessment		between 2024 and 2027
Thanet	1	Built and operational	Fully commissioned Sep 2010	74	78	Complete for the ornithology receptors being assessed	Yes	Included as an operational project that does not yet form part of the baseline.
Teesside	1	Built and operational	Fully commissioned Aug 2013	335	324	Complete but limited quantitative species assessment	Yes	Included as an operational project that does not yet form part of the baseline.
Westermost Rough	1	Built and operational	Fully commissioned May 2015	209	199	Complete for the ornithology receptors being assessed	Yes	Included as an operational project that does not yet form part of the baseline.
Humber Gateway	1	Built and operational	Fully commissioned May 2015	190	179	Complete but limited quantitative species assessment	Yes	Included as an operational project that does not yet form part of the baseline.



Project	Tier	Status	Development status	Distance from East Anglia TWO windfarm site (km)	Distance from offshore cable corridor (km)	Project data status	Included in CIA	Rationale
Galloper	1	Built and operational	Fully commissioned March 2018	7	17	Complete for the ornithology receptors being assessed	Yes	Included as an operational project that does not yet form part of the baseline.
Dudgeon	1	Built and operational	Fully commissioned November 2017	110	103	Complete but limited quantitative species assessment	Yes	Included as an operational project that does not yet form part of the baseline.
Race Bank	1	Built and operational	Fully commissioned February 2018	131	121	Complete but limited quantitative species assessment	Yes	Included as an operational project that does not yet form part of the baseline.
Beatrice	2	Under construction	Consent Mar 2014. Construction commenced Jan 2017	727	719	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
East Anglia ONE	2	Under	Consent Jun 2014, offshore	11	19	Complete for the	Yes	Included as a consented project that does not yet



Project	Tier	Status	Development status	Distance from East Anglia TWO windfarm site (km)	Distance from offshore cable corridor (km)	Project data status	Included in CIA	Rationale
		construction	construction due to commence August 2018			ornithology receptors being assessed		form part of the baseline.
EOWDC (Aberdeen OWF)	2	Under construction	Consent August 2014, offshore construction commenced April 2018	605	598	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Hornsea Project 1	2	Under construction	Consent Dec 2014, offshore construction commenced January 2018	167	165	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Rampion	2	Under construction	Consent Aug 2014. Construction commenced Apr 2017 (expected to be commissioned 2018)	221	207	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Blyth (NaREC Demonstration)	3	Consented	Consent Nov 2013, no construction	386	377	Complete but limited	Yes	Included as a consented project that does not yet



Project	Tier	Status	Development status	Distance from East Anglia TWO windfarm site (km)	Distance from offshore cable corridor (km)	Project data status	Included in CIA	Rationale
			start date			quantitative species assessment		form part of the baseline.
Dogger Bank Creyke Beck A and B	3	Consented	Consent Feb 2015, no construction start date.	261	258	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Inch Cape	3	Consented	Consent Sep 2014, no construction start date, revised application submitted August 2018	534	526	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Neart na Gaoithe	3	Consented	Consent Oct 2014, no construction start date, revised application with fewer wind turbines (max 54) submitted August 2018	516	507	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Firth of Forth Alpha and Bravo	3	Consented	Consent Oct 2014, no construction	525	517	Complete for the	Yes	Included as a consented project that does not yet



Project	Tier	Status	Development status	Distance from East Anglia TWO windfarm site (km)	Distance from offshore cable corridor (km)	Project data status	Included in CIA	Rationale
			start date, revised application submitted Sept 2018			ornithology receptors being assessed		form part of the baseline.
Moray East Offshore Windfarm	3	Consented	Consent Mar 2014, no construction start date	667	659	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Dogger Bank Teesside A and B (now Sofia)	3	Consented	Consent Aug 2015, no construction start date	280	278	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Hornsea Project 2	3	Consented	Consent Aug 2016, no construction start date	172	168	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Triton Knoll	3	Consented	Consent Jul 2013, no construction	144	135	Complete for the	Yes	Included as a consented project that does not yet



Project	Tier	Status	Development status	Distance from East Anglia TWO windfarm site (km)	Distance from offshore cable corridor (km)	Project data status	Included in CIA	Rationale
			start date, Non- material variation submitted Feb 2018			ornithology receptors being assessed		form part of the baseline.
East Anglia THREE	3	Consented	Consent Aug 2017. No construction start date	45	45	Complete for the ornithology receptors being assessed	Yes	Included as a consented project that does not yet form part of the baseline.
Hornsea Project 3	4	Application accepted	Pre-examination	158	156	Complete for the ornithology receptors being assessed	Yes	Included as a foreseeable project
Thanet Extension	4	Application accepted	Pre-examination	74	78	Complete for the ornithology receptors being assessed	Yes	Included as a foreseeable project
Norfolk Vanguard	4	Application accepted	Pre-examination	56	55	Complete for the	Yes	Included as a foreseeable project



Project	Tier	Status	Development status	Distance from East Anglia TWO windfarm site (km)	Distance from offshore cable corridor (km)	Project data status	Included in CIA	Rationale
						ornithology receptors being assessed		
East Anglia ONE North	5	Pre-planning application	Submission expected Q1 2019	10	10	ES not yet available	Yes	In the absence of final data, the outputs from the PEIR have been included.
Norfolk Boreas	5	Pre-planning application	Submission expectedQ2 2019	73	72	Not yet available	Yes	In the absence of final data, the outputs of the PEIR have been included.
Hornsea Project 4	5	Pre-planning application		175	170	Not yet available	Yes	In the absence of data, the inclusion of this project is only on a qualitative basis.



263. The level of data available and the ease with which impacts can be combined across the windfarms is quite variable, reflecting the availability of relevant data for older projects and the approach to assessment taken. Wherever possible the cumulative assessment is quantitative (i.e. where data in an appropriate format have been obtained). Where this has not been possible (e.g. for older projects) a qualitative assessment has been undertaken.

### **12.7.3 Cumulative Assessment of Operational Displacement**

- 264. The species assessed for project alone operational displacement impacts (and the relevant seasons) were red-throated diver (autumn, winter, spring), gannet (autumn, spring), guillemot (breeding, nonbreeding) and razorbill (breeding, autumn, winter, spring).
- 265. A review of the BDMPS regions for each species indicated that for gannet, guillemot, and razorbill, all the windfarms identified for inclusion in the CIA in **Table 12.35** have the potential to contribute a cumulative effect. For red-throated diver, the BDMPS is the southwest North Sea. Thus windfarms located in the north-west North Sea (all offshore windfarms located from the Northumbria coast northwards) and in the English Channel were not considered likely to contribute to a cumulative displacement effect for this species. In addition, as the species tends to be found in estuarine and near-shore shallow waters during the non-breeding season, offshore wind farms further from the coast (Hornsea, Dogger Bank) were also excluded.

### 12.7.3.1 Red-throated Diver

- 266. The project alone assessment concluded that during the midwinter period when divers are most at risk of impacts due to displacement, the maximum number of individuals at risk of mortality due to displacement was sufficiently small (0-1 birds) that there was no risk of a significant impact. Although a higher number was considered to be at risk of displacement-caused mortality during spring migration (2-15), this includes birds passing through the site for a brief period on migration and therefore the consequences of displacement are minimal and no significant project alone effects were predicted.
- 267. The recent Environmental Statement for Norfolk Vanguard Offshore Windfarm (Norfolk Vanguard Ltd 2018) and the PEIR for Norfolk Boreas (Norfolk Boreas Ltd 2018) include estimates of potential displacement mortality for red-throated divers from windfarms in the southern North Sea. These were based on a review of information in environmental statements. Windfarms at which wind turbines were installed before or during 2012 were considered to form part of the baseline, since any displacement effect at these sites would have occurred prior to the surveys for Norfolk Vanguard and Norfolk Boreas, and therefore any modifications in red-throated diver distribution as a result of displacement would



be fully reflected in the baseline data. The combined total and the windfarm sites contributing to this total is shown in *Table 12.36*.

- 268. As baseline surveys for the proposed East Anglia TWO project were carried out between November 2015 and August 2018 (*Appendix 12.1*), a similar argument can be applied as for Norfolk Vanguard, that displacement effects of windfarms with wind turbines installed before or during 2012 would be taken account of in baseline surveys. None of the other windfarms included in the cumulative assessment for Norfolk Vanguard has been completed in this timescale, so the cumulative total is applied to this assessment.
- 269. The predicted cumulative displacement mortality for red-throated divers from offshore windfarms in the southern North Sea, assuming a maximum of 80% displacement from the windfarms and a 4km buffer, and 1-5% mortality of displaced birds, is between 30 and 150.5 birds per year (*Table 12.36*).
- 270. The largest BDMPS for red-throated diver is 13,277 during spring and autumn migration (Furness 2015). At the average baseline mortality rate for red-throated diver of 0.228 (*Table 12.16*) the number of individuals expected to die is 3,027 (13,277 x 0.228). The addition of 30-150.5 to this would increase the mortality rate by 1-5%. The biogeographic population for red-throated diver with connectivity to UK waters is 27,000 (Furness 2015). At the average baseline mortality rate for red-throated diver of 0.228 the number of individuals expected to die is 6,156 (27,000 x 0.228). The addition of 30-150.5 to this would increase the mortality rate by 0.5-2.4%.
- 271. The cumulative red-throated diver displacement mortality total combines several sources of precaution:
  - Each windfarm assessment has assumed that all birds within 4km of the windfarm lease boundary are potentially affected to the same extent, whereas there is evidence that displacement declines with distance from windfarm boundaries and in some cases has been reported as zero by 2km;
  - It includes an unknown degree of double counting across seasons since some individuals will be present within more than one season;
  - There is some overlap between survey areas for different sites (the Norfolk Vanguard East 4km buffer includes part of the East Anglia THREE windfarm and 4km buffer and vice versa) so including both sites double counts birds in the overlapping area;
  - Displacement mortality may be less than 1-5% and could be as low as zero; and
  - More than half the total annual mortality (76%) is predicted to occur during the autumn and spring migration periods when the potential consequences



of displacement are expected to be much lower due to the brief duration that birds spend in the area at this time.

- 272. Looking at the winter period, the BDMPS is 10,177 (Furness 2015). At the average baseline mortality rate for red-throated diver of 0.228 the number of individuals expected to die is 2,320 (10,177 x 0.228). The addition of a maximum of 8-28 birds to this would increase the mortality rate by 0.3-1.2%.
- Generally based on findings from population viability analyses for bird species, 273. it would be considered that increases in mortality rates of less than 1% would be undetectable in terms of changes in population size, whereas above 1% there could be detectable effects. Using a range of mortality of 1-5% for displaced birds predicts changes in population mortality rates which are likely to be undetectable at the lower end and may be detectable at the upper end of the range. A review of the possible behavioural, energetic and demographic effects of displacement on red-throated divers (Dierschke et al. 2017) considered that because they use a range of marine habitats and take a variety of prey species, and are generally highly mobile in winter, increased density of birds in area of sea due to displacement from offshore wind farms would be unlikely to limit prev intake, both for displaced individuals and those already present in areas to which displaced birds move. Thus there might not be any effects of displacement on energy budgets, body condition and therefore mortality rates of displaced birds, unless prey abundance is depleted over extensive areas due to environmental conditions such as adverse weather. This was qualified with the evidence that red-throated divers do show a strong behavioural avoidance to anthropogenic disturbance which is likely to reflect a hormonal stress response and could affect their ability to forage normally for a period after displacement, and that some individuals are site faithful during the non-breeding season and if displaced might suffer to a greater extent than more mobile individuals.
- 274. A further source of precaution is that the assessment methodology makes no allowance for the fact that wind turbine densities (and hence the negative stimulus to which the birds respond) within the built windfarms will be much lower than the worst case designs on which the projects were consented. For example, East Anglia ONE was originally assessed on the basis of 333 wind turbines, reduced to 240 for consent and currently being constructed with 102. Thus, the final windfarm will have less than one third the original number of proposed (and assessed) wind turbines. Similar reductions are likely for other consented windfarms which have not yet been built. This is likely to further reduce the magnitude of displacement.
- 275. Given the various additive sources of precaution in this assessment, there is a very high likelihood that cumulative displacement would be lower than the worst



case totals presented here, resulting in increases in background mortality below 1%, and thus the magnitude of cumulative displacement is assessed as negligible. Therefore, as the species is of high sensitivity to disturbance, the cumulative impact significance would be **minor adverse**.



Offshore Windfarm(s)	No. red-throat	ed divers at r	ment mortality*	Source of information	
	Autumn migration	Winter	Spring migration	Annual	
East Anglia One, East Anglia THREE, Norfolk Vanguard, Galloper and Thanet Extension	0-6	3-19	6-27	12-59.5	Norfolk Vanguard Ltd (2018); Norfolk Boreas Ltd (2018)
East Anglia TWO	0-2	0-1	3-15	4-19	Table 12.17, Table 12.18, Table 12.19
East Anglia ONE North	0-1	1-3	3-14	3-17	East Anglia ONE North Ltd (2019)
Norfolk Boreas	0-1	1-5	10-49	11-55	Norfolk Boreas Ltd (2018)
Totals	2-10	8-28	22-105	30-150.5	

## Table 12.36 Cumulative Numbers of Red-throated Divers at Risk of Displacement Mortality from Offshore Windfarms in the North Sea

\* For the windfarm and a 4km buffer, assuming a maximum of 80% displacement and 1-5% mortality of displaced birds. Numbers for East Anglia TWO and East Anglia ONE North are rounded to the nearest integer, so annual totals may appear to be less or more than the sum of seasonal totals.



## 12.7.3.2 Gannet

276. The East Anglia TWO windfarm site is located beyond the mean maximum foraging range of gannets from breeding colonies in the North Sea. Therefore, displacement risk is primarily of concern outside the breeding season. There is evidence that gannets avoid flying through windfarms (Krijgsveld et al. 2011; Skov et al. 2018, Cook et al. in press). If this prevents them accessing important foraging areas this could have an impact on displaced individuals. However, for the reasons set out below, the potential for the proposed East Anglia TWO project to contribute to a cumulative effect such as this is considered to be very unlikely. The period when gannet displacement is of potential concern is during autumn migration. At this time, very large numbers of gannets are migrating from breeding colonies in Northern Europe to wintering areas farther south, predominantly off the coast of West Africa (Kubetzki et al. 2009; Furness et al. 2018a). Thus, displacement due to windfarms in the North Sea is trivial when compared with the range over which individuals of this species travel (Garthe et al. 2012, see also Masden et al. 2010, 2012). Furthermore, gannets are considered to be highly flexible in their foraging requirements, and exclusion from windfarms in the southern North Sea, on the basis of the low overall numbers of seabirds present, is very unlikely to represent a loss of any importance. Consequently, the potential for the proposed East Anglia TWO project to contribute to a significant cumulative displacement effect on gannets during migration is considered to be negligible and the impact significance of cumulative displacement on a receptor of low to medium sensitivity is negligible.

### 12.7.3.3 Razorbill

- 277. The East Anglia TWO windfarm site is located beyond the mean maximum foraging range of any razorbill breeding colonies (see *section 12.6.2.1.4*). Outside the breeding season razorbills migrate southwards from their breeding colonies. Large numbers are found in the North Sea throughout the non-breeding seasons (the spring and autumn migration periods and winter, between August and March; Furness 2015).
- 278. The annual total of razorbills at risk of displacement from the East Anglia TWO windfarm site is estimated as 754 individuals (summing the seasonal peak means on the East Anglia TWO windfarm site (and 2km buffer) for the migration-free breeding, autumn migration, winter, and spring migration periods; *Table 12.15*).
- 279. The recent cumulative assessments for offshore ornithology for East Anglia THREE (EATL 2016) and the Norfolk Vanguard project (Norfolk Vanguard Limited 2018) include cumulative estimates of the total numbers of razorbills at risk of displacement from other offshore windfarms in the North Sea. These



estimates are included in *Table 12.37*, along with updates for projects where new information has become available since the publication of Norfolk Vanguard Limited (2018). These totals omit windfarms for which no data are available (Gunfleet Sands, Kentish Flats, Lynn and Inner Dowsing, Scroby Sands), but they are also likely to over-estimate the numbers present due to the precautionary use of seasonal peak numbers at each site rather than average numbers, which is likely to lead to double counting as birds move through the North Sea.

- 280. The estimated annual cumulative total of razorbills at risk of displacement from windfarms in the North Sea is 98,927 individuals (*Table 12.37*). Considering a range of displacement of 30-70%, and mortality of displaced individuals from 1-10%, based on advice from Natural England, the estimated number of razorbills subject to mortality from displacement throughout the year is between 297 and 6,925 (*Table 12.38*).
- 281. This is a large range so the assessment considers the most realistic value within this range.
- 282. Post-construction monitoring of auks at offshore windfarms has found evidence of avoidance behaviour, and indications that wind turbine density may affect the magnitude of avoidance (Leopold et al. 2011; Krijgsveld et al. 2011). The estimated macro-avoidance rate of auks from studies at OWEZ was around 68%, although it should be noted that this was based on birds flying towards the windfarm and this value may not be appropriate for swimming birds. During daylight, wintering auks at OWEZ were observed mainly to sit on the water and float with tidal currents, with flight movements considered to be mainly corrections when birds drifted too far from favoured foraging areas (Krijgsveld et al. 2011). Razorbills and guillemots were seen inside OWEZ, although razorbills were never observed inside a neighbouring windfarm (PAWP) with a higher wind turbine density (Leopold et al. 2011).
- 283. The above studies were conducted at sites with relatively closely spaced wind turbines (e.g. 550m), while the minimum spacing at East Anglia TWO will be 1980m in row and 1210m between rows (see *Table 12.2*). Thus, a figure of 70% displacement represents a precautionary estimate.
- 284. Most of the windfarms in *Table 12.37* are beyond foraging range of breeding colonies for razorbills, so the majority of birds at risk of displacement would be non-breeding birds. The pressures on nonbreeding birds in terms of energy requirements are lower than during the breeding season, as they only need to obtain sufficient food to maintain their own survival (and not also to feed nestlings). In addition, auks can remain on the sea for extended periods and thus flight costs are minimised. Recoveries of ringed razorbills from UK colonies



have indicated a wide distribution in winter, with birds tending to move southwards and spreading throughout the southern and eastern North Sea, Celtic Sea, Channel, and Bay of Biscay (Furness 2015). It is likely that razorbills are relatively flexible in terms of where they spend the winter and are not dependent on particular foraging locations. There may be variation between years, and long-term changes in migration patterns, relating to prey availability and abundance, as is described for guillemot. Hence, the consequence of winter displacement from windfarms in terms of increased mortality is likely to be minimal. Given that, even when fish stocks have collapsed, survival rates of adult seabirds have shown declines of no more than 6 to 7% (e.g. kittiwake, Frederiksen et al. 2004) an increase in mortality due to displacement from windfarm sites seems likely to be at the low end of the proposed 1 to 10% range, and a value of 1% when combined with the precautionary 70% displacement rate is considered appropriate. On this basis, a precautionary cumulative nonbreeding displacement mortality of 692 is obtained (Table 12.38).

285. The largest BDMPS for razorbill in UK North Sea waters is 591,874 (Furness 2015). At the average baseline mortality rate of 0.174 (*Table 12.16*) the number of individuals expected to die in a year is 102,986 (591,874 x 0.174). The addition of a maximum of 692 individuals to this increases the background mortality rate by 0.7%. This magnitude of increase would not materially alter the background mortality of the population and would be undetectable. Therefore, the magnitude of effect is assessed as negligible. As the species is of low to medium sensitivity to disturbance, the potential for the proposed East Anglia TWO project to contribute to a significant cumulative displacement effect on razorbill is considered to be very small and the impact significance of cumulative displacement is **negligible**.



Offshore Windfarm(s)	No. razorbill	s at risk of displa	cement*		Source of information
	Breeding	Autumn migration	Winter	Spring migration	
Aberdeen	161	64	7	26	EATL (2016), Norfolk Vanguard Ltd (2018)
Beatrice	873	833	555	833	EATL (2016), Norfolk Vanguard Ltd (2018)
Blyth Demonstration	121	91	61	91	EATL (2016), Norfolk Vanguard Ltd (2018)
Dogger Bank Creyke Beck A	1250	1,576	1,728	4,149	EATL (2016), Norfolk Vanguard Ltd (2018)
Dogger Bank Creyke Beck B	1538	2,097	2,143	5,119	EATL (2016), Norfolk Vanguard Ltd (2018)
Dogger Bank Teesside A	834	310	959	1,919	EATL (2016), Norfolk Vanguard Ltd (2018)
Dogger Bank Teesside B	1153	592	1,426	2,953	EATL (2016), Norfolk Vanguard Ltd (2018)
Dudgeon	256	346	745	346	EATL (2016), Norfolk Vanguard Ltd (2018)
East Anglia ONE	16	26	155	336	EATL (2016), Norfolk Vanguard Ltd (2018)
East Anglia THREE	1807	1,122	1,499	1,524	East Anglia Three Ltd (2015)
Galloper	44	43	106	394	EATL (2016), Norfolk Vanguard Ltd (2018)
Greater Gabbard	0	0	387	84	EATL (2016), Norfolk Vanguard Ltd (2018)
Hornsea Project One	1109	4,812	1,518	1,803	EATL (2016), Norfolk Vanguard Ltd (2018)
Hornsea Project Two	2511	4,221	720	1,668	EATL (2016), Norfolk Vanguard Ltd (2018)
Hornsea Project Three	630	2,020	3,649	1,236	Orsted (2018)

#### Table 12.37 Cumulative Numbers of Razorbills at Risk of Displacement from Offshore Windfarms in the North Sea



Offshore Windfarm(s)	No. razorbill	s at risk of displa	cement*	Source of information		
	Breeding	Autumn migration	Winter	Spring migration		
Humber Gateway	27	20	13	20	EATL (2016), Norfolk Vanguard Ltd (2018)	
Inch Cape	1436	2,870	651		EATL (2016), Norfolk Vanguard Ltd (2018)	
Lincs and LID6	45	34	22	34	EATL (2016), Norfolk Vanguard Ltd (2018)	
London Array I and II	14	20	14	20	EATL (2016), Norfolk Vanguard Ltd (2018)	
Moray	2423	1,103	30	168	EATL (2016), Norfolk Vanguard Ltd (2018)	
Neart na Gaoithe	331	5,492	508		EATL (2016), Norfolk Vanguard Ltd (2018)	
Norfolk Vanguard East	599	491	279	752	Norfolk Vanguard Ltd (2018)	
Norfolk Vanguard west	280	375	348	172	Norfolk Vanguard Ltd (2018)	
Race Bank	28	42	28	42	EATL (2016), Norfolk Vanguard Ltd (2018)	
Seagreen A	3208				EATL (2016), Norfolk Vanguard Ltd (2018)	
Seagreen B	886				EATL (2016), Norfolk Vanguard Ltd (2018)	
Sheringham Shoal	106	1,343	211	30	EATL (2016), Norfolk Vanguard Ltd (2018)	
Teesside	16	61	2	20	EATL (2016), Norfolk Vanguard Ltd (2018)	
Thanet	3	0	14	21	EATL (2016), Norfolk Vanguard Ltd (2018)	
Thanet Extension	0	6	56	124	Vattenfall Wind Power Ltd (2018) Annex 4-3.	
Triton Knoll	40	254	855	117	EATL (2016), Norfolk Vanguard Ltd (2018)	



Offshore Windfarm(s)	No. razorbi	lls at risk of dis	placement*		Source of information
	Breeding	Autumn migration	Winter	Spring migration	
Westermost Rough	91	121	152	91	EATL (2016), Norfolk Vanguard Ltd (2018)
East Anglia TWO	288	55	148	263	Table 12.15
East Anglia North ONE	403	85	54	207	East Anglia ONE North Ltd (2019)
Norfolk Boreas	750	221	885	414	Norfolk Boreas Ltd. (2018)
Cumulative total	23,277	30,746	19,928	24,967	
Cumulative Annual total				98,927	
*Totals for the windfarm and a	a 2km buffer; blar	nks indicate no da	ata available for	a particular devel	lopment and season



### Table 12.38 Cumulative Displacement Matrix for Razorbill

Annual		Mortality rate										
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	99	198	297	396	495	989	1979	2968	4946	7914	9893
	20%	198	396	594	791	989	1979	3957	5936	9893	15828	19785
	30%	297	594	890	1187	1484	2968	5936	8903	14839	23743	29678
	40%	396	791	1187	1583	1979	3957	7914	11871	19785	31657	39571
	50%	495	989	1484	1979	2473	4946	9893	14839	24732	39571	49464
	60%	594	1187	1781	2374	2968	5936	11871	17807	29678	47485	59356
	70%	692	1385	2077	2770	3462	6925	13850	20775	34625	55399	69249
	80%	791	1583	2374	3166	3957	7914	15828	23743	39571	63313	79142
	90%	890	1781	2671	3561	4452	8903	17807	26710	44517	71228	89034
	100%	989	1979	2968	3957	4946	9893	19785	29678	49464	79142	98927



### 12.7.3.4 Guillemot

- 286. The East Anglia TWO windfarm site is located beyond the mean maximum foraging range of guillemot breeding colonies. Outside the breeding season, guillemots disperse from their breeding sites. Large numbers are found throughout the North Sea in the nonbreeding season (defined as August to February, Furness 2015).
- 287. The annual total of guillemots at risk of displacement from the proposed East Anglia TWO project is estimated as 4,146 individuals (summing the seasonal peak means on the East Anglia TWO windfarm site (and 2km buffer) for the breeding and non-breeding periods (*Table 12.15)*.
- 288. The recent cumulative assessments for offshore ornithology for East Anglia THREE (EATL 2016) and the Norfolk Vanguard project (Norfolk Vanguard Limited 2018) include estimates of the total numbers of guillemots at risk of displacement from other offshore windfarms in the North Sea. These estimates are included in *Table 12.39*, along with updates for projects where new information has become available since the publication of Norfolk Vanguard Limited (2018). These totals omit windfarms for which no data are available (Gunfleet Sands, Kentish Flats, Lynn and Inner Dowsing, Scroby Sands), but they are also likely to over-estimate the numbers present due to the precautionary use of seasonal peak numbers at each site rather than average numbers, which is likely to lead to double counting as birds move through the North Sea.
- 289. The estimated annual cumulative total of guillemots at risk of displacement from windfarms in the North Sea is 240,939 individuals (*Table 12.39*). Considering a range of displacement of 30 to 70%, and mortality of displaced individuals from 1 to 10%, based on advice from Natural England, the estimated number of guillemots subject to mortality from displacement throughout the year is between 723 and 16,866 (*Table 12.40*).
- 290. This is a large range so the assessment considers the most realistic value within this range. As discussed above under razorbill (*paragraphs 282* to *284*), a precautionary estimate of 70% displacement and of 1% mortality, 1697 birds (Table 12.40) is considered most appropriate.
- 291. The largest BDMPS for guillemot in UK North Sea waters is 1,617,306 (Furness 2015). At the average baseline mortality rate of 0.14 (*Table 12.16*) the number of individuals expected to die in a year is 226,423 (1,617,306 x 0.14). The addition of a maximum of 1,687 individuals to this increases the background mortality rate by 0.7%. This magnitude of increase would not materially alter the background mortality of the population and would be undetectable. Therefore, the magnitude of effect is assessed as negligible. As the species is



of low to medium sensitivity to disturbance, the potential for the proposed East Anglia TWO project to contribute to a significant cumulative displacement effect on guillemot is considered to be very small and the impact significance of cumulative displacement is **negligible**.



Offshore Windfarm(s)	No. guillemots at risk of disp	lacement*	Source of information
	Breeding	Non-breeding	
Aberdeen	547	225	EATL (2016), Norfolk Vanguard Ltd (2018)
Beatrice	13,610	2,755	EATL (2016), Norfolk Vanguard Ltd (2018)
Blyth Demonstration	1,220	1,321	EATL (2016), Norfolk Vanguard Ltd (2018)
Dogger Bank Creyke Beck A	5407	6,142	EATL (2016), Norfolk Vanguard Ltd (2018)
Dogger Bank Creyke Beck B	9479	10,621	EATL (2016), Norfolk Vanguard Ltd (2018)
Dogger Bank Teesside A	3283	2,268	EATL (2016), Norfolk Vanguard Ltd (2018)
Dogger Bank Teesside B	5211	3,701	EATL (2016), Norfolk Vanguard Ltd (2018)
Dudgeon	334	542	EATL (2016), Norfolk Vanguard Ltd (2018)
East Anglia ONE	274	640	EATL (2016), Norfolk Vanguard Ltd (2018)
East Anglia THREE	1669	2,859	EATL (2015)
Galloper	305	593	EATL (2016), Norfolk Vanguard Ltd (2018)
Greater Gabbard	345	548	EATL (2016), Norfolk Vanguard Ltd (2018)
Hornsea Project One	9836	8,097	EATL (2016), Norfolk Vanguard Ltd (2018)
Hornsea Project Two	7735	13,164	EATL (2016), Norfolk Vanguard Ltd (2018)
Hornsea Project Three	13,374	17,772	Orsted (2018)
Humber Gateway	99	138	EATL (2016), Norfolk Vanguard Ltd (2018)

#### Table 12.39 Cumulative Numbers of Guillemots at Risk of Displacement from Offshore Windfarms in the North Sea



Offshore Windfarm(s)	No. guillemots at risk of displ	acement*	Source of information
	Breeding	Non-breeding	
Inch Cape	4371	3,177	EATL (2016), Norfolk Vanguard Ltd (2018)
Lincs and LID6	582	814	EATL (2016), Norfolk Vanguard Ltd (2018)
London Array I and II	192	377	EATL (2016), Norfolk Vanguard Ltd (2018)
Moray	9820	547	EATL (2016), Norfolk Vanguard Ltd (2018)
Neart na Gaoithe	1755	3,761	EATL (2016), Norfolk Vanguard Ltd (2018)
Norfolk Vanguard East	2931	2,197	Norfolk Vanguard Ltd (2018)
Norfolk Vanguard west	1389	2,579	Norfolk Vanguard Ltd (2018)
Race Bank	361	708	EATL (2016), Norfolk Vanguard Ltd (2018)
Seagreen A	16500		EATL (2016), Norfolk Vanguard Ltd (2018)
Seagreen B	16054		EATL (2016), Norfolk Vanguard Ltd (2018)
Sheringham Shoal	390	715	EATL (2016), Norfolk Vanguard Ltd (2018)
Teesside	267	901	EATL (2016), Norfolk Vanguard Ltd (2018)
Thanet	18	124	EATL (2016), Norfolk Vanguard Ltd (2018)
Thanet Extension	12	1,178	Vattenfall Wind Power Ltd (2018) Annex 4-3.
Triton Knoll	425	746	EATL (2016), Norfolk Vanguard Ltd (2018)
Westermost Rough	347	486	EATL (2016), Norfolk Vanguard Ltd (2018)



Offshore Windfarm(s)	No. guillemots at risk of disp	lacement*	Source of information		
	Breeding	Non-breeding			
East Anglia TWO	2,126	2,020	Table 12.15		
East Anglia North ONE	4,183	1,888	East Anglia ONE North Ltd (2019)		
Norfolk Boreas	2,413	10,471	Norfolk Boreas Ltd (2018)		
Cumulative total	136,864	104,075			
Cumulative Annual total	240,939				
*Totals for the windfarm and a 21	km buffer; blanks indicate no data	available for a particular develop	oment and season		



#### Table 12.40 Cumulative Displacement Matrix for Guillemot

Annual		Mortality	rate									
		1%	2%	3%	4%	5%	10%	20%	30%	50%	80%	100%
Displacement	10%	241	482	723	964	1205	2409	4819	7228	12047	19275	24094
	20%	482	964	1446	1928	2409	4819	9638	14456	24094	38550	48188
	30%	723	1446	2168	2891	3614	7228	14456	21685	36141	57825	72282
	40%	964	1928	2891	3855	4819	9638	19275	28913	48188	77101	96376
	50%	1205	2409	3614	4819	6023	12047	24094	36141	60235	96376	120470
	60%	1446	2891	4337	5783	7228	14456	28913	43369	72282	115651	144564
	70%	1687	3373	5060	6746	8433	16866	33732	50597	84329	134926	168658
	80%	1928	3855	5783	7710	9638	19275	38550	57825	96376	154201	192752
	90%	2168	4337	6505	8674	10842	21685	43369	65054	108423	173476	216845
	100%	2409	4819	7228	9638	12047	24094	48188	72282	120470	192752	240939



### 12.7.4 Cumulative Assessment of Operational Collision Risk

- 292. The species assessed for project alone collision impacts (and the relevant seasons) were those for which a collision mortality greater than one individual for the project alone was estimated, on the grounds that the potential for the proposed East Anglia TWO project to contribute to a cumulative mortality effect was negligible for annual mortalities below this. Thus, cumulative collision risk both annually and for key seasons was assessed for gannet, kittiwake, lesser black-backed gull and great black-backed gull.
- 293. It is considered that all of the windfarms identified for inclusion in the CIA in *Table 12.35* have the potential to contribute to a cumulative effect.

#### 12.7.4.1 Gannet

- 294. The cumulative gannet collision risk prediction is set out in *Table 12.41*. This collates collision predictions from other windfarms which may contribute to the cumulative total. This table takes the recently submitted windfarm assessment for Norfolk Vanguard as its starting point.
- 295. The cumulative totals of collision mortality in each season, and summed across seasons, are presented in *Table 12.41*. Assessments at other windfarms have been conducted using a range of avoidance rates and alternative collision model Options. In order to simplify interpretation of the data across sites and also to bring these assessments up to date with the current Natural England Advice, the values in *Table 12.41* are those estimated using the Band Model Option 1 (or 2, if that was the one presented) standardised at an avoidance rate of 98.9%.
- 296. All windfarm estimates have also been updated to reflect the evidence based nocturnal flight activity rates reported in Furness et al. (2018b). Furness et al. (2018b) recommended precautionary nocturnal activity rates for gannet in the breeding and nonbreeding seasons of 8% and 4% respectively. However, the actual average rates from their study were 7.1% and 2.3% respectively. Furthermore, the breeding season value was very heavily influenced by the results from the smallest study in the review, which was based on only three tagged birds in Shetland (Garthe et al., 1999). This study yielded a nocturnal activity rate of 20.9% (compared to daytime) but the total duration of flight activity recorded was only 215 hours, which was less than 3% of the > 8,000 hours covered by the remaining studies. If the average rate is calculated without this study a breeding season rate of 4.3% (SE 2.7%) is obtained. This is considered to be more robust and has been used in the current assessment. Similarly, the actual nonbreeding season rate of 2.3% (SE 0.4%) has been used here in preference to the rounded-up value of 4% reported in Furness et al. (2018). Use of these evidence-based rates for gannet improves the scientific basis of the assessment while also reducing unnecessary precaution.


#### Table 12.41 Cumulative Collision Risk Assessment for Gannet

Tier	Windfarm	Breeding	Autumn	Spring	Annual
		CRM	CRM	CRM	CRM
1	Beatrice Demonstrator	0.6	0.9	0.7	2.2
1	Greater Gabbard	0	0	0	0
1	Gunfleet Sands	0	0	0	0
1	Kentish Flats	0.2	0	0	0.2
1	Lincs	4.3	0	0	4.3
1	London Array	0	0	0	0
1	Lynn and Inner Dowsing	0	0	0	0
1	Scroby Sands	0	0	0	0
1	Sheringham Shoal	10.3	4.7	0	15
1	Teeside	5.6	0	0	5.6
1	Thanet	0.8	0	0	0.8
1	Humber Gateway	2.9	0	0	2.9
1	Westernmost Rough	0.2	0	0	0.2
2	Beatrice	20.0	48.2	10	78.2
2	Dudgeon	20.1	30.3	15.4	65.8
2	Galloper	9.4	24.9	13.5	47.8
2	Race Bank	29.6	6.9	3.1	39.6
2	Rampion	36.4	50.1	1.1	87.6
2	Hornsea Project One	10.3	23.6	17.5	51.4
3	Blythe Demonstration	0	0	0	0
3	Dogger Bank Creyke Beck Projects A and B	10.0	10.3	6.7	27
3	East Anglia ONE	1.8	66.0	3.6	71.4
3	Aberdeen (EOWDC)	2.9	2.2	0.0	5.1
3	Forth (Seagreen) Alpha and Bravo	742.5	111.1	83.8	937.4
3	Inch Cape	337.5	30.3	7.2	375

### East Anglia TWO Offshore Windfarm Preliminary Environmental Information Report



Tier	Windfarm	Breeding	Autumn	Spring	Annual
		CRM	CRM	CRM	CRM
3	Moray Firth (EDA)	48.3	50.4	30.2	128.9
3	Neart na Gaoithe	130.6	38.6	18.5	187.7
3	Dogger Bank Teeside A and B (now Sofia)	13.6	8.1	8.7	30.4
3	Triton Knoll	24.4	64.6	40.8	129.8
3	Hornsea Project Two	7.9	10.2	4.0	22.1
4	East Anglia THREE	5.2	24.7	7.1	37
4	Hornsea Project three	18.3	12.1	8.1	38.5
4	Thanet Extension	0	2.9	7.1	10.0
4	Norfolk Vanguard	18.3	62.3	29.9	110.5
5	East Anglia TWO	8.8	8.6	1.2	18.6
5	East Anglia ONE North	8.8	5.5	1.3	15.6
5	Norfolk Boreas	15	42.1	11.1	68.2
	TOTALS	1544.6	739.6	330.6	2614.8

297. The annual cumulative total for estimated collision mortality is 2,615. Note, however that many of the collision estimates for other windfarms were calculated on the basis of designs with higher total rotor swept areas than have been installed (or are planned), which is a key factor in collision risk. For example, the Beatrice windfarm, which is currently under construction, was consented on the basis of up to 125 x 7MW wind turbines but only 84 (of the same model) will be installed, leading to a reduction in mortality risk of 33%. A method for updating collision estimates for changes in windfarm design such as this was presented in MacArthur Green (2017). This uses ratios of consented and as-built turbine parameters to adjust the collision risk mortality estimates for a consented wind farm. Updating the collision estimates for the Beatrice windfarm using this approach reduces the predicted annual mortality from 96 to 64 (not accounting for revised nocturnal activity rates. Similar reductions in collision mortality are likely to result from changes in windfarm design at other consented but not constructed windfarms in the North Sea which are included in the cumulative and in-combination totals. This could include Firth of Forth Alpha and Bravo, Hornsea Project One and Two, Inch Cape, Moray Firth, Neart na Gaoithe, Race Bank, Rampion, Dogger Bank Creyke Beck A and B and Teesside A and B and Triton Knoll. Applying similar reductions to these



windfarms in **Table 12.41** would reduce the cumulative annual mortality by about 400. Therefore, the values presented in **Table 12.41**, as well as being based on precautionary calculation methods, can be seen to overestimate the total risk by around 15% due to the reduced collision risks for projects which undergo design revisions post-consent.

- Previous gannet collision assessments for the windfarms listed in Table 12.41 298. have been made on the basis of Band model Option 1 and a range of avoidance rates between 95% and 99%. The current rate of 98.9% dates from November 2014 (JNCC et al., 2014) and followed the review conducted by Cook et al. (2014). Therefore, the decisions for some of the projects consented prior to this date were on the basis of estimated cumulative collision mortality numbers which were higher than the values presented in Table 12.41. However, given the variation in rates presented in different assessments and the rates used in reaching consent decisions, it is difficult to confidently determine the avoidance rate used for each windfarm consent decision. Nonetheless, it can be stated with a good degree of certainty that none of the previous windfarms have been consented on the basis of an avoidance rate higher than 99%, and many will have been based on assessment at 98%. It therefore follows that the cumulative total including the proposed East Anglia TWO project (2,615) is almost certainly lower than those on which some recent consent decisions have been granted.
- 299. Work conducted at the Greater Gabbard windfarm (APEM, 2014) has also found that gannet avoidance of windfarms during the autumn migration period may be even higher than the current estimate of 98.9%. Of 336 gannets observed during this study, only 8 were recorded within the windfarm, indicating a high degree of windfarm (macro) avoidance. Analysis of their data indicated a macro-avoidance rate in excess of 95% compared with the current guidance value of 64% (see *paragraph 168* above). When combined with meso- and micro-avoidance this would result in higher overall avoidance than the current 98.9% and would further reduce the total collision mortality prediction.
- 300. A bird flight behaviour study commissioned by ORJIP provides further evidence relating to the precautionary nature of current avoidance rates and other parameters used in windfarm assessment (Skov et al. 2018). Based on a combination of video, radar and field observations at Thanet Offshore Windfarm, the empirical avoidance rate for gannet was calculated as 99.9%.
- 301. Demographic data were collated for the British gannet population to produce a population model which was used to consider the potential impact of additional mortality (WWT, 2012). Two versions of the model were developed, with and without density dependence. Of these two models, the density independent one was considered to provide more reliable predictions since it predicted baseline



growth at a rate close to that recently observed (1.28% per year compared with an observed rate of 1.33%) while the density dependent model predicted baseline growth of 0.9%. While density-dependent regulation of populations is to be expected as the norm, in the case of gannet the population has been increasing for many decades, suggesting that the population has not yet reached a level where density-dependent regulation is a major influence on its dynamics.

- 302. The study concluded that, using the density independent model, population growth, on average, would remain positive until additional mortality exceeded 10,000 individuals per year while the lower 95% confidence interval on population growth remained positive until additional mortality exceeded 3,500 individuals, which is greater than the cumulative total in *Table 12.41*. Consideration was also given to the risk of population decline. The risk of a 5% population decline was less than 5% for additional annual mortalities below 5,000 (using either the density dependent or density independent model; WWT, 2012).
- 303. It is important to note that the gannet model presented in WWT (2012) was based on the entire British population, so collisions at windfarms on the west coast also need to be considered for consistency. However, a review of applications in the Irish Sea and Solway Firth (Barrow, Burbo Bank, Burbo Bank Extension, Gwynt Y Mor, North Hoyle, Ormonde, Rhyl Flats, Robin Rigg, Walney 1 and 2, Walney Extension and West of Duddon Sands) gave a gannet annual collision cumulative total of 243 at an avoidance rate of 98.9%. Therefore, inclusion of these windfarms in the assessment does not alter the conclusion that cumulative collisions are below a level at which a significant impact on the British gannet population would result.
- 304. Furthermore, the WWT (2012) analysis was conducted using the estimated gannet population in 2004 (the most recent census available at that time), when the British population was estimated to be 261,000 breeding pairs. The most recent census indicates the equivalent number of breeding pairs is now a third higher at 349,498 (Murray et al., 2015). This increase in size will raise the thresholds at which impacts would be predicted and therefore further reduces the risk of significant impacts.
- 305. In conclusion, the cumulative impact on the gannet population due to collisions both year round and within individual seasons is considered to be of low magnitude, and the relative contribution of the proposed East Anglia TWO project to this cumulative total is small. Gannets are considered to be of low to medium sensitivity to collision mortality and the impact significance is therefore **minor adverse**.



### 12.7.4.1 Kittiwake

- 306. The cumulative collision risk predictions for kittiwake are set out in **Table 12.42**. This collates collision predictions from other windfarms which may contribute to the cumulative total. This table takes the recently submitted windfarm assessment for Norfolk Vanguard as its starting point.
- 307. The cumulative totals of collision mortality in each season, and summed across seasons, are presented in *Table 12.42*. Assessments at other windfarms have been conducted using a range of avoidance rates and alternative collision model Options. In order to simplify interpretation of the data across sites and also to bring these assessments up to date with the current Natural England Advice, the values in *Table 12.42* are those estimated using the Band Model Option 1 (or 2, if that was the one presented) standardised at an avoidance rate of 98.9%.
- 308. All windfarm collision estimates have been updated to reflect the evidence based nocturnal flight activity rates estimated by Furness et al. (in prep). This is a review of nocturnal activity in kittiwakes derived from analysis of extensive geolocator tag data from several studies and a limited amount of GPS tracking data. The previous estimated value for nocturnal flight activity (50%) was found to be a considerable overestimate. Evidence-based rates of 20% during the breeding season and 17% during the nonbreeding season were found. It is straightforward to adjust mortality estimates for other windfarms considered in the cumulative assessment using the new and old nocturnal activity rates and the monthly number of daytime and night-time hours (i.e. it is not necessary to rerun the collision model for this update). Use of these evidence-based rates for kittiwake improves the scientific basis of the assessment while also reducing unnecessary precaution.

lier	windfarm	Breeding	Autumn	Spring	Annual
		CRM	CRM	CRM	CRM
1	Beatrice Demonstrator	0	2.1	1.7	3.8
1	Greater Gabbard	1.1	15	11.4	27.5
1	Gunfleet Sands	0	0	0	0
1	Kentish Flats	2.4	0	0	2.4
1	Lincs	0	0	0	0
1	London Array	0	0	0	0
1	Lynn and Inner	0	0	0	0

#### Table 12.42 Cumulative Collision Risk Assessment for Kittiwake

# East Anglia TWO Offshore Windfarm Preliminary Environmental Information Report



Tier	Windfarm	Breeding	Autumn	Spring	Annual
		CRM	CRM	CRM	CRM
	Dowsing				
1	Scroby Sands	0	0	0	0
1	Sheringham Shoal	0	0	0	0
1	Teeside	14.0	0	0	14
1	Thanet	0.8	0	0	0.8
1	Humber Gateway	5.7	0	0	5.7
1	Westernmost Rough	0.4	0	0	0.4
2	Beatrice	62.1	11.2	48.4	121.7
2	Dudgeon	0	0	0	0
2	Galloper	8.6	37.2	12.9	58.7
2	Race Bank	1.3	17.1	4.4	22.8
2	Rampion	62.6	31.3	21.9	115.8
2	Hornsea Project One	39.3	41.8	18.9	120.8
3	Blythe Demonstration	0	0	0	0
3	Dogger Bank Creyke Beck Projects A and B	197	100.1	277.6	574.7
3	East Anglia ONE	0.9	73.7	23.4	98
3	Aberdeen (EOWDC)	14.7	5.7	1.0	21.1
3	Forth (Seagreen) Alpha and Bravo	356.7	229.8	255.6	842.1
3	Inch Cape	10.3	193.1	25	228.4
3	Moray Firth (EDA)	36.6	8.1	25.1	69.8
3	Neart na Gaoithe	9.1	17.9	2.2	29.2
3	Dogger Bank Teeside A and B (now Sofia)	86.7	65.6	197.6	349.9
3	Triton Knoll	12.4	93.3	53.4	159.1
3	Hornsea Project Two	11.0	7.8	3.7	22.5
4	East Anglia THREE	5.2	46.5	27.1	78.8



Tier	Windfarm	Breeding	Autumn	Spring	Annual
		CRM	CRM	CRM	CRM
4	Hornsea Project three	86.0	70.3	71.0	227.3
4	Thanet Extension	1.8	2.1	7.3	11.2
4	Norfolk Vanguard	20.8	61.3	76.2	158.3
5	East Anglia TWO	13.6	2.9	9.3	25.8
5	East Anglia ONE North	6.0	4.3	17.4	27.7
5	Norfolk Boreas	26.6	83.4	43.7	155.8
	TOTALS	1093.7	1221.6	1236.2	3574.1

- 309. The estimated annual cumulative total is 3,418. Note, however that many of the collision estimates for other windfarms were calculated on the basis of designs with higher total rotor swept areas than have been installed (or are planned), which is a key factor in collision risk. For example, the Beatrice windfarm, which is currently under construction, was consented on the basis of up to 125 x 7MW wind turbines but only 84 (of the same model) will be installed, leading to a reduction in mortality risk of 33%. A method for updating collision estimates for changes in windfarm design was presented in Macarthur Green (2017). Updating the collision estimates for the Beatrice windfarm using this approach reduces the predicted annual mortality from 145 to 97 (not accounting for revised nocturnal activity rates). Similar reductions in collision mortality are likely to result from changes in windfarm design at other consented but not constructed windfarms in the North Sea which are included in the cumulative and in-combination totals. This could include Firth of Forth Alpha and Bravo, Hornsea Project One and Two, Inch Cape, Moray Firth, Neart na Gaoithe, Race Bank, Rampion, Dogger Bank, Creyke Beck A and B, Teesside A and B and Triton Knoll. Applying the same method to these windfarms could achieve a reduction in the cumulative annual mortality of around 550. Therefore, the values presented in Table 12.42, as well as being based on precautionary calculation methods, can be seen to overestimate the total risk by around 15% due to the reduced collision risks for projects which undergo design revisions post consent.
- 310. For the assessment of the nearby East Anglia THREE windfarm, a kittiwake population model was developed to assess the potential effects of cumulative predicted mortality from collisions with offshore windfarms on the kittiwake BDMPS populations (EATL 2015). Both density independent and density dependent models were developed. For annual mortality of 4,000, the density



dependent model predicted the population after 25 years would be 3.6% to 4.4% smaller than that predicted in the absence of additional mortality from collisions with offshore wind farms, while the more precautionary density independent model predicted equivalent declines of 10.3% to 10.9%. There is evidence that kittiwake populations are limited by food supply, and therefore are subject to density-dependent regulation (Frederiksen et al. 2004, 2007; Cury et al. 2011; Sandvik et al. 2012; Carroll et al. 2017), and therefore the densitydependent model is more appropriate for this species. To place these predicted magnitudes of change in context, over three approximately 15 year periods (between censuses) the British kittiwake population changed by +24% (1969 to 1985). -25% (1985 to 1998) and -44% (2000)to 2015) (http://jncc.defra.gov.uk/page-3201 accessed 15<sup>th</sup> October 2018). Changes of between 3% and 10% across a longer (25 year) period against a background of changes an order of magnitude larger will almost certainly be undetectable. It is possible that the longer term decline will continue and the population is unlikely to recover over this period. However even precautionary estimates of additional mortality from offshore windfarms are not predicted to significantly increase the rate of decline or to prevent the population from recovering should environmental conditions become more favourable.

- 311. Evidence for density dependent regulation of the North Sea kittiwake population was summarised in EATL (2015). Trinder (2014) explored a range of strengths of density dependence for this species and identified model parameters which produced population predictions consistent with patterns of seabird population growth which have been observed across a wide range of taxa (including kittiwake) worldwide (Cury et al. 2011). Thus, there is robust evidence for density dependent regulation of the North Sea kittiwake population (and for seabirds more widely) and its inclusion in the kittiwake population model (EATL 2015) balanced this evidence with reasonable precaution. Consequently, the density dependent kittiwake model results are considered to be the more robust ones on which to base this assessment.
- 312. Kittiwake is considered to be of low to medium sensitivity and the magnitude of effect described above is considered to be low. Consequently, the worst case cumulative collision mortality is considered to be of low magnitude, resulting in impacts of **minor adverse** significance. However, when the various sources of precaution are taken in to account (precautionary avoidance rate estimates, reduction in construction versus consented windfarm sizes, over-estimated nocturnal activity) the cumulative collision risk impact magnitude is almost certainly smaller still.



### 12.7.4.2 Lesser Black-backed Gull

- 313. The cumulative collision risk prediction for lesser black-backed gull is set out in *Table 12.43*. This collates collision predictions from other windfarms which may contribute to the cumulative total. This table takes the recently submitted windfarm assessment for Norfolk Vanguard as its starting point.
- 314. The collision values presented in *Table 12.43* include totals for breeding, nonbreeding and annual periods. However, not all projects provide a seasonal breakdown of collision impacts, therefore it is not possible to extract data from these periods for cumulative assessment. Natural England has previously noted that an 80:20 split between the nonbreeding and breeding seasons is appropriate for lesser black-backed gull in terms of collision estimates (Natural England 2013). Therefore, for those sites where a seasonal split was not presented the annual numbers in *Table 12.43* have been multiplied by 0.8 to estimate the nonbreeding component and 0.2 to estimate the breeding component.
- 315. Assessments for other windfarms have been conducted using a range of avoidance rates and alternative collision model Options. In order to simplify interpretation of the data across sites and also to bring these assessments up to date with the current Natural England advice, the values in *Table 12.43* are those estimated using the Band Model Option 1 (or 2, if that was the one presented) at an avoidance rate of 99.5%. (Note that estimates for the Dogger Bank projects have only been presented using Band model Option 3. Therefore, these values in *Table 12.43* have been converted to the Natural England advised rate for this model of 98.9%).

Tier	Windfarm	Breeding	Non-breeding	Annual
		CRM	CRM	CRM
1	Beatrice Demonstrator	0	0	0
1	Greater Gabbard	12.4	49.6	62
1	Gunfleet Sands	0	0	0
1	Kentish Flats	0.3	1.3	1.6
1	Lincs	1.7	6.8	8.5
1	London Array	0	0	0
1	Lynn and Inner Dowsing	0	0	0

Table 12.43 Cumulative Collision Risk Assessment for Lesser Black-backed Gull

# East Anglia TWO Offshore Windfarm Preliminary Environmental Information Report



Tier	Windfarm	Breeding	Non-breeding	Annual
		CRM	CRM	CRM
1	Scroby Sands	0	0	0
1	Sheringham Shoal	1.7	6.6	8.3
1	Teeside	0	0	0
1	Thanet	3.2	12.8	16
1	Humber Gateway	0.2	1.1	1.3
1	Westernmost Rough	0.0	0.3	0.3
2	Beatrice	0	0	0
2	Dudgeon	7.7	30.6	38.3
2	Galloper	27.8	111	138.8
2	Race Bank	43.2	10.8	54
2	Rampion	1.6	6.3	7.9
2	Hornsea Project One	4.4	17.4	21.8
3	Blythe Demonstration	0	0	0
3	Dogger Bank Creyke Beck Projects A and B	2.6	10.4	13
3	East Anglia ONE	4	23	27
3	Aberdeen (EOWDC)	0	0	0
3	Forth (Seagreen) Alpha and Bravo	2.1	8.4	10.5
3	Inch Cape	0	0	0
3	Moray Firth (EDA)	0	0	0
3	Neart na Gaoithe	0.3	1.2	1.5
3	Dogger Bank Teeside A and B (now Sofia)	2.4	9.6	12
3	Triton Knoll	7.4	29.6	37
3	Hornsea Project Two	2	2	4
4	East Anglia THREE	1.8	8.2	10
4	Hornsea Project three	15	3	18



Tier	Windfarm	Breeding	Non-breeding	Annual
		CRM	CRM	CRM
4	Thanet Extension	1.5	0.8	2.3
4	Norfolk Vanguard	23.3	4.1	27.4
5	East Anglia TWO	0.5	0	0.5
5	East Anglia ONE North	0.6	0	0.6
5	Norfolk Boreas	22.0	5.2	27.2
	TOTALS	189.7	360.1	549.8

- The cumulative predicted annual total is 550. Note, however that many of the 316. collision estimates for other windfarms were calculated on the basis of designs with higher total rotor swept areas than have been installed (or are planned), which is a key factor in collision risk. For example, the Galloper windfarm, which is currently under construction, was consented on the basis of 140 wind turbines but only 56 have been installed. A method for updating collision estimates for changes in windfarm design was presented in MacArthur Green (2017). Updating the collision estimates for the Galloper windfarm using this approach reduces the predicted annual mortality from 139 to 60. Turbine numbers may also be reduced at a number of other windfarms, including Firth of Forth Alpha and Bravo, Hornsea Project One and Two, Inch Cape, Moray Firth, Neart na Gaoithe, Race Bank, Rampion, Dogger Bank Creyke Beck A and B and Teesside A and B and Triton Knoll. Applying the same method to the other windfarms in Table 12.43 can achieve a reduction in the cumulative annual mortality of around 200. Therefore, the values presented in Table 12.43, as well as being based on precautionary calculation methods, can be seen to overestimate the total risk by around 36% due to the reduced collision risks for projects which undergo design revisions post consent.
- 317. Lesser black-backed gull collision assessments undertaken prior to 2014 were made on the basis of Band Model Option 1 and an avoidance rate of 98%, with the change to 99.5% dating from November 2014 (JNCC et al., 2014). Therefore, projects consented prior to this date were on the basis of a cumulative collision mortality 4 times that presented in *Table 12.43*. Projects included in the cumulative total at this time were those to Triton Knoll in *Table 12.43* (with reference to information on the timing of consents in *Table 12.35*) with a cumulative total of 461 predicted collisions (adding annual totals for all sites up to triton Knoll in *Table 12.43*). This total includes four projects



consented after November 2014 (Hornsea Project 1, 22 annual collisions at 99.5%; Dogger Bank Creyke Beck A and B, 13 annual collisions at 98.9% Option 3; Dogger Bank Teesside A and B, 12 annual collisions at 98.9% Option 3). Thus, at this time, the previous cumulative collision total (at 98%) excluding these three projects would have been 1,656 (461 – (22+13+12) x 4). The current worst case cumulative total of 550, including all consented and still to be consented projects, is therefore much lower than this previously accepted cumulative total.

- A review of nocturnal activity in seabirds (EATL 2015) has indicated that the 318. value currently used for this parameter (50%) to estimate collision risk at night for lesser black-backed gull is almost certainly an overestimate, possibly by as much as a factor of two (i.e. study data suggest that 25% is more appropriate). Reducing the nocturnal activity factor to 25% reduced collision estimates by around 15%. Natural England have recognised this aspect of precaution and advised recent projects to undertake collision modelling with nocturnal activity set to both 25% and 50%. This was included in the proposed East Anglia TWO project collision modelling (by setting the nocturnal factor in simulated model runs to be randomly selected as one of these two values). However, this adjustment to nocturnal activity is also applicable to the other cumulative collision estimates. A similar correction applied to the other windfarms would further reduce the overall collision estimate for all windfarms by a significant amount (e.g. between 7% and 25%; note the magnitude of reduction varies depending on the time of year and windfarm latitude due to the variation in day and night length). This further emphasises the precautionary nature of the current assessment.
- 319. In conclusion, the current cumulative total is considerably lower than previously consented cumulative totals (as much as 3 times lower), and yet this total still includes several sources of precaution (e.g. consented vs. built impacts and overestimated nocturnal activity). Therefore, the cumulative impact on the lesser black-backed gull population due to collisions both year round and within individual seasons is considered to be of low magnitude and lesser black-backed gull is considered to be of medium sensitivity, therefore the impact significance is **minor adverse**.

### 12.7.4.3 Great Black-backed Gull

- 320. The cumulative predicted collision risk for great black-backed gull is set out in *Table 12.44*. This collates collision predictions from other windfarms which may contribute to the cumulative total. This table takes the recently submitted windfarm assessment for Norfolk Vanguard as its starting point.
- 321. The collision values presented in *Table 12.44* include breeding, nonbreeding and annual collision totals. However, not all projects provide a seasonal



breakdown of collision impacts, therefore it is not possible to extract data from these periods for cumulative assessment. Natural England has previously noted that an 80:20 split between the nonbreeding and breeding seasons is appropriate for lesser black-backed gull in terms of collision estimates (Natural England, 2013). This ratio is considered to also be appropriate for great black-backed gull, therefore for those sites where a seasonal split was not presented the annual numbers in **Table 12.44** have been multiplied by 0.8 to estimate the nonbreeding component and 0.2 to estimate the breeding component.

322. Assessments for other windfarms have been conducted using a range of avoidance rates and alternative collision model Options. In order to simplify interpretation of the data across sites and also to bring these assessments up to date with the current Natural England advice, the values in *Table 12.44* are those estimated using the Band Model Option 1 (or 2, if that was the one presented) at an avoidance rate of 99.5%. Note that estimates for the Dogger Bank projects have only been presented using Band Model Option 3. Therefore, these values in *Table 12.44* have been converted to the Natural England advised rate for this model of 98.9%).

Tier	windrarm	Бгееанд	Non-breeding	Annual
		CRM	CRM	CRM
1	Greater Gabbard	15	60	75
1	Gunfleet Sands	0	0	0
1	Kentish Flats	0.1	0.2	0.3
1	Lincs	0	0	0
1	London Array	0	0	0
1	Lynn and Inner Dowsing	0	0	0
1	Scroby Sands	0	0	0
1	Sheringham Shoal	0	0	0
1	Teeside	8.7	34.8	43.6
1	Thanet	0.1	0.4	0.5
1	Humber Gateway	1.3	5.1	6.3
1	Westernmost Rough	0	0	0.1
2	Beatrice	30.2	120.8	151

Table 12.44 Cumulative Collision Risk Assessment for Great Black-backed Gull

# East Anglia TWO Offshore Windfarm Preliminary Environmental Information Report



Tier	Windfarm	Breeding	Non-breeding	Annual
		CRM	CRM	CRM
2	Dudgeon	0	0	0
2	Galloper	4.5	18	22.5
2	Race Bank	0	0	0
2	Rampion	5.2	20.8	26
2	Hornsea Project One	17.2	68.6	85.8
3	Blythe Demonstration	1.3	5.1	6.3
3	Dogger Bank Creyke Beck Projects A and B	5.8	23.3	29.1
3	East Anglia ONE	0	32	32
3	Aberdeen (EOWDC)	0.6	2.4	3
3	Forth (Seagreen) Alpha and Bravo	13.4	53.4	66.8
3	Inch Cape	0	36.8	36.8
3	Moray Firth (EDA)	9.5	25.5	35
3	Neart na Gaoithe	0.9	3.6	4.5
3	Dogger Bank Teeside A and B (now Sofia)	6.4	25.5	31.9
3	Triton Knoll	24.4	97.6	122
3	Hornsea Project Two	3	20	23
4	East Anglia THREE	4.6	34.4	39
4	Hornsea Project three	16	50	66
4	Thanet Extension	1.3	20.8	22.2
4	Norfolk Vanguard	0	22.2	22.2
5	Norfolk Boreas	16.4	58.3	74.7
5	East Anglia TWO	3	0.6	3.6

Tier	Windfarm	Breeding	Non-breeding	Annual
		CRM	CRM	CRM
5	East Anglia ONE North	0	0.5	0.5
	TOTALS	188.9	840.7	1029.7

- 323. The annual cumulative total of predicted collisions is 1,030. Note, however that many of the collision estimates for other windfarms were calculated on the basis of designs with higher total rotor swept areas than have been installed (or are planned), which is a key factor in collision risk. For example, the Beatrice windfarm, which is currently under construction, was consented on the basis of 125 wind turbines but only 84 are being installed. A method for updating collision estimates for changes in windfarm design was presented in MacArthur Green (2017). Updating the collision estimates for the Beatrice windfarm using this approach reduces the predicted annual mortality from 151 to 101. Turbine numbers may also be reduced at a number of other windfarms, including Firth of Forth Alpha and Bravo, Hornsea Project One and Two, Inch Cape, Moray Firth, Neart na Gaoithe, Race Bank, Rampion, Dogger Bank Creyke Beck A and B and Teesside A and B and Triton Knoll. Applying the same method to other windfarms can achieve a reduction in the cumulative annual mortality (Table 12.44) of around 260. Therefore, the values presented in Table 12.44, as well as being based on precautionary calculations, can be seen to overestimate the total risk by around 25% due to the reduced collision risks for projects which undergo design revisions post consent.
- 324. A review of nocturnal activity in seabirds (EATL 2015) has indicated that the value currently used for this parameter (50%) to estimate collision risk at night for great black-backed gull is almost certainly an overestimate, possibly by as much as a factor of two (i.e. study data suggest that 25% is more appropriate). Reducing the nocturnal activity factor to 25% reduced collision estimates by around 15%. Natural England have recognised this aspect of precaution and advised recent projects to undertake collision modelling with nocturnal activity set to both 25% and 50%. This was included in the proposed East Anglia TWO project collision modelling (by setting the nocturnal factor in simulated model runs to be randomly selected as one of these two values). However, this adjustment to nocturnal activity is also applicable to the other cumulative collision estimates. A correction applied to the other windfarms similar to that used for Norfolk Vanguard along these lines would further reduce the overall collision estimate for all windfarms by a significant amount (e.g. between 7%) and 25%; note the magnitude of reduction varies depending on the time of year



and windfarm latitude due to the variation in day and night length). This further emphasises the precautionary nature of the current assessment.

- 325. Great black-backed gull collision assessments undertaken prior to 2014 were made on the basis of Band Model Option 1 and an avoidance rate of 98%, with the change to 99.5% dating from November 2014 (JNCC et al. 2014). This has resulted in a large reduction in predicted cumulative totals to the extent that the current cumulative estimate of 955 is much lower than cumulative totals for consented developments on which it has previously been concluded there will be no adverse effect on the population in the long term (DECC 2014).
- 326. A population model for great black-backed gull was developed to inform that East Anglia THREE assessment (EATL 2016a). Four versions of the model were presented, using two different sets of demographic rates (from the literature) and both with and without density dependent regulation of reproduction. Comparison of the historical population trend with the outputs from these models indicated that the density dependent versions generated population predictions which were much more closely comparable to the population trend. The density dependent models were also less sensitive to which set of demographic rates was used. The density dependent versions were therefore considered to provide a more reliable predictive tool.
- 327. Using the density dependent model, application of an additional annual mortality of 900 to the great black-backed gull BDMPS resulted in impacted populations after 25 years which were 6.1% to 7.7% smaller than in the absence of impact. The equivalent density independent predictions generated population reductions of 21.3% to 21.5%. Based on the modelling, Natural England concluded that whilst a significant cumulative effect could not be ruled out, the contribution of East Anglia THREE was so small that it would not materially affect the overall cumulative impact magnitude. The final East Anglia THREE annual collision impact for great black-backed gull was 39, compared with only 3.6 for the proposed East Anglia TWO project.
- 328. In conclusion, the cumulative impact on the great black-backed gull population due to predicted collisions both year round and within individual seasons is considered to be of low magnitude and great black-backed gull is considered to be of low to medium sensitivity, therefore the impact significance is **minor adverse**.



## **12.8Transboundary Impacts**

329. With regard to the potential for transboundary cumulative impacts, there is clearly potential for collisions and displacement at windfarms outside UK territorial waters. However, the operational offshore windfarms in Belgium, the Netherlands and Germany are comparatively small (in combination these projects are of a similar size to no more than one to two of the more recent UK windfarms, such as East Anglia ONE). Since the spatial scale and hence seabird populations sizes for a transboundary assessment would be much larger, it is apparent that the scale of windfarm development would be relatively much smaller. Therefore, the inclusion of non-UK windfarms is considered very unlikely to alter the conclusions of the existing cumulative assessment, and highly likely to reduce the cumulative impact assessed on the larger population present over a larger spatial scale.

Table 12.45 List of Other EO Member States Retained in the Transboundary impact Assessme	ŧητ
in Relation to the Topic	

EU member state	Commentary
Netherlands	<ul> <li>Rijkswaterstaat, Ministry of Infrastructure and Water Management, responded to the transboundary consultation requesting a meeting in relation to a number of issues including ornithology. A teleconference was held on 10 September 2018. Rijkswaterstaat provided details of the following offshore windfarm proposals. It is understood that these developments have been consented but not yet constructed and are due to become operational in the period 2019-2025</li> <li>Borssele 1,2, 0.7GW</li> <li>Borssele 3,4, 0.7GW</li> </ul>
	<ul> <li>Hollandse Kust (HK) zuid 1,2, 0.7GW</li> <li>HK zuid 3,4, 0.7GW</li> </ul>
	HK noord 1,2 0.7GW
	HK west, 1.4GW
	At the time of writing no specific information has been provided, or appears to be available (based on internet searches) in relation to turbine numbers and specifications, or ornithology assessments (and specifically estimates of collision risk and displacement for bird species) for individual developments.



## 12.9Interactions

330. The impacts identified and assessed in this chapter have the potential to interact with each other, which could give rise to synergistic impacts as a result of that interaction. The worst case impacts assessed within the chapter take these interactions into account and therefore the impact assessments are considered conservative and robust. For clarity, the areas of interaction between impacts are presented in *Table 12.46.* 

Potential interaction between impacts							
Construction	1 Disturbance and displace from increased vessel activ	ment ity	2 Indirect effects as a result of displacement of prey species due to increased noise and disturbance to sea bed				
1 Disturbance and displacement from increased vessel activity			Yes possible medium to long term effects on birds, but spatial magnitude very small				
2 Indirect effects as a result of displacement of prey species due to increased noise and disturbance to sea bed	Yes possible medium to lor effects on birds, but spatial magnitude very small	ng term					
Operation	3 Disturbance and displacement from offshore infrastructure and operational activity	4 Collision risk		5 Indirect impacts through effects on habitats and prey species			
3 Disturbance and displacement from offshore infrastructure and operational activity		No (birds that are displaced would not be at risk of collision)		No (direct displacement of birds overrides prey effects)			
4 Collision risk	No (birds that are displaced would not be at risk of collision)			No			
5 Indirect impacts through effects on habitats and prey species	No (mutually exclusive)	No					
Decommissioning				·			
It is anticipated that the decommissioning impacts will be similar in nature to those of construction.							

#### Table 12.46 Potential for Interactions Between Ornithology Impacts



# **12.10Inter-relationships**

- 331. The construction, operation and decommissioning phases of the proposed East Anglia TWO project would cause a range of effects on offshore ornithological interests. The magnitude of these effects has been assessed individually above in **section 12.6** using expert knowledge and judgement, drawing from a wide science base that includes project-specific surveys and previously acquired knowledge of the bird ecology of the North Sea (from published scientific papers and books, and 'grey' literature).
- 332. Impacts to offshore ornithological interests may be inter-related with other receptor groups. With respect to the impacts assessed for offshore ornithology (*section 12.6*), this is considered to be the case for indirect impacts through effects on habitats and prey species only. For direct disturbance/displacement and collision risk there is considered to be no potential for interaction with other receptor groups.
- 333. Inter-relationships are summarised in *Table 12.47*, which indicates where assessments carried out in other ES chapters have been used to inform the offshore ornithology assessment.

Impact	Related Chapter	Where addressed in this Chapter	Rationale
2 Indirect impacts through effects on habitats and prey during construction	Chapter 9 Benthic Ecology	Section 12.6.1.2	Potential impacts on benthic ecology and fish and shellfish during construction could affect the prey resource for birds.
	Chapter 10 Fish and Shellfish Ecology		
5 Indirect impacts through effects on habitats and prey during operation	Chapter 9 Benthic Ecology	Section 12.6.2.2	Potential impacts on benthic ecology and fish and shellfish during operation could affect the prey resource for birds.
	Chapter 10 Fish and Shellfish Ecology		
7 Indirect impacts through effects on habitats and prey during decommissioning	Chapter 9 Benthic Ecology Chapter 10 Fish and Shellfish Ecology	Section 12.6.3.2	Potential impacts on benthic ecology and fish and shellfish during decommissioning could affect the prey resource for birds.

## Table 12.47 Ornithology Inter-relationships



# 12.11Summary

- This chapter provides an assessment of the potential impacts on offshore 334. ornithology that may arise from the construction. operation and decommissioning of the offshore components (offshore windfarm site and export cable corridor to MLWS at the landfall site). It describes the offshore components of the proposed project; the consultation that has been held with stakeholders; the scope and methodology of the assessment; the avoidance and mitigation measures that have been embedded through project design; the baseline data on birds and important sites and habitats for birds acquired through desk study and survey (Appendix 12.1) and assesses the potential impacts on birds.
- 335. Detailed consultation and iteration of the overall approach to the impact assessment on ornithology receptors has informed this assessment through the ornithology ETG for East Anglia TWO, which involved Natural England and the RSPB.
- 336. A standard survey area for offshore ornithology, covering the East Anglia TWO offshore windfarm site and a 4km buffer was surveyed using high resolution aerial survey methods over periods of 24 months. The results of these surveys have been used to estimate the abundance and assemblage of birds using or passing across the area. The analysis and assessment in this PEIR has been undertaken prior to the data from the final aerial surveys being available, so is based on the first 21 monthly surveys and will be updated when 24 surveys are available.
- 337. The impacts that could potentially arise for offshore ornithology during the construction, operation and decommissioning of the proposed East Anglia TWO project were discussed with Natural England and the RSPB as part of the Evidence Plan process. As a result of those discussions it was agreed that the potential impacts that required detailed assessment were:
  - In the construction phase:
    - Impact 1: Disturbance/displacement
    - Impact 2: Indirect impacts through effects on habitats and prey species
  - In the operational phase:
    - Impact 3: Disturbance and displacement from offshore infrastructure and due to increased vessel and helicopter activity
    - o Impact 4: Collision risk
    - Impact 5: Indirect impacts through effects on habitats and prey species
  - In the decommissioning phase:
    - Impact 6: Disturbance/displacement



- Impact 7: Indirect impacts through effects on habitats and prey species
- 338. During the construction phase of the proposed project no impacts have been assessed to be greater than of minor adverse significance for any bird species.
- 339. During operation, displacement effects on red-throated divers, gannets, razorbills and guillemots would not create impacts of more than minor adverse significance during any biological season. The risk to birds from collisions with wind turbines from the proposed East Anglia TWO project alone is assessed as no greater than minor adverse significance for gannet, kittiwake, lesser black-backed gull and great black-backed gull when considered for all biological seasons against the most appropriate population scale.
- 340. Two potential effects of the proposed East Anglia TWO project were screened in for cumulative assessment: operational displacement and collision risk. Other potential effects would be temporary, small scale and localised and given the distances to other activities in the region (e.g. other offshore windfarms and aggregate extraction) it was concluded that there is no pathway for cumulative interaction.
- 341. A screening process was also carried out for potential plans and projects that might affect ornithological receptors cumulatively with the proposed project. In the offshore environment only other UK windfarms that were operational, under construction, consented but not constructed, subject to current applications or subject to consultation were screened in. This list of windfarms with their status is provided in *Table 12.35*.
- 342. The risk to ornithological receptors from cumulative displacement and collisions is assessed as no greater than minor adverse significance for all species.
- 343. The potential for collisions and displacement from windfarms outside UK territorial waters (transboundary) to contribute to cumulative impacts was considered. However, the operational offshore windfarms which might contribute to cumulative effects are comparatively small (in combination these projects are of a similar size to no more than one to two of the more recent UK windfarms, such as East Anglia TWO). Since the spatial scale and hence seabird population sizes for a transboundary assessment would be much larger, therefore, the inclusion of non-UK windfarms is considered very unlikely to alter the conclusions of the existing cumulative assessment, and highly likely to reduce the cumulative impact assessed on the larger population present over a larger spatial scale.
- 344. The identified impacts for the project alone are summarised in *Table 12.48* and cumulative impacts in *Table 12.49*.



Potential Impact	Receptor	Value/ Sensitivity	Magnitude	Significance	Mitigation	Residual Impact
Construction						
Direct disturbance and displacement during export cable construction	Red-throated diver	High	Negligible	Minor adverse	N/A	Minor adverse
Direct disturbance	Razorbill	Medium	Negligible	Negligible	N/A	Negligible
and displacement from construction activity on windfarm site	Guillemot	Medium	Negligible	Negligible	N/A	Negligible
Indirect effects due to prey species displacement	All species	Low to high	Negligible	Negligible to minor adverse	N/A	Negligible to minor adverse
Operation						
Direct disturbance and	Red-throated diver	High	Negligible	Minor adverse	N/A	Minor adverse
displacement	Gannet	Low to medium	Negligible	Negligible	N/A	Negligible
	Razorbill	Medium	Negligible	Negligible	N/A	Negligible
	Guillemot	Medium	Negligible	Negligible	N/A	Negligible
Indirect effects due to prey species displacement	All species	Low to high	Negligible	Negligible to minor adverse	N/A	Negligible to minor adverse
Collision risk	Gannet –	Low to medium	Negligible	Negligible	N/A	Negligible to minor adverse
	Kittiwake	Low to medium	Negligible	Negligible	N/A	Negligible to minor adverse

#### Table 12.48 Potential Impacts Identified for Offshore Ornithology



Potential Impact	Receptor	Value/ Sensitivity	Magnitude	Significance	Mitigation	Residual Impact	
	Lesser black- backed gull	Low to medium	Negligible	Negligible	N/A	Negligible to minor adverse	
	Great black- backed gull	Low to medium	Negligible	Negligible	N/A	Negligible to minor adverse	
Decommissioning							
Direct disturbance and displacement	All species	Low to high	Negligible	Negligible to minor adverse	N/A	Negligible to minor adverse	
Indirect effects due to prey species displacement	All species	Low to high	Negligible	Negligible to minor adverse	N/A	Negligible to minor adverse	

## Table 12.49 Potential Cumulative Impacts Identified for Ornithology

Potential Impact	Receptor	Value/ Sensitivity	Magnitude	Significance	Mitigation	Residual Impact
Operation						
Disturbance and displacement	Red- throated diver	High	Negligible	Minor adverse	N/A	Minor adverse
	Gannet	Low to medium	Negligible	Negligible	N/A	Negligible
	Razorbill	Low to medium	Negligible	Negligible	N/A	Negligible
	Guillemot	Low to medium	Negligible	Negligible	N/A	Negligible
Collision risk	Gannet	Low to medium	Low	Minor adverse	N/A	Minor adverse
	Kittiwake	Low to medium	Low	Minor adverse	N/A	Minor adverse
	Lesser black- backed gull	Low to medium	Low	Minor adverse	N/A	Minor adverse



# East Anglia TWO Offshore Windfarm Preliminary Environmental Information Report

Potential Impact	Receptor	Value/ Sensitivity	Magnitude	Significance	Mitigation	Residual Impact
	Great black- backed gull	Low to medium	Low	Minor adverse	N/A	Minor adverse



## 12.12 References

APEM (2011). East Anglia Offshore Wind: Zonal Assessment Report.

APEM. (2014) Assessing Northern gannet avoidance of offshore wind farms. Report for East Anglia Offshore Wind Ltd.

APEM (2016). Assessment of Displacement Impacts of Offshore Windfarms and Other Human Activities on Red-throated Divers and Alcids. Natural England Commissioned Reports, Number 227.

Austin, G.E. and Rehfisch, M.M. (2005) Shifting nonbreeding distributions of migratory fauna in relation to climate change. Global Change Biology, 11, 31-38.

Balmer, D., Gillings, S., Caffrey, B., Swann, B., Downie, I. and Fuller, R. (2013) Bird Atlas 2007-11: The Breeding and Wintering Birds of Britain and Ireland. BTO Books, Thetford.

Band, W. (2012) Using a collision risk model to assess bird collision risks for offshore wind farms. The Crown Estate Strategic Ornithological Support Services (SOSS) report SOSS-02. SOSS Website. Original published Sept 2011, extended to deal with flight height distribution data March 2012.

Bellebaum, J., Diederichs, A., Kube, J., Schulz, A. and Nehls, G. (2006) Flucht- und Meidedistanzen überwinternder Seetaucher und Meeresenten gegenüber Schiffen auf See. Ornithologischer Rundbrief Mecklenburg-Vorpommern, 45, 86–90.

Bicknell, A.W.J., Oro, D., Camphuysen, C.J. and Votier, S.C. (2013) Potential consequences of discard reform for seabird communities. Journal of Applied Ecology, 50, 649-658.

BirdLife International. (2004) Birds in Europe: population estimates, trends and conservation status. (Birdlife Conservation Series No. 12). BirdLife, Cambridge.

Braasch, A., Michalik, A. and Todeskino, E. (2015) Assessing impacts of offshore wind farms on two highly pelagic seabird species. Poster presented at Conference on Wind farms and Wildlife in Berlin.

Bradbury G., Trinder M., Furness B., Banks A.N., Caldow R.W.G., et al. (2014) Mapping Seabird Sensitivity to Offshore Wind farms. PLoS ONE, 9(9), e106366. doi:10.1371/journal.pone. 0106366

Brooke, M.D., Bonnaud, E., Dilley, B.J., Flint, E.N., Holmes, N.D., Jones, H.P., Provost, P., Rocamora, G., Ryan, P.G., Surman, C. and Buxton, R.T. (2018) Seabird population changes following mammal eradications on islands. Animal Conservation, 21, 3-12.

Brown, A. and Grice, P. (2005) Birds in England. T and AD Poyser, London.



Burthe, S.J., Wanless, S., Newell, M.A., Butler, A. and Daunt, F. (2014) Assessing the vulnerability of the marine bird community in the western North Sea to climate change and other anthropogenic impacts. Marine Ecology Progress Series, 507, 277-295.

Camphuysen, C.J. (1995) Herring Gull (Larus argentatus) and Lesser Black-backed Gull (L. fuscus) feeding at fishing vessels in the breeding season: competitive scavenging versus efficient flying. Ardea, 83, 365-380.

Capuzzo, E., Lynam, C.P., Barry, J., Stephens, D., Forster, R.M., Greenwood, N., McQuatters-Gollop, A., Silva, T., Van Leeuwen, S.M. and Engelhard, G.H. (2018) A decline in primary production in the North Sea over 25 years, associated with reductions in zooplankton abundance and fish stock recruitment. Global Change Biology, 24, E352-E364.

Carroll, M.J., Butler, A., Owen, E., Ewing, S.R., Cole, T., Green, J.A., Soanes, L.M., Arnould, J.P.Y., Newton, S.F., Baer, J., Daunt, F., Wanless, S., Newell, M.A., Robertson, G.S., Mavor, R.A. and Bolton, M. (2015) Effects of sea temperature and stratification changes on seabird breeding success. Climate Research, 66, 75-89.

Carroll, M.J., Bolton, M., Owen, E., Anderson, G.Q.A., Mackley, E.K., Dunn, E.K. and Furness, R.W. (2017) Kittiwake breeding success in the southern North Sea correlates with prior sandeel fishing mortality. Aquatic Conservation – Marine and Freshwater Ecosystems, 27, 1164-1175.

CIEEM. (2018) Guidelines for Ecological Impact Assessment in the UK and Ireland: Terrestrial, Freshwater, Coastal and Marine. Chartered Institute of Ecology and Environmental Management, Winchester. https://www.cieem.net/data/files/ECIA%20Guidelines.pdf

Cook, A.S.C.P., Wright, L.J., and Burton, N.H.K. (2012) A review of flight heights and avoidance rates of birds in relation to offshore wind farms. The Crown Estate Strategic Ornithological Support Services (SOSS). SOSS Website.

Cook, A.S.C.P., Humphries, E.M., Masden, E.A., and Burton, N.H.K. (2014) The avoidance rates of collision between birds and offshore turbines. BTO research Report No 656 to Marine Scotland Science. BTO, Thetford.

Cook, A,S.C.P, Humphreys, E.M., Bennet, F., Masden, E.A., and Burton, N.H.K. (in press). Quantifying avian avoidance of offshore wind turbines: Current evidence and key knowledge gaps. Marine Environmental Research.

Cramp S. and Simmons K.E.L. (Eds.) (1977 - 1994). The Birds of the Western Palearctic. Oxford University Press, Oxford.

Cury, P., Boyd, I.L., Bonhommeau, S., Anker-Nilssen, T., Crawford, R.J.M., Furness, R.W., Mills, J.A., Murphy, E.J., Österblom, H., Paleczny, M., Piatt, J.F., Roux, J-P., Shannon, L. and Sydeman, W.J. (2011) Global seabird response to forage fish depletion – one-third for the birds. Science, 334, 1703-1706.



Daunt, F., Benvenuti, S., Harris, M.P., Dall'Antonia, L., Elston, D.A. and Wanless, S. (2002) Foraging strategies of the black-legged kittiwake *Rissa tridactyla* at a North Sea colony: evidence for a maximum foraging range. Marine Ecology Progress Series, 245, 239-247.

Department of Energy and Climate Change (DECC) (2009) Future leasing for offshore wind farms and licensing for offshore oil and gas and gas storage. Environmental Report January 2009. Appendix 3d. Water Environment

DECC (2014) Environmental Assessment Report Comprising: Habitats Regulations Assessment, Transboundary Considerations and Consideration of Greater Backblacked Gulls Available online at: <u>http://infrastructure.planningportal.gov.uk/wp-content/ipc/uploads/projects/EN010032/3.%20Post%20Decision%20Information/Decision/Rampion%20Environmental%20Assessment%20Report.pdf</u>

Del Hoyo, J., Elliott, A. and Sargatal, J. (Eds.) (1992 – 2011) Handbook of the Birds of the World. Lynx Editions, Madrid.

Dierschke, V., Furness, R.W. and Garthe, S. (2016) Seabirds and offshore wind farms in European waters: avoidance and attraction. Biological Conservation, 202, 59-68.

Dierschke, V., Furness, R.W., Gray, C.E., Petersen, I.K., Schmutz, J., Zydelis, R. and Daunt, F. (2017) Possible behavioural, energetic and demographic effects of displacement of red-throated divers. JNCC Report No. 605. JNCC, Peterborough.

Dooling, R.J. and Therrien, S.C. (2012) Hearing in birds: What changes from air to water. Advances in Experimental Medicine and Biology, 730, 77-82.

Drewitt, A.L. and Langston, R.H.W. (2006) Assessing the impacts of wind farms on birds. Ibis, 148 (Suppl. 1), 4-7.

East Anglia ONE North Ltd (2019). East Anglia ONE North Offshore Wind Farm Chapter 12 Offshore Ornithology. Preliminary Environmental Report Volume 1.

EATL (2015) East Anglia THREE Chapter 13 Offshore Ornithology. Vol 1 Ref 6.1.13. Available online at: <u>https://infrastructure.planninginspectorate.gov.uk/wp-content/ipc/uploads/projects/EN010056/EN010056-000418-6.1.13%20Volume%201%20Chapter%2013%20Offshore%20Ornithology.pdf</u>

EATL (2016) Revised CRM. Submitted for Deadline 5: Available online at: <u>https://infrastructure.planninginspectorate.gov.uk/wp-</u>

content/ipc/uploads/projects/EN010056/EN010056-001644-EA3%20-%20Revised%20CRM.pdf

Eaton, M.A., Aebischer, N.J., Brown, A.F., Hearn, R.D., Lock, L., Musgrove, A.J., Noble, D.G., Stroud, D.A. and Gregory, R.D. (2015) Birds of Conservation Concern 4: the population status of birds in the United Kingdom, Channel Islands and Isle of Man. British Birds, 108, 708–746.



EATL (2016a). Applicants Comments on Written Representations. Deadline 3. appendix 1 Great black-backed gull PVA. https://infrastructure.planninginspectorate.gov.uk/wp-content/ipc/uploads/projects/EN010056/EN010056-001424-East%20Anglia%20THREE%20Limited%202

Exo, K-M., Hüppop, O. and Garthe, S. (2003) Offshore wind farms and bird protection. Seevögel, 23, 83-95.

Foster, S. and Marrs, S. (2012) Seabirds In Scotland. Scottish Natural Heritage Trend Note No. 21.

Foster, S., Swann, R.L. and Furness, R.W. (2017) Can changes in fishery landings explain long-term population trends in gulls? Bird Study, 64, 90-97.

Fox, A.D. and 19 Others. (2016) Seeking explanations for recent changes in abundance of wintering Eurasian wigeon (Anas penelope) in Northwest Europe. Ornis Fennica, 93, 12-25.

Frederiksen, M., Wanless, S., Harris, M.P., Rothery, P. and Wilson, L.J. (2004) The role of industrial fisheries and oceanographic change in the decline of North Sea black-legged kittiwakes. Journal of Applied Ecology, 41, 1129-1139.

Frederiksen, M., Furness, R.W. and Wanless, S. (2007) Regional variation in the role of bottom-up and top-down processes in controlling sandeel abundance in the North Sea. Marine Ecology Progress Series, 337, 279-286.

Frederiksen, M., Daunt, F., Harris, M.P. and Wanless, S. (2008) The demographic impact of extreme events: Stochastic weather drives survival and population dynamics in a long-lived seabird. Journal of Animal Ecology, 77, 1020-1029.

Frederiksen, M., Anker-Nilssen, T., Beaugrand, G. and Wanless, S. (2013) Climate, copepods and seabirds in the Boreal Northeast Atlantic – Current state and future outlook. Global Change Biology, 19, 364-372.

Furness, R.W. (2015) Non-breeding season populations of seabirds in UK waters: Population sizes for Biologically Defined Minimum Population Scales (BDMPS). Natural England Commissioned Report Number 164.

Furness, R.W. (2016) Impacts and effects of ocean warming on seabirds. In: Laffoley, D. and Baxter, J.M. (Editors). Explaining Ocean Warming: Causes, Scale, Effects And Consequences. Full Report. Gland, Switzerland: IUCN. Pp. 271-288.

Furness, R.W. (2018). Consequences for birds of obstruction lighting on offshore wind turbines.



Furness, R.W. and Tasker, M.L. (2000) Seabird-fishery interactions: quantifying the sensitivity of seabirds to reductions in sandeel abundance and identification of key areas for sensitive seabirds in the North Sea. Marine Ecology Progress Series, 202, 253-264.

Furness, R.W., Wade, H.M. and Masden, E.A. (2013) Assessing vulnerability of marine bird populations to offshore wind farms. Journal of Environmental Management, 119, 56-66.

Furness, R.W., Hallgrimsson, G.T., Montevecchi, W.A., Fifield, D., Kubetzki, U., Mendel, B. and Garthe, S. (2018a) Adult gannet migrations frequently loop clockwise around Britain and Ireland. Ringing and Migration, doi: 10.1080/03078698.2018.1472971.

Furness, R.W., Garthe, S., Trinder, M., Matthiopoulos, J., Wanless, S. and Jeglinski, J. (2018b) Nocturnal flight activity of northern gannets *Morus bassanus* and implications for modelling collision risk at offshore wind farms. Environmental Impact Assessment Review, 73, 73, 1-6. <u>https://www.sciencedirect.com/science/article/abs/pii/S019592551830091X</u>

Furness et al. (in prep). Nocturnal flight activity of black-legged kittiwakes, *Rissa tridactyla* and implications for modelling collision risk at offshore wind farms.

Garthe, S and Hüppop, O. (2004) Scaling possible adverse effects of marine wind farms on seabirds: developing and applying a vulnerability index. Journal of Applied Ecology, 41, 724-734.

Garthe, S., Grémillet, D. and Furness, R.W. (1999) At-sea-activity and foraging efficiency in chick-rearing northern gannets *Sula bassana*: a case study in Shetland. Marine Ecology Progress Series, 185, 93-99.

Garthe, S., Ludynia, K., Hüppop, O., Kubetzki, U., Meraz, J.F. and Furness, R.W. (2012) Energy budgets reveal equal benefits of varied migration strategies in northern gannets. Marine Biology, 159, 1907-1915.

GGOWL. (2011) Quarterly Ornithological Monitoring Report (Q3): December 2010-February 2011 for the Greater Gabbard Offshore Wind farm. Produced by ESS and Royal Haskoning on behalf of Greater Gabbard Offshore Wind Limited (GGOWL). April 2011.

Green, R.E., Langston, R.H.W., McCluskie, A., Sutherland, R. and Wilson, J.D. (2016) Lack of sound science in assessing wind farm impacts on seabirds. Journal of Applied Ecology, 53, 1635-1641.

Hamer, K.C., Monaghan, P., Uttley, J.D., Walton, P. and Burns, M.D. (1993) The influence of food supply on the breeding ecology of kittiwakes *Rissa tridactyla* in Shetland. Ibis, 135, 255-263.



Harris, M. and Wanless, S. (1989) The breeding biology of razorbills Alca torda on the Isle of May. Bird Study, 36, 105-114.

Holling, M. and the Rare Breeding Birds Panel. (2011) Rare breeding birds in the United Kingdom in 2009. British Birds, 104, 476–537.

Holt, C.A., Austin, G.E., Calbrade, N.A., Mellan, H.J., Hearn, R.D., Stroud, D.A., Wotton, S.R. and Musgrove, A.J. (2012) Waterbirds in the UK 2010/11: The Wetland Bird Survey. BTO/RSPB/JNCC, Thetford.

Horswill, C. and Robinson R. A. (2015) Review of seabird demographic rates and density dependence. JNCC Report No. 552. Joint Nature Conservation Committee, Peterborough

Hüppop, O. and Wurm, S. (2000) Effect of winter fishery activities on resting numbers, food and body condition of large gulls *Larus argentatus* and *L. marinus* in the south-eastern North Sea. Marine Ecology Progress Series, 194, 241-247.

Jarrett, D., Cook, A. S. C. P., Woodward, I., Ross, K., Horswill, C., Dadam, D. & Humphreys, E.M. (2018). Short-Term Behavioural Responses of Wintering Waterbirds to Marine

Activity (CR/2015/17). Scottish Marine and Freshwater Science Vol 9, No 7.

JNCC. (2013) The mean surface density map data was from NE's aerial visual surveys undertaken during winter (October – March) between 2001/01 and 2009/10 (GIS data files supplied by JNCC attached to e-mail dated 7th May 2013)

JNCC. (2015) Seabird Displacement Impacts from Offshore Wind Farms: report of the MROG Workshop, 6-7th May 2015. JNCC Report No 568. JNCC Peterborough.

JNCC. (2016) Seabird Population Trends And Causes Of Change: 1986-2015 Report Http://Jncc.Defra.Gov.Uk/Page-3201 Joint Nature Conservation Committee, Peterborough.

JNCC, NE, NIEA, NRW and SNH (2014) Joint Response from the Statutory Nature Conservation Bodies to the Marine Scotland Science Avoidance Rate Review.

Johnston, A., Cook, A.S.C.P., Wright, L.J., Humphreys, E.M. and Burton, E.H.K. (2014a) Modelling flight heights of marine birds to more accurately assess collision risk with offshore wind turbines. Journal of Applied Ecology, 51, 31-41.

Johnston, A., Cook, A.S.C.P., Wright, L.J., Humphreys, E.M. and Burton, N.H.K. (2014b) corrigendum. Journal of Applied Ecology, 51, doi: 10.1111/1365-2664.12260.

King, S., Maclean, I.M.D., Norman, T., and Prior, A. (2009) Developing Guidance on Ornithological Cumulative Impact Assessment for Offshore Wind Farm Developers. COWRIE.



Kober, K., Webb, A., Win, I., Lewis, M., O'Brien, S., Wilson, L.J. and Reid, J.B. (2010) An analysis of the numbers and distribution of seabirds within the British Fishery Limit aimed at identifying areas that qualify as possible marine SPAs. JNCC Report, No. 431. JNCC, Peterborough.

Kotzerka, J., Garthe, S. and Hatch, S.A. (2010) GPS tracking devices reveal foraging strategies of black-legged kittiwakes. Journal of Ornithology, 151, 495-467.

Krijgsveld, K. L., Fijn, R. C., Japink, M., van Horssen, P. W., Heunks, C., Collier, M. P., Poot, M. J. M., Beuker, D. and Dirksen, S. (2011) Effect Studies Offshore Wind farm Egmond aan Zee. Final report on fluxes, flight altitudes and behaviour of flying birds. Bureau Waardenburg report 10-219, NZW-Report Bureau Waardenburg, Culmeborg, Netherlands.

Kubetzki, U., Garthe, S., Fifield, D., Mendel, B. and Furness, R.W. (2009) Individual migratory schedules and wintering areas of northern gannets. Marine Ecology Progress Series, 391, 257-265.

Langston, R.H.W. (2010) Offshore wind farms and birds: Round 3 zones, extensions to Round 1 and Round 2 sites and Scottish Territorial Waters. RSPB Research Report No. 39. RSPB, Sandy.

Langston, R.H.W., Teuten, E. and Butler, A. (2013) Foraging ranges of northern gannets *Morus bassanus* in relation to proposed offshore wind farms in the UK: 2010-2012. Report to DECC. Reference DECC URN:13D/306.

Leopold, M.F. and Camphuysen, C.J. (2007) Did the pile driving during the construction of the Offshore Wind farm Egmond aan Zee, the Netherlands, impact local seabirds? Report CO62/07. Wageningen IMARES Institute for Marine Resources and Ecosystem Studies.

Leopold, M.F., Dijkman, E.M. and Teal, L. (2011) Local birds in and around the Offshore Wind farm Egmond aan Zee (OWEZ) (T-0 and T-1, 2002-2010). NoordzeeWind report OWEZ\_R\_221\_T1\_20110915\_localbirds\_final. Imares / NoordzeeWind, Wageningen /IJmuiden.

Lindgren, M., van Duers, M., Mackenzie, B.R., Clausen, L.W., Christensen, A. and Rindorf, A. (2018) Productivity and recovery of forage fish under climate change and fishing: North Sea sandeel as a case study. Fisheries Oceanography, 27, 212-221.

MacArthur Green (2017) Estimates of Ornithological Headroom in Offshore Wind Farm Collision Mortality. Report to The Crown Estate.

Macdonald, A., Heath, M.R., Edwards, M., Furness, R.W., Pinnegar, J.K., Wanless, S., Speirs, D.C. and Greenstreet, S.P.R. (2015) Climate driven trophic cascades affecting seabirds around the British Isles. Oceanography and Marine Biology, 53, 55-80.



Maclean, I.M.D., Wright, L.J., Showler, D.A. and Rehfisch, M.M. (2009) A Review of Assessment Methodologies for Offshore Wind Farms. BTO report commissioned by COWRIE Ltd.

Masden, E.A., Haydon, D.T., Fox, A.D. and Furness, R.W. (2010) Barriers to movement: Modelling energetic costs of avoiding marine wind farms amongst breeding seabirds. Marine Pollution Bulletin, 60, 1085-1091.

Masden E.A., Reeve, R., Desholm, M., Fox, A.D., Furness, R.W. and Haydon, D.T. (2012) Assessing the impact of marine wind farms on birds through movement modelling. Journal of the Royal Society Interface, 9, 2120-2130.

Mendel, B., Sonntag, N., Wahl, J., Schwemmer, P., Dries, H., Guse, N., Müller, S. and Garthe, S. (2008) Profiles of seabirds and waterbirds of the German North and Baltic Seas: Distribution, ecology and sensitivities to human activities within the marine environment. Bundesamt Für Naturschutz, Bonn – Bad Godesberg.

Mendel, B., Kotzerka, J., Sommerfeld, J., Schwemmer, H., Sonntag, N., and Garthe, S. (2014) Effects of the alpha ventus offshore test site on distribution patterns, behaviour and flight heights of seabirds. In Ecological Research at the Offshore Wind farm alpha ventus pp. 95-110. Springer Fachmedien, Wiesbaden.

Mendel, B, Schwemmer, P., Peschko, V., Müller, S., Schwemmer, H., Mercker, M. & Garther, S. (2019) Operational offshore wind farms and associated ship traffic cause profound changes in distribution patterns of Loons (Gavia spp.). Journal of Environmental Management 231:429-438. https://doi.org/10.1016/j.jenvman.2018.10.053

Mitchell, P.I., Newton, S.F., Ratcliffe, N. and Dunn, T.E. (2004) Seabird Populations of Britain and Ireland. T. and A.D. Poyser, London.

Mitschke, A., Garthe, S. and Hüppop, O. (2001) Erfassung der Verbreitung, Häufigkeiten und Wanderungen von See- und Wasservögeln in der deutschen Nordsee und Entwicklung eines Konzeptes zur Umsetzung internationaler Naturschutzziele. BfN-Skripten 34, Bonn–Bad Godesberg.

MMO (2018) Displacement and habituation of seabirds in response to marine activities. A report produced for the Marine Management Organisation. MMO Project No: 1139, May 2018.

Murray, S., Harris, M.P. and Wanless, S. (2015) The status of the gannet in Scotland in 2013-14. Scottish Birds, 35, 3-18.

Musgrove, A.J., Aebischer, N.J., Eaton, M.A., Hearn, R.D., Newson, S.E., Noble, D.G., Parsons, M., Risely, K. and Stroud, D.A. (2013) Population estimates on birds in Great Britain and the United Kingdom. British Birds, 106, 64–100.



Natural England (2012) TIN131. Black-throated diver: species information for marine Special Protection Area consultations.

Norfolk Boreas Ltd (2018). Norfolk Boreas Offshore Wind Farm Chapter 13 Offshore Ornithology. Preliminary Environmental Report Volume 1. Document Reference PB5640-005-13.

Norfolk Vanguard Ltd (2018). Norfolk Vanguard Offshore Wind Farm Chapter 13 Offshore Ornithology. Environmental Statement Volume 1. Document Reference 6.1.13.

Ost, M., Lehikoinen, A., Poysa, H. and Linden, A. (2016) European ducks in a changing world: Human impacts, population processes and species interactions. Ornis Fennica, 93, 1-2.

Pearce-Higgins, J.W. and Holt, C.A. (2013) Impacts of climate change on waterbirds. Marine Climate Change Impacts Partnership Science Review, 2013, 149-154.

Pennycuick, C.J. (1987) Flight of auks (Alcidae) and other northern seabirds compared with southern Procellariiformes: ornithodolite observations. Journal of Experimental Biology, 128, 335-347.

Percival, S. (2010) Kentish Flats Offshore Wind farm: Diver Surveys 2009-10. Report to Vattenfall. Ecology Consulting, Durham.

Percival, S. (2013) Thanet Offshore Wind Farm. Ornithological Monitoring 2012-2013 Final Report. (http://ecologyconsult.co.uk/index\_htm\_files/Ornithological%20monitoring%202012-13%20v3.pdf)

Percival, S.M. (2014) Kentish Flats Offshore Wind farm: Diver Surveys 2011-12 and 2012-13. Ecology Consulting report to Vattenfall.

Petersen, I.K., Christensen, T.K., Kahlert, J., Desholm, M. and Fox, A.D. (2006) Final results of bird studies at the offshore wind farms at Nysted and Horns Rev, Denmark. NERI report commissioned by DONG energy and Vattenfall A/S 2006.

Piotrowski (2012)Lesser black-backed gull and herring gull breeding colonies in<br/>Suffolk.Suffolk.SuffolkBirds,62,23-30.https://issuu.com/suffolknaturalistssociety/docs/sb62a, accessed July 2018.

Planning inspectorate (2015) Advice note seventeen: Cumulative Effects Assessment relevant to nationally significant infrastructure projects. https://infrastructure.planninginspectorate.gov.uk/wp-content/uploads/2015/12/Advice-note-17V4.pdf, accessed October 2018.

Planning Inspectorate (2018) Advice note nine: Using the Rochdale Envelope, Version3.PlanningInspectorate,Bristol.



https://infrastructure.planninginspectorate.gov.uk/wp-content/uploads/2013/05/Advicenote-9.-Rochdale-envelope-web.pdf, accessed October 2018.

Planning Inspectorate (2014) Rampion Offshore Wind Farm and connection works Examining Authority's Report of Findings and Conclusions and Recommendation to the Secretary of State for Energy and Climate Change. Available online at: <u>http://infrastructure.planningportal.gov.uk/wp-</u>

content/ipc/uploads/projects/EN010032/3.%20Post%20Decision%20Information/Decision/Rampion%20Recommendation%20Report.pdf

R Core Team (2016) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. <u>www.R-project.org</u>

Ratcliffe, N. (2004) Causes of seabird population change. Pp 407-437 In Mitchell, P.I., Newton, S.F., Ratcliffe, N. And Dunn, T.E. (Eds.) Seabird Populations of Britain and Ireland. T.and A.D. Poyser, London.

Ratcliffe, N., Mitchell, I., Varnham, K., Verboven, N. and Higson, P. (2009) How to prioritize rat management for the benefit of petrels: a case study of the UK, Channel Islands and Isle of Man. Ibis, 151, 699-708.

RenewableUK. (2013) Cumulative Impact Assessment Guidelines – Guiding Principles for Cumulative Impacts Assessment in Offshore Wind farms. RenewableUK, London.

Robinson, R.A. (2005) Bird Facts: profiles of birds occurring in Britain and Ireland. BTO Research Report 407, BTO, Thetford.

Rock, P. and Vaughan, I.P. (2013) Long-term estimates of adult survival rates of urban herring gulls *Larus argentatus* and lesser black-backed gulls *Larus fuscus*. Ringing and Migration, 28, 21-29.

Sandvik, H., Erikstad, K.E. and Saether, B.-E. (2012) Climate affects seabird population dynamics both via reproduction and adult survival. Marine Ecology Progress Series, 454, 273-284.

Schwemmer, P. Mendel, B., Sonntag, N., Dierschke, V. and Garthe, S. (2011) Effects of ship traffic on seabirds in offshore waters: implications for marine conservation and spatial planning. Ecological Applications, 21, 1851-1860.

Scottish Power Renewables (2017). East Anglia TWO Offshore Windfarm Scoping report.

Searle, K., Mobbs, D., Butler, A., Bogdanova, M., Freeman, S., Wanless, S. and Daunt, F. (2014) Population consequences of displacement from proposed offshore wind energy developments for seabirds breeding at Scottish SPAs (CR/2012/03). Report to Marine Scotland Science. Centre for Ecology and Hydrology, Penicuik.



Searle, K.R., Mobbs, D.C., Butler, A., Furness, R.W., Trinder, M.N. and Daunt, F. (2017) Fate of displaced birds. Report to Marine Scotland Science. Centre for Ecology and Hydrology, Penicuik.

Skov, H., Heinänen, S., Norman, T., Ward, R.M., Méndez-Roldán, S. and Ellis, I. (2018) ORJIP Bird Collision and Avoidance Study. Final report – April 2018. The Carbon Trust, United Kingdom.

SNCB (2017) Joint SNCB Interim Displacement Advice Note.

Speakman, J., Gray, H. and Furness, L. (2009) University of Aberdeen report on effects of offshore wind farms on the energy demands of seabirds. Report to the Department of Energy and Climate Change.

Stienen, E.W., Waeyenberge, V., Kuijken, E. and Seys, J. (2007) Trapped within the corridor of the southern North Sea: the potential impact of offshore wind farms on seabirds. In Birds and Wind farms. de Lucas, M., Janss, G.F.E. and Ferrer, M. (Eds). Quercus, Madrid.

Stone, C.J. Webb, A., Barton, C., Ratcliffe, N., Reed, T.C. Tasker, M.L. Camphuysen, C.J. and Pienkowski, M.W. (1995) An atlas of seabird distribution in north-west European waters. JNCC, Peterborough.

Sydeman, W.J., Thompson, S.A., Anker-Nilssen, T., Arimitsu, M., Bennison, A., Bertrand, S., Boersch-Supan, P., Boyd, C., Bransome, N., Crawford, R.J.M., Daunt, F., Furness, R.W., Gianuca, D., Gladics, A., Koehn, L., Lang, J., Logerwell, E., Morris, T.L., Phillips, E.M., Provencher, J., Punt, A.E., Saraux, C., Shannon, L., Sherley, R.B., Simeone, A., Wanless, R.M., Wanless, S. and Zador, S. (2017) Best practices for assessing forage fish fisheries – seabird resource competition. Fisheries Research, 194, 209-221.

Tasker, M.L., Camphuysen, C.J., Cooper, J., Garthe, S., Montevecchi, W.A. and Blaber, S.J.M. (2000) The impacts of fishing on marine birds. ICES Journal of Marine Science, 57, 531-547.

Thaxter, C.B., Lascelles, B., Sugar, K., Cook, A.S.C.P., Roos, S., Bolton, M., Langston, R.H.W. and Burton, N.H.K. (2012) Seabird foraging ranges as a preliminary tool for identifying Marine Protected Areas. Biological Conservation, 156, 53-61.

Thaxter, C.B., Ross-Smith, V.H., Bouten, W., Masden, E.A., Clark, N.A., Conway, G.J., Barber, L., Clewley, G.D. and Burton, N.H.K. (2018) Dodging the blades: new insights into three-dimensional space use of offshore wind farms by lesser black-backed gulls *Larus fuscus*. Marine Ecology Progress series, 587, 247-253.

Thaxter, C.B, Ross- Smith, V.H., Bouten, W., Clark, N.A., Conway, G.J., Rehfisch, M.M., Burton, N.H.K. 2015. Seabird–wind farm interactions during the breeding season vary within and between years: A case study of lesser black-backed gull Larus fuscus in the UK. Biological conservation 186:347-358



Topping, C. and Peterson, I.K. (2011) Report on a red-throated diver agent-based model to assess the cumulative impact from offshore wind farms. Report commissioned by the Environment Group. Aarhus University. Danish Centre for Environment and Energy.

Trinder, M. (2014) Flamborough and Filey Coast pSPA Seabird PVA Final Report, submitted for Hornsea Wind Farm Project ONE, Appendix N, Deadline V, 14 May 2014.

Uttley, J.D., Walton, P., Monaghan, P. and Austin, G. (1994) The effects of food abundance on breeding performance and adult time budgets of guillemots *Uria aalge*. Ibis, 136, 205–213.

Vallejo, G.C., Grellier, K., Nelson, E.J., McGregor, R.M., Canning, S.J., Caryl, F.M. and McLean, N. (2017) Responses of two marine top predators to an offshore wind farm. Ecology and Evolution, 7, 8698-8708.

Vanermen, N., Stienen, E.W.M., Onkelinx, T., Courtens, W., Van De Walle, M., Verschelde, P. and Verstraete, H. (2012) Seabirds and Offshore Wind Farms Monitoring Results 2011. Research Institute for Nature and Forest: Study commissioned by the Royal Belgian Institute for Natural Sciences, Management Unit of the North Sea Mathematical Models.

Vanermen, N., Stienen, E.W.M., Courtens, W., Onkelinx, T., Van de walle, M. and Verstraete, H. (2013) Bird monitoring at offshore wind farms in the Belgian part of the North Sea - Assessing seabird displacement effects. Rapporten van het Instituut voor Natuur- en Bosonderzoek 2013 (INBO.R.2013.755887). Instituut voor Natuur- en Bosonderzoek, Brussel.

Vanermen, N, Onkelinx, T, Courtens, W., Van de walle, M., Hilbran Verstraete, and Stienen E.W.M. (2014) Seabird avoidance and attraction at an offshore wind farm in the Belgian part of the North Sea. Hydrobiologia. DOI 10.1007/s10750-014-2088-x

Votier, S.C., Furness, R.W., Bearhop, S., Crane, J.E., Caldow, R.W.G., Catry, P., Ensor, K., Hamer, K.C., Hudson, A.V., Kalmbach, E., Klomp, N.I., Pfeiffer, S., Phillips, R.A., Prieto, I., and Thompson, D.R. (2004) Changes in fisheries discard rates and seabird communities. Nature, 427, 727-730.

Votier, S.C., Hatchwell, B.J., Beckerman, A., Mccleery, R.H., Hunter, F.M., Pellatt, J., Trinder, M. and Birkhead, T.R. (2005) Oil pollution and climate have wide-scale impacts on seabird demographics. Ecology Letters, 8, 1157-1164.

Votier, S.C., Birkhead, T.R., Oro, D., Trinder, M., Grantham, M.J., Clark, J.A., Mccleery, R.H. and Hatchwell, B.J. (2008) Recruitment and survival of immature seabirds in relation to oil spills and climate variability. Journal of Animal Ecology, 77, 974-983.


Votier, S.C., Bicknell, A., Cox, S.L., Scales, K.L. and Patrick, S.C. (2013) A bird's eye view of discard reforms: Bird-borne cameras reveal seabird/fishery interactions. Plos One, 8(3), E57376.

Wade, H.M., Masden E.M., Jackson, A.C. and Furness, R.W. (2016) Incorporating data uncertainty when estimating potential vulnerability of Scottish seabirds to marine renewable energy developments. Marine Policy, 70, 108-113.

Walls, R., Canning, S., Lye, G., Givens, L., Garrett, C. and Lancaster, J. (2013) Analysis of Marine Environmental Monitoring Plan Data from the Robin Rigg Offshore Wind farm, Scotland (Operational Year 1). Natural Power report for E.ON Climate and Renewables UK.

Wernham, C.V., Toms, M.P., Marchant, J.H., Clark, J.A., Siriwardena, G.M. and Baillie, S.R. (eds). (2002) The Migration Atlas: Movements of the birds of Britain and Ireland. T. and A.D. Poyser, London.

Wright, L.J., Ross-Smith, V.H., Massimino, D., Dadam, D., Cook, A.S.C.P. and Burton, N.H.K. (2012) Assessing the risk of offshore windfarm development to migratory birds designated as features of UK Special Protection Areas (and other Annex I species). Strategic Ornithological Support Services. Project SOSS-05. BTO Research Report No. 592.

WWT (2012) SOSS-04 Gannet PVA – Demographic data, population model and outputs. Available from <u>www.bto.org/science/wetland-and-marine/soss/projects</u>



This page is intentionally blank.